

1.8 Determining Limits Using the Squeeze Theorem

Name: _____ Class: _____ Date: _____

Total: 9 marks

KEY WORDS squeeze theorem 夹逼定理 • limit 极限

Objective

Build the skills to answer exam questions on **the Squeeze Theorem**.

You must be able to:

- state the **Squeeze Theorem** 夹逼定理
- apply it when a function is bounded between two others with equal limits

1 Worked examples

Study these first. Each one shows the method for a question type used later.

■ The Squeeze Theorem

If $g(x) \leq f(x) \leq h(x)$ near a and

$$\lim_{x \rightarrow a} g(x) = \lim_{x \rightarrow a} h(x) = L,$$

then $\lim_{x \rightarrow a} f(x) = L$.

■ Example

Since $-x^2 \leq x^2 \sin \frac{1}{x} \leq x^2$ and both bounds $\rightarrow 0$ at 0, the middle limit is 0.

2 Practice

2.1 State the Squeeze Theorem.

[1]

2.2 State what must be true of the two bounding limits.

[1]

2.3 If $-x^2 \leq f(x) \leq x^2$ and both bounds $\rightarrow 0$ at 0, state $\lim_{x \rightarrow 0} f(x)$. [2]

3 Exam-style questions

3.1 The Squeeze Theorem requires $g \leq f \leq h$ and [1]

A	B
C	D

A $\lim g = \lim h$

B $g = h$ everywhere

C $f = 0$

D $h = 0$

3.2 If both bounds approach 3, then $\lim f$ is [1]

A	B
C	D

A 0

B 3

C 6

D undefined

3.3 $-x^2 \leq f(x) \leq x^2$ near 0.

(a) State $\lim_{x \rightarrow 0} (-x^2)$. [1]

(b) State $\lim_{x \rightarrow 0} x^2$. [1]

(c) State $\lim_{x \rightarrow 0} f(x)$. [1]