


CALCULUS AB
SECTION II, Part A
Time—30 minutes
Number of problems—2

A graphing calculator is required for these problems.

1. Grass clippings are placed in a bin, where they decompose. For $0 \leq t \leq 30$, the amount of grass clippings remaining in the bin is modeled by $A(t) = 6.687(0.931)^t$, where $A(t)$ is measured in pounds and t is measured in days.
- (a) Find the average rate of change of $A(t)$ over the interval $0 \leq t \leq 30$. Indicate units of measure.
- (b) Find the value of $A'(15)$. Using correct units, interpret the meaning of the value in the context of the problem.
- (c) Find the time t for which the amount of grass clippings in the bin is equal to the average amount of grass clippings in the bin over the interval $0 \leq t \leq 30$.
- (d) For $t > 30$, $L(t)$, the linear approximation to A at $t = 30$, is a better model for the amount of grass clippings remaining in the bin. Use $L(t)$ to predict the time at which there will be 0.5 pound of grass clippings remaining in the bin. Show the work that leads to your answer.

t (minutes)	0	2	5	8	12
$v_A(t)$ (meters/minute)	0	100	40	-120	-150

4. Train A runs back and forth on an east-west section of railroad track. Train A 's velocity, measured in meters per minute, is given by a differentiable function $v_A(t)$, where time t is measured in minutes. Selected values for $v_A(t)$ are given in the table above.
- (a) Find the average acceleration of train A over the interval $2 \leq t \leq 8$.
- (b) Do the data in the table support the conclusion that train A 's velocity is -100 meters per minute at some time t with $5 < t < 8$? Give a reason for your answer.
- (c) At time $t = 2$, train A 's position is 300 meters east of the Origin Station, and the train is moving to the east. Write an expression involving an integral that gives the position of train A , in meters from the Origin Station, at time $t = 12$. Use a trapezoidal sum with three subintervals indicated by the table to approximate the position of the train at time $t = 12$.
- (d) A second train, train B , travels north from the Origin Station. At time t the velocity of train B is given by $v_B(t) = -5t^2 + 60t + 25$, and at time $t = 2$ the train is 400 meters north of the station. Find the rate, in meters per minute, at which the distance between train A and train B is changing at time $t = 2$.



2. Let f and g be the functions defined by $f(x) = 1 + x + e^{x^2-2x}$ and $g(x) = x^4 - 6.5x^2 + 6x + 2$. Let R and S be the two regions enclosed by the graphs of f and g shown in the figure above.
- (a) Find the sum of the areas of regions R and S .
- (b) Region S is the base of a solid whose cross sections perpendicular to the x -axis are squares. Find the volume of the solid.
- (c) Let h be the vertical distance between the graphs of f and g in region S . Find the rate at which h changes with respect to x when $x = 1.8$.

END OF PART A OF SECTION II


t (years)	2	3	5	7	10
$H(t)$ (meters)	1.5	2	6	11	15

4. The height of a tree at time t is given by a twice-differentiable function H , where $H(t)$ is measured in meters and t is measured in years. Selected values of $H(t)$ are given in the table above.

- (a) Use the data in the table to estimate $H'(6)$. Using correct units, interpret the meaning of $H'(6)$ in the context of the problem.
- (b) Explain why there must be at least one time t , for $2 < t < 10$, such that $H'(t) = 2$.
- (c) Use a trapezoidal sum with the four subintervals indicated by the data in the table to approximate the average height of the tree over the time interval $2 \leq t \leq 10$.
- (d) The height of the tree, in meters, can also be modeled by the function G , given by $G(x) = \frac{100x}{1+x}$, where x is the diameter of the base of the tree, in meters. When the tree is 50 meters tall, the diameter of the base of the tree is increasing at a rate of 0.03 meter per year. According to this model, what is the rate of change of the height of the tree with respect to time, in meters per year, at the time when the tree is 50 meters tall?

5. Let f be the function defined by $f(x) = e^x \cos x$.

- (a) Find the average rate of change of f on the interval $0 \leq x \leq \pi$.
- (b) What is the slope of the line tangent to the graph of f at $x = \frac{3\pi}{2}$?
- (c) Find the absolute minimum value of f on the interval $0 \leq x \leq 2\pi$. Justify your answer.
- (d) Let g be a differentiable function such that $g\left(\frac{\pi}{2}\right) = 0$. The graph of g' , the derivative of g , is shown below. Find the value of $\lim_{x \rightarrow \pi/2} \frac{f(x)}{g(x)}$ or state that it does not exist. Justify your answer.

Graph of g'

t (days)	0	3	7	10	12
$r'(t)$ (centimeters per day)	-6.1	-5.0	-4.4	-3.8	-3.5

4. An ice sculpture melts in such a way that it can be modeled as a cone that maintains a conical shape as it decreases in size. The radius of the base of the cone is given by a twice-differentiable function r , where $r(t)$ is measured in centimeters and t is measured in days. The table above gives selected values of $r'(t)$, the rate of change of the radius, over the time interval $0 \leq t \leq 12$.

(a) Approximate $r''(8.5)$ using the average rate of change of r' over the interval $7 \leq t \leq 10$. Show the computations that lead to your answer, and indicate units of measure.

(b) Is there a time t , $0 \leq t \leq 3$, for which $r'(t) = -6$? Justify your answer.

(c) Use a right Riemann sum with the four subintervals indicated in the table to approximate the value of

$$\int_0^{12} r'(t) \, dt.$$

(d) The height of the cone decreases at a rate of 2 centimeters per day. At time $t = 3$ days, the radius is 100 centimeters and the height is 50 centimeters. Find the rate of change of the volume of the cone with respect to time, in cubic centimeters per day, at time $t = 3$ days. (The volume V of a cone with radius r and height h is $V = \frac{1}{3} \pi r^2 h$.)

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

t (minutes)	0	3	7	12
$C(t)$ (degrees Celsius)	100	85	69	55

1. The temperature of coffee in a cup at time t minutes is modeled by a decreasing differentiable function C , where $C(t)$ is measured in degrees Celsius. For $0 \leq t \leq 12$, selected values of $C(t)$ are given in the table shown.

(a) Approximate $C'(5)$ using the average rate of change of C over the interval $3 \leq t \leq 7$. Show the work that leads to your answer and include units of measure.

(b) Use a left Riemann sum with the three subintervals indicated by the data in the table to approximate the value of $\int_0^{12} C(t) \, dt$. Interpret the meaning of $\frac{1}{12} \int_0^{12} C(t) \, dt$ in the context of the problem.

(c) For $12 \leq t \leq 20$, the rate of change of the temperature of the coffee is modeled by $C'(t) = \frac{-24.55e^{0.01t}}{t}$,

where $C'(t)$ is measured in degrees Celsius per minute. Find the temperature of the coffee at time $t = 20$.

Show the setup for your calculations.

(d) For the model defined in part (c), it can be shown that $C''(t) = \frac{0.2455e^{0.01t}(100 - t)}{t^2}$. For $12 < t < 20$,

determine whether the temperature of the coffee is changing at a decreasing rate or at an increasing rate.

Give a reason for your answer.

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.