

1. Male birds of a certain species arrive at a nesting area over a thirty-day period. The rate at which the male birds arrive at the nesting area at time t days is modeled by a differentiable function M , where $M(t)$ is measured in number of birds per day. Selected values of $M(t)$ are shown in the table.

t (days)	0	5	10	15	20	25	30
$M(t)$ (birds per day)	2	7	16	6	5	2	0

(Note: Your calculator should be in radian mode.)

Part A

Approximate $M'(7.5)$ using the average rate of change of M over the interval $5 \leq t \leq 10$. Show the work that leads to your answer, and indicate units of measure.

Part B

- i. Use a midpoint Riemann sum with the three subintervals $[0, 10]$, $[10, 20]$, and $[20, 30]$ to approximate $\int_0^{30} M(t) dt$. Show the work that leads to your answer.
- ii. Interpret the meaning of $\int_0^{30} M(t) dt$ in the context of the problem.

Part C

The rate at which female birds of the same species arrive at the same nesting area, in birds per day, is modeled by the function F defined as follows.

$$F(t) = \begin{cases} 0 & \text{for } 0 \leq t < 15 \\ 18 + 16 \sin\left(\frac{\pi}{20}(t + 15)\right) & \text{for } 15 \leq t \leq 45 \end{cases}$$

How many female birds of this species arrive at the nesting area from $t = 15$ to $t = 45$? Show the setup for your calculations, and round your answer to the nearest integer.

Part D

On the interval $15 < t < 30$, the difference in the rates at which male and female birds of this species arrive at the nesting area can be modeled by the differentiable function $D(t) = M(t) - F(t)$, where F is the function defined in part C. Is there a time t in the interval $15 < t < 20$ when $D(t) = 0$? Justify your answer.

3. A student starts reading a book at time $t = 0$ minutes and continues reading for the next 10 minutes. The rate at which the student reads is modeled by the differentiable function R , where $R(t)$ is measured in words per minute. Selected values of $R(t)$ are given in the table shown.

t (minutes)	0	2	8	10
$R(t)$ (words per minute)	90	100	150	162

- A. Approximate $R'(1)$ using the average rate of change of R over the interval $0 \leq t \leq 2$. Show the work that leads to your answer. Indicate units of measure.
- B. Must there be a value c , for $0 < c < 10$, such that $R(c) = 155$? Justify your answer.
- C. Use a trapezoidal sum with the three subintervals indicated by the data in the table to approximate the value of $\int_0^{10} R(t) dt$. Show the work that leads to your answer.
- D. A teacher also starts reading at time $t = 0$ minutes and continues reading for the next 10 minutes. The rate at which the teacher reads is modeled by the function W defined by $W(t) = -\frac{3}{10}t^2 + 8t + 100$, where $W(t)$ is measured in words per minute. Based on the model, how many words has the teacher read by the end of the 10 minutes? Show the work that leads to your answer.

r (centimeters)	0	1	2	2.5	4
$f(r)$ (milligrams per square centimeter)	1	2	6	10	18

1. The density of a bacteria population in a circular petri dish at a distance r centimeters from the center of the dish is given by an increasing, differentiable function f , where $f(r)$ is measured in milligrams per square centimeter. Values of $f(r)$ for selected values of r are given in the table above.

(a) Use the data in the table to estimate $f'(2.25)$. Using correct units, interpret the meaning of your answer in the context of this problem.

(b) The total mass, in milligrams, of bacteria in the petri dish is given by the integral expression

$$2\pi \int_0^4 r f(r) dr.$$

Approximate the value of $2\pi \int_0^4 r f(r) dr$ using a right Riemann sum with the four

subintervals indicated by the data in the table.

(c) Is the approximation found in part (b) an overestimate or underestimate of the total mass of bacteria in the petri dish? Explain your reasoning.

(d) The density of bacteria in the petri dish, for $1 \leq r \leq 4$, is modeled by the function g defined by $g(r) = 2 - 16(\cos(1.57\sqrt{r}))^3$. For what value of k , $1 < k < 4$, is $g(k)$ equal to the average value of $g(r)$ on the interval $1 \leq r \leq 4$?

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

6. Functions f , g , and h are twice-differentiable functions with $g(2) = h(2) = 4$. The line $y = 4 + \frac{2}{3}(x - 2)$ is tangent to both the graph of g at $x = 2$ and the graph of h at $x = 2$.

(a) Find $h'(2)$.

(b) Let a be the function given by $a(x) = 3x^3h(x)$. Write an expression for $a'(x)$. Find $a'(2)$.

(c) The function h satisfies $h(x) = \frac{x^2 - 4}{1 - (f(x))^3}$ for $x \neq 2$. It is known that $\lim_{x \rightarrow 2} h(x)$ can be evaluated using

L'Hospital's Rule. Use $\lim_{x \rightarrow 2} h(x)$ to find $f(2)$ and $f'(2)$. Show the work that leads to your answers.

(d) It is known that $g(x) \leq h(x)$ for $1 < x < 3$. Let k be a function satisfying $g(x) \leq k(x) \leq h(x)$ for $1 < x < 3$. Is k continuous at $x = 2$? Justify your answer.

STOP
END OF EXAM

CALCULUS AB
SECTION II, Part A
Time—30 minutes
Number of problems—2

A graphing calculator is required for these problems.

t (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

1. Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.
- Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.
 - Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
 - Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
 - For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

CALCULUS AB
SECTION II, Part B
Time—60 minutes
Number of problems—4

No calculator is allowed for these problems.

t (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	−220	150

3. Johanna jogs along a straight path. For $0 \leq t \leq 40$, Johanna's velocity is given by a differentiable function v . Selected values of $v(t)$, where t is measured in minutes and $v(t)$ is measured in meters per minute, are given in the table above.
- Use the data in the table to estimate the value of $v'(16)$.
 - Using correct units, explain the meaning of the definite integral $\int_0^{40} |v(t)| dt$ in the context of the problem.
 Approximate the value of $\int_0^{40} |v(t)| dt$ using a right Riemann sum with the four subintervals indicated in the table.
 - Bob is riding his bicycle along the same path. For $0 \leq t \leq 10$, Bob's velocity is modeled by $B(t) = t^3 - 6t^2 + 300$, where t is measured in minutes and $B(t)$ is measured in meters per minute. Find Bob's acceleration at time $t = 5$.
 - Based on the model B from part (c), find Bob's average velocity during the interval $0 \leq t \leq 10$.