


2. The shaded region R is bounded by the graphs of the functions f and g , where $f(x) = x^2 - 2x$ and $g(x) = x + \sin(\pi x)$, as shown in the figure.



(Note: Your calculator should be in radian mode.)

- A. Find the area of R . Show the setup for your calculations.
- B. Region R is the base of a solid. For this solid, at each x the cross section perpendicular to the x -axis is a rectangle with height x and base in region R . Find the volume of the solid. Show the setup for your calculations.
- C. Write, but do not evaluate, an integral expression for the volume of the solid generated when the region R is rotated about the horizontal line $y = -2$.
- D. It can be shown that $g'(x) = 1 + \pi \cos(\pi x)$. Find the value of x , for $0 < x < 1$, at which the line tangent to the graph of f is parallel to the line tangent to the graph of g .

END OF PART A

2. A particle moves along the x -axis with velocity given by $v(t) = \frac{10 \sin(0.4t^2)}{t^2 - t + 3}$ for time $0 \leq t \leq 3.5$.

The particle is at position $x = -5$ at time $t = 0$.

(a) Find the acceleration of the particle at time $t = 3$.

(b) Find the position of the particle at time $t = 3$.

(c) Evaluate $\int_0^{3.5} v(t) \, dt$, and evaluate $\int_0^{3.5} |v(t)| \, dt$. Interpret the meaning of each integral in the context of the problem.

(d) A second particle moves along the x -axis with position given by $x_2(t) = t^2 - t$ for $0 \leq t \leq 3.5$. At what time t are the two particles moving with the same velocity?

END OF PART A OF SECTION II