

Part B (AB): Graphing calculator not allowed**Question 5****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Consider the curve defined by the equation $x^2 + 3y + 2y^2 = 48$. It can be shown that $\frac{dy}{dx} = \frac{-2x}{3 + 4y}$.

Model Solution**Scoring**

- (a) There is a point on the curve near $(2, 4)$ with x -coordinate 3. Use the line tangent to the curve at $(2, 4)$ to approximate the y -coordinate of this point.

$\left. \frac{dy}{dx} \right _{(x, y)=(2, 4)} = \frac{-2(2)}{3 + 4(4)} = -\frac{4}{19}$	Slope of tangent line	1 point
$y \approx 4 - \frac{4}{19}(3 - 2) = \frac{72}{19}$	Approximation	1 point

Scoring notes:

- A response earns the first point for finding $\left. \frac{dy}{dx} \right|_{(x, y)=(2, 4)} = -\frac{4}{19}$, even if this is not labeled or used as the slope of a tangent line.
- A response that does not explicitly find the value of $\left. \frac{dy}{dx} \right|_{(x, y)=(2, 4)}$ but uses a slope of $-\frac{4}{19}$ in a linear approximation also earns the first point.
- A response that declares $\left. \frac{dy}{dx} \right|_{(x, y)=(2, 4)}$ equal to any nonzero value other than $-\frac{4}{19}$ does not earn the first point but is eligible for the second point for a linear approximation at $x = 3$ through the point $(2, 4)$ with a slope of the declared value.
 - The second point cannot be earned with a linear approximation using a slope other than $-\frac{4}{19}$ if that slope has not been declared to be the value of $\left. \frac{dy}{dx} \right|_{(x, y)=(2, 4)}$.
- The second point cannot be earned for an approximation at any value of x other than 3.
- A response does not have to write the tangent line equation but must clearly demonstrate its use at $x = 3$ in finding the requested approximation in order to earn both points.
- The minimal work required to earn both points is $4 - \frac{4}{19}(3 - 2)$.

Total for part (a) 2 points

- (b) Is the horizontal line $y = 1$ tangent to the curve? Give a reason for your answer.

$\frac{dy}{dx} = \frac{-2x}{3 + 4y} = 0 \Rightarrow x = 0$	Considers $\frac{dy}{dx} = 0$	1 point
And so, if the horizontal line $y = 1$ is tangent to the curve, the point of tangency must be $(0, 1)$.	Answer with reason	1 point
However, the point $(0, 1)$ is not on the curve, because $0^2 + 3 \cdot 1 + 2 \cdot 1^2 = 5 \neq 48$.		
Therefore, the horizontal line $y = 1$ is not tangent to the curve.		

Scoring notes:

- The first point can be earned with a response of $\frac{dy}{dx} = 0$, $-2x = 0$, or $x = 0$, OR by identifying the point $(0, 1)$.
- To earn the second point a response must provide a reason that the line $y = 1$ is not tangent to the curve. Merely stating “ $(0, 1)$ does not lie on the curve” is not sufficient to earn the second point.
- Alternate solution:
 - If $y = 1$, then $x^2 + 3 \cdot 1 + 2 \cdot 1^2 = 48 \Rightarrow x = \pm\sqrt{43}$.

$$\left. \frac{dy}{dx} \right|_{(x, y)=(\pm\sqrt{43}, 1)} = \frac{\pm 2\sqrt{43}}{7} \neq 0$$
 Therefore, the horizontal line $y = 1$ is not tangent to the curve.
 - A response that uses this method earns the first point by using $y = 1$ in $x^2 + 3y + 2y^2 = 48$.
 - A response that fails to consider both $x = +\sqrt{43}$ and $x = -\sqrt{43}$ does not earn the second point.

Total for part (b) 2 points

- (c) The curve intersects the positive x -axis at the point $(\sqrt{48}, 0)$. Is the line tangent to the curve at this point vertical? Give a reason for your answer.

At the point $(\sqrt{48}, 0)$, the slope of the line tangent to the curve is $\frac{dy}{dx} = \frac{-2\sqrt{48}}{3 + 4(0)}$.	Answer with reason	1 point
The denominator of $\frac{dy}{dx}$ is $3 + 4(0)$, which does not equal 0.		
Therefore, the line tangent to the curve at this point is not vertical.		

Scoring notes:

- A response does not need to consider the numerator of $\frac{dy}{dx}\bigg|_{(x,y)=(\sqrt{48},0)}$ in order to earn this point; considering the denominator is sufficient.
- To earn this point a response must clearly demonstrate that the slope of the tangent line at the point $(\sqrt{48}, 0)$ is defined and answer “no.”
 - Such demonstrations include, but are not limited to, the following:
 - $3 + 4(0) \neq 0$
 - At $(\sqrt{48}, 0)$, $\frac{dy}{dx} = \frac{-2\sqrt{48}}{3}$.
 - $\frac{-2\sqrt{48}}{3}$
 - When $3 + 4y = 0$, $y \neq 0$.

Total for part (c) 1 point

- (d) For time $t \geq 0$, a particle is moving along another curve defined by the equation $y^3 + 2xy = 24$. At the instant the particle is at the point $(4, 2)$, the y -coordinate of the particle's position is decreasing at a rate of 2 units per second. At that instant, what is the rate of change of the x -coordinate of the particle's position with respect to time?

$3y^2 \frac{dy}{dt} + 2x \frac{dy}{dt} + 2y \frac{dx}{dt} = 0$	Attempts implicit differentiation	1 point
	$3y^2 \frac{dy}{dt} + 2x \frac{dy}{dt} + 2y \frac{dx}{dt} = 0$	1 point
$\frac{dy}{dt} = -2$	Uses $\frac{dy}{dt} = \pm 2$	1 point
$3(2)^2(-2) + 2(4)(-2) + 2(2)\frac{dx}{dt} = 0 \Rightarrow \frac{dx}{dt} = \frac{40}{4} = 10$	Answer	1 point
The rate of change with respect to time in the x -coordinate is 10 units per second.		

Scoring notes:

- The first point is earned for implicitly differentiating $y^3 + 2xy = 24$ with respect to t with at most one error.
 - The first point can also be earned by correctly differentiating with respect to x :

$$3y^2 \frac{dy}{dx} + 2x \frac{dy}{dx} + 2y = 0.$$
- The second point is earned for an equation equivalent to $3y^2 \frac{dy}{dt} + 2x \frac{dy}{dt} + 2y \frac{dx}{dt} = 0$.
- A response does not need to explicitly declare $\frac{dy}{dt} = -2$ or $\frac{dy}{dt} = 2$ in order to earn the third point; this point may be earned by correctly substituting $\frac{dy}{dt} = -2$ or $\frac{dy}{dt} = 2$ in the implicitly differentiated equation. However, a response that uses both $\frac{dy}{dt} = -2$ and $\frac{dy}{dt} = 2$ in the implicitly differentiated equation does not earn the third point.
- The fourth point cannot be earned without the first 3 points. The fourth point is earned only for the value of 10 with no mistakes in supporting work.
 - Note that a response that uses $\frac{dy}{dt} = 2$ and then mishandles subtracting 40 from both sides of the equation, e.g., $3(2)^2(2) + 2(4)(2) + 2(2)\frac{dx}{dt} = 0 \Rightarrow \frac{dx}{dt} = \frac{40}{4}$, does not earn the fourth point.

Total for part (d) 4 points**Total for question 5 9 points**

AP Calculus AB

Scoring Guidelines

2018 SCORING GUIDELINES

Question 1

(a) $\int_0^{300} r(t) \, dt = 270$

According to the model, 270 people enter the line for the escalator during the time interval $0 \leq t \leq 300$.

(b) $20 + \int_0^{300} (r(t) - 0.7) \, dt = 20 + \int_0^{300} r(t) \, dt - 0.7 \cdot 300 = 80$

According to the model, 80 people are in line at time $t = 300$.

(c) Based on part (b), the number of people in line at time $t = 300$ is 80.

The first time t that there are no people in line is

$$300 + \frac{80}{0.7} = 414.286 \text{ (or 414.285) seconds.}$$

(d) The total number of people in line at time t , $0 \leq t \leq 300$, is modeled by $20 + \int_0^t r(x) \, dx - 0.7t$.

$$r(t) - 0.7 = 0 \Rightarrow t_1 = 33.013298, t_2 = 166.574719$$

t	People in line for escalator
0	20
t_1	3.803
t_2	158.070
300	80

The number of people in line is a minimum at time $t = 33.013$ seconds, when there are 4 people in line.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{considers rate out} \\ 1 : \text{answer} \end{cases}$

1 : answer

4 : $\begin{cases} 1 : \text{considers } r(t) - 0.7 = 0 \\ 1 : \text{identifies } t = 33.013 \\ 1 : \text{answers} \\ 1 : \text{justification} \end{cases}$

2018 SCORING GUIDELINES

Question 2

(a) $v'(3) = -2.118$

The acceleration of the particle at time $t = 3$ is -2.118 .

1 : answer

(b) $x(3) = x(0) + \int_0^3 v(t) dt = -5 + \int_0^3 v(t) dt = -1.760213$

The position of the particle at time $t = 3$ is -1.760 .

3 : $\begin{cases} 1 : \int_0^3 v(t) dt \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

(c) $\int_0^{3.5} v(t) dt = 2.844$ (or 2.843)

$$\int_0^{3.5} |v(t)| dt = 3.737$$

The integral $\int_0^{3.5} v(t) dt$ is the displacement of the particle over the time interval $0 \leq t \leq 3.5$.

The integral $\int_0^{3.5} |v(t)| dt$ is the total distance traveled by the particle over the time interval $0 \leq t \leq 3.5$.

3 : $\begin{cases} 1 : \text{answers} \\ 2 : \text{interpretations of } \int_0^{3.5} v(t) dt \\ \text{and } \int_0^{3.5} |v(t)| dt \end{cases}$

(d) $v(t) = x_2'(t)$

$$v(t) = 2t - 1 \Rightarrow t = 1.57054$$

The two particles are moving with the same velocity at time $t = 1.571$ (or 1.570).

2 : $\begin{cases} 1 : \text{sets } v(t) = x_2'(t) \\ 1 : \text{answer} \end{cases}$

2018 SCORING GUIDELINES

Question 3

$$\begin{aligned} \text{(a)} \quad f(-5) &= f(1) + \int_1^{-5} g(x) dx = f(1) - \int_{-5}^1 g(x) dx \\ &= 3 - \left(-9 - \frac{3}{2} + 1\right) = 3 - \left(-\frac{19}{2}\right) = \frac{25}{2} \end{aligned}$$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b)} \quad \int_1^6 g(x) dx &= \int_1^3 g(x) dx + \int_3^6 g(x) dx \\ &= \int_1^3 2 dx + \int_3^6 2(x-4)^2 dx \\ &= 4 + \left[\frac{2}{3}(x-4)^3\right]_{x=3}^{x=6} = 4 + \frac{16}{3} - \left(-\frac{2}{3}\right) = 10 \end{aligned}$$

3 : $\begin{cases} 1 : \text{split at } x = 3 \\ 1 : \text{antiderivative of } 2(x-4)^2 \\ 1 : \text{answer} \end{cases}$

(c) The graph of f is increasing and concave up on $0 < x < 1$ and $4 < x < 6$ because $f'(x) = g(x) > 0$ and $f''(x) = g(x)$ is increasing on those intervals.

2 : $\begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$

(d) The graph of f has a point of inflection at $x = 4$ because $f''(x) = g(x)$ changes from decreasing to increasing at $x = 4$.

2 : $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

2018 SCORING GUIDELINES

Question 4

$$(a) \quad H'(6) \approx \frac{H(7) - H(5)}{7 - 5} = \frac{11 - 6}{2} = \frac{5}{2}$$

$H'(6)$ is the rate at which the height of the tree is changing, in meters per year, at time $t = 6$ years.

$$(b) \quad \frac{H(5) - H(3)}{5 - 3} = \frac{6 - 2}{2} = 2$$

Because H is differentiable on $3 \leq t \leq 5$, H is continuous on $3 \leq t \leq 5$.

By the Mean Value Theorem, there exists a value c , $3 < c < 5$, such that $H'(c) = 2$.

(c) The average height of the tree over the time interval $2 \leq t \leq 10$ is given by $\frac{1}{10 - 2} \int_2^{10} H(t) \, dt$.

$$\begin{aligned} \frac{1}{8} \int_2^{10} H(t) \, dt &\approx \frac{1}{8} \left(\frac{1.5 + 2}{2} \cdot 1 + \frac{2 + 6}{2} \cdot 2 + \frac{6 + 11}{2} \cdot 2 + \frac{11 + 15}{2} \cdot 3 \right) \\ &= \frac{1}{8} (65.75) = \frac{263}{32} \end{aligned}$$

The average height of the tree over the time interval $2 \leq t \leq 10$ is $\frac{263}{32}$ meters.

$$(d) \quad G(x) = 50 \Rightarrow x = 1$$

$$\frac{d}{dt}(G(x)) = \frac{d}{dx}(G(x)) \cdot \frac{dx}{dt} = \frac{(1+x)100 - 100x \cdot 1}{(1+x)^2} \cdot \frac{dx}{dt} = \frac{100}{(1+x)^2} \cdot \frac{dx}{dt}$$

$$\left. \frac{d}{dt}(G(x)) \right|_{x=1} = \frac{100}{(1+1)^2} \cdot 0.03 = \frac{3}{4}$$

According to the model, the rate of change of the height of the tree with respect to time when the tree is 50 meters tall is $\frac{3}{4}$ meter per year.

2 : $\begin{cases} 1 : \text{estimate} \\ 1 : \text{interpretation with units} \end{cases}$

2 : $\begin{cases} 1 : \frac{H(5) - H(3)}{5 - 3} \\ 1 : \text{conclusion using Mean Value Theorem} \end{cases}$

2 : $\begin{cases} 1 : \text{trapezoidal sum} \\ 1 : \text{approximation} \end{cases}$

3 : $\begin{cases} 2 : \frac{d}{dt}(G(x)) \\ 1 : \text{answer} \end{cases}$

Note: max 1/3 [1-0] if no chain rule

2018 SCORING GUIDELINES

Question 5

(a) The average rate of change of f on the interval $0 \leq x \leq \pi$ is

$$\frac{f(\pi) - f(0)}{\pi - 0} = \frac{-e^\pi - 1}{\pi}.$$

(b) $f'(x) = e^x \cos x - e^x \sin x$

$$f'\left(\frac{3\pi}{2}\right) = e^{3\pi/2} \cos\left(\frac{3\pi}{2}\right) - e^{3\pi/2} \sin\left(\frac{3\pi}{2}\right) = e^{3\pi/2}$$

The slope of the line tangent to the graph of f at $x = \frac{3\pi}{2}$ is $e^{3\pi/2}$.

(c) $f'(x) = 0 \Rightarrow \cos x - \sin x = 0 \Rightarrow x = \frac{\pi}{4}, x = \frac{5\pi}{4}$

x	$f(x)$
0	1
$\frac{\pi}{4}$	$\frac{1}{\sqrt{2}} e^{\pi/4}$
$\frac{5\pi}{4}$	$-\frac{1}{\sqrt{2}} e^{5\pi/4}$
2π	$e^{2\pi}$

The absolute minimum value of f on $0 \leq x \leq 2\pi$ is $-\frac{1}{\sqrt{2}} e^{5\pi/4}$.

(d) $\lim_{x \rightarrow \pi/2} f(x) = 0$

Because g is differentiable, g is continuous.

$$\lim_{x \rightarrow \pi/2} g(x) = g\left(\frac{\pi}{2}\right) = 0$$

By L'Hospital's Rule,

$$\lim_{x \rightarrow \pi/2} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \pi/2} \frac{f'(x)}{g'(x)} = \frac{-e^{\pi/2}}{2}.$$

1 : answer

2 : $\begin{cases} 1 : f'(x) \\ 1 : \text{slope} \end{cases}$

3 : $\begin{cases} 1 : \text{sets } f'(x) = 0 \\ 1 : \text{identifies } x = \frac{\pi}{4}, x = \frac{5\pi}{4} \\ \text{as candidates} \\ 1 : \text{answer with justification} \end{cases}$


3 : $\begin{cases} 1 : g \text{ is continuous at } x = \frac{\pi}{2} \\ \text{and limits equal 0} \\ 1 : \text{applies L'Hospital's Rule} \\ 1 : \text{answer} \end{cases}$

Note: max 1/3 [1-0-0] if no limit notation attached to a ratio of derivatives

2018 SCORING GUIDELINES

Question 6

(a)



- 2 : $\begin{cases} 1 : \text{solution curve through } (0, 2) \\ 1 : \text{solution curve through } (1, 0) \end{cases}$

Curves must go through the indicated points, follow the given slope lines, and extend to the boundary of the slope field.

(b) $\left. \frac{dy}{dx} \right|_{(x,y)=(1,0)} = \frac{4}{3}$

An equation for the line tangent to the graph of $y = f(x)$ at

$x = 1$ is $y = \frac{4}{3}(x - 1)$.

$f(0.7) \approx \frac{4}{3}(0.7 - 1) = -0.4$

- 2 : $\begin{cases} 1 : \text{equation of tangent line} \\ 1 : \text{approximation} \end{cases}$

(c) $\frac{dy}{dx} = \frac{1}{3}x(y - 2)^2$
 $\int \frac{dy}{(y - 2)^2} = \int \frac{1}{3}x \, dx$

$\frac{-1}{y - 2} = \frac{1}{6}x^2 + C$

$\frac{1}{2} = \frac{1}{6} + C \Rightarrow C = \frac{1}{3}$

$\frac{-1}{y - 2} = \frac{1}{6}x^2 + \frac{1}{3} = \frac{x^2 + 2}{6}$

$y = 2 - \frac{6}{x^2 + 2}$

Note: this solution is valid for $-\infty < x < \infty$.

- 5 : $\begin{cases} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ \text{and uses initial condition} \\ 1 : \text{solves for } y \end{cases}$

Note: 0/5 if no separation of variables

Note: max 3/5 [1-2-0-0] if no constant of integration