

1. An invasive species of plant appears in a fruit grove at time $t = 0$ and begins to spread. The function C defined by $C(t) = 7.6 \arctan(0.2t)$ models the number of acres in the fruit grove affected by the species t weeks after the species appears. It can be shown that $C'(t) = \frac{38}{25 + t^2}$.

(Note: Your calculator should be in radian mode.)

- A. Find the average number of acres affected by the invasive species from time $t = 0$ to time $t = 4$ weeks. Show the setup for your calculations.
- B. Find the time t when the instantaneous rate of change of C equals the average rate of change of C over the time interval $0 \leq t \leq 4$. Show the setup for your calculations.
- C. Assume that the invasive species continues to spread according to the given model for all times $t > 0$. Write a limit expression that describes the end behavior of the rate of change in the number of acres affected by the species. Evaluate this limit expression.
- D. At time $t = 4$ weeks after the invasive species appears in the fruit grove, measures are taken to counter the spread of the species. The function A , defined by $A(t) = C(t) - \int_4^t 0.1 \cdot \ln(x) dx$, models the number of acres affected by the species over the time interval $4 \leq t \leq 36$. At what time t , for $4 \leq t \leq 36$, does A attain its maximum value? Justify your answer.

t (seconds)	0	60	90	120	135	150
$f(t)$ (gallons per second)	0	0.1	0.15	0.1	0.05	0

1. A customer at a gas station is pumping gasoline into a gas tank. The rate of flow of gasoline is modeled by a differentiable function f , where $f(t)$ is measured in gallons per second and t is measured in seconds since pumping began. Selected values of $f(t)$ are given in the table.

- (a) Using correct units, interpret the meaning of $\int_{60}^{135} f(t) dt$ in the context of the problem. Use a right

Riemann sum with the three subintervals $[60, 90]$, $[90, 120]$, and $[120, 135]$ to approximate the value of

$$\int_{60}^{135} f(t) dt.$$

- (b) Must there exist a value of c , for $60 < c < 120$, such that $f'(c) = 0$? Justify your answer.


- (c) The rate of flow of gasoline, in gallons per second, can also be modeled by $g(t) = \left(\frac{t}{500}\right) \cos\left(\left(\frac{t}{120}\right)^2\right)$ for

$0 \leq t \leq 150$. Using this model, find the average rate of flow of gasoline over the time interval $0 \leq t \leq 150$.

Show the setup for your calculations.

- (d) Using the model g defined in part (c), find the value of $g'(140)$. Interpret the meaning of your answer in the context of the problem.

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

Graph of f

Let f be a continuous function defined on the closed interval $-4 \leq x \leq 6$. The graph of f , consisting of four line segments, is shown above. Let G be the function defined by $G(x) = \int_0^x f(t) \, dt$.

(a) On what open intervals is the graph of G concave up? Give a reason for your answer.

(b) Let P be the function defined by $P(x) = G(x) \cdot f(x)$. Find $P'(3)$.

(c) Find $\lim_{x \rightarrow 2} \frac{G(x)}{x^2 - 2x}$.

(d) Find the average rate of change of G on the interval $[-4, 2]$. Does the Mean Value Theorem guarantee a value c , $-4 < c < 2$, for which $G'(c)$ is equal to this average rate of change? Justify your answer.

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

t (hours)	0	0.3	1.7	2.8	4
$v_P(t)$ (meters per hour)	0	55	-29	55	48


2. The velocity of a particle, P , moving along the x -axis is given by the differentiable function v_P , where $v_P(t)$ is measured in meters per hour and t is measured in hours. Selected values of $v_P(t)$ are shown in the table above. Particle P is at the origin at time $t = 0$.
- (a) Justify why there must be at least one time t , for $0.3 \leq t \leq 2.8$, at which $v_P'(t)$, the acceleration of particle P , equals 0 meters per hour per hour.
- (b) Use a trapezoidal sum with the three subintervals $[0, 0.3]$, $[0.3, 1.7]$, and $[1.7, 2.8]$ to approximate the value of $\int_0^{2.8} v_P(t) dt$.
- (c) A second particle, Q , also moves along the x -axis so that its velocity for $0 \leq t \leq 4$ is given by $v_Q(t) = 45\sqrt{t} \cos(0.063t^2)$ meters per hour. Find the time interval during which the velocity of particle Q is at least 60 meters per hour. Find the distance traveled by particle Q during the interval when the velocity of particle Q is at least 60 meters per hour.
- (d) At time $t = 0$, particle Q is at position $x = -90$. Using the result from part (b) and the function v_Q from part (c), approximate the distance between particles P and Q at time $t = 2.8$.
-

END OF PART A OF SECTION II

t (years)	2	3	5	7	10
$H(t)$ (meters)	1.5	2	6	11	15

4. The height of a tree at time t is given by a twice-differentiable function H , where $H(t)$ is measured in meters and t is measured in years. Selected values of $H(t)$ are given in the table above.
- (a) Use the data in the table to estimate $H'(6)$. Using correct units, interpret the meaning of $H'(6)$ in the context of the problem.
- (b) Explain why there must be at least one time t , for $2 < t < 10$, such that $H'(t) = 2$.
- (c) Use a trapezoidal sum with the four subintervals indicated by the data in the table to approximate the average height of the tree over the time interval $2 \leq t \leq 10$.
- (d) The height of the tree, in meters, can also be modeled by the function G , given by $G(x) = \frac{100x}{1+x}$, where x is the diameter of the base of the tree, in meters. When the tree is 50 meters tall, the diameter of the base of the tree is increasing at a rate of 0.03 meter per year. According to this model, what is the rate of change of the height of the tree with respect to time, in meters per year, at the time when the tree is 50 meters tall?
-

x	$g(x)$	$g'(x)$
-5	10	-3
-4	5	-1
-3	2	4
-2	3	1
-1	1	-2
0	0	-3

Graph of h

6. Let f be the function defined by $f(x) = \cos(2x) + e^{\sin x}$.

Let g be a differentiable function. The table above gives values of g and its derivative g' at selected values of x .

Let h be the function whose graph, consisting of five line segments, is shown in the figure above.

- (a) Find the slope of the line tangent to the graph of f at $x = \pi$.
- (b) Let k be the function defined by $k(x) = h(f(x))$. Find $k'(\pi)$.
- (c) Let m be the function defined by $m(x) = g(-2x) \cdot h(x)$. Find $m'(2)$.
- (d) Is there a number c in the closed interval $[-5, -3]$ such that $g'(c) = -4$? Justify your answer.

STOP
END OF EXAM

x	-2	$-2 < x < -1$	-1	$-1 < x < 1$	1	$1 < x < 3$	3
$f(x)$	12	Positive	8	Positive	2	Positive	7
$f'(x)$	-5	Negative	0	Negative	0	Positive	$\frac{1}{2}$
$g(x)$	-1	Negative	0	Positive	3	Positive	1
$g'(x)$	2	Positive	$\frac{3}{2}$	Positive	0	Negative	-2

5. The twice-differentiable functions f and g are defined for all real numbers x . Values of f , f' , g , and g' for various values of x are given in the table above.

(a) Find the x -coordinate of each relative minimum of f on the interval $[-2, 3]$. Justify your answers.

(b) Explain why there must be a value c , for $-1 < c < 1$, such that $f''(c) = 0$.

(c) The function h is defined by $h(x) = \ln(f(x))$. Find $h'(3)$. Show the computations that lead to your answer.

(d) Evaluate $\int_{-2}^3 f'(g(x))g'(x) dx$.