

Part B (AB or BC): Graphing calculator not allowed**Question 3****9 points****General Scoring Notes**


The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

The depth of seawater at a location can be modeled by the function H that satisfies the differential equation $\frac{dH}{dt} = \frac{1}{2}(H - 1)\cos\left(\frac{t}{2}\right)$, where $H(t)$ is measured in feet and t is measured in hours after noon ($t = 0$). It is known that $H(0) = 4$.

Model Solution**Scoring**

- (a) A portion of the slope field for the differential equation is provided. Sketch the solution curve, $y = H(t)$, through the point $(0, 4)$.



Solution curve

1 point**Scoring notes:**

- The solution curve must pass through the point $(0, 4)$, extend to at least $t = 4.5$, and have no obvious conflicts with the given slope lines.
- Only portions of the solution curve within the given slope field are considered.

Total for part (a) 1 point

- (b) For $0 < t < 5$, it can be shown that $H(t) > 1$. Find the value of t , for $0 < t < 5$, at which H has a critical point. Determine whether the critical point corresponds to a relative minimum, a relative maximum, or neither a relative minimum nor a relative maximum of the depth of seawater at the location. Justify your answer.

Because $H(t) > 1$, then $\frac{dH}{dt} = 0$ implies $\cos\left(\frac{t}{2}\right) = 0$.

Considers sign of $\frac{dH}{dt}$ **1 point**

This implies that $t = \pi$ is a critical point.

Identifies $t = \pi$ **1 point**

For $0 < t < \pi$, $\frac{dH}{dt} > 0$ and for $\pi < t < 5$, $\frac{dH}{dt} < 0$. Therefore, $t = \pi$ is the location of a relative maximum value of H .

Answer with justification **1 point**

Scoring notes:

- The first point is earned for considering $\frac{dH}{dt} = 0$, $\frac{dH}{dt} > 0$, $\frac{dH}{dt} < 0$, $\cos\left(\frac{t}{2}\right) = 0$, $\cos\left(\frac{t}{2}\right) > 0$, or $\cos\left(\frac{t}{2}\right) < 0$.
- The second point is earned for identifying $t = \pi$, with or without supporting work. A response may consider $H = 1$ or $t = 1$ as potential critical points without penalty.
- The third point cannot be earned without the first point. The third point is earned only for a correct justification and a correct answer of "relative maximum."
- The justification can be shown by determining the sign of $\frac{dH}{dt}$ (or $\cos\left(\frac{t}{2}\right)$) at a single value in $0 < t < \pi$ and at a single value in $\pi < t < 5$. It is not necessary to state that $\frac{dH}{dt}$ does not change sign on these intervals.
- The third point can also be earned by using the Second Derivative Test. For example:

$$\frac{d^2H}{dt^2} = \frac{1}{2}(H - 1)\left(-\frac{1}{2}\sin\left(\frac{t}{2}\right)\right) + \cos\left(\frac{t}{2}\right) \cdot \frac{1}{2} \cdot \frac{dH}{dt}$$

$$\frac{d^2H}{dt^2}\bigg|_{t=\pi} < 0$$
 Therefore, $t = \pi$ is the location of a relative maximum value of H .

Total for part (b) 3 points

- (c) Use separation of variables to find $y = H(t)$, the particular solution to the differential equation

$$\frac{dH}{dt} = \frac{1}{2}(H-1)\cos\left(\frac{t}{2}\right) \text{ with initial condition } H(0) = 4.$$

$\frac{dH}{H-1} = \frac{1}{2}\cos\left(\frac{t}{2}\right)dt$	Separation of variables	1 point
$\int \frac{dH}{H-1} = \int \frac{1}{2}\cos\left(\frac{t}{2}\right)dt$	One antiderivative	1 point
$\Rightarrow \ln H-1 = \sin\left(\frac{t}{2}\right) + C$	Second antiderivative	1 point
$\ln 4-1 = \sin\left(\frac{0}{2}\right) + C \Rightarrow C = \ln 3$ Because $H(0) = 4$, $H > 1$, so $ H-1 = H-1$. $\ln(H-1) = \sin\left(\frac{t}{2}\right) + \ln 3$	Constant of integration and uses initial condition	1 point
$H-1 = e^{\sin(t/2)+\ln 3} = 3e^{\sin(t/2)}$ $H(t) = 1 + 3e^{\sin(t/2)}$	Solves for H	1 point

Scoring notes:

- A response with no separation of variables earns 0 out of 5 points.
- A response that presents $\int \frac{dH}{H-1} = \ln(H-1)$ without absolute value symbols earns that antiderivative point.
- A response with no constant of integration can earn at most the first 3 points.
- A response is eligible for the fourth point only if it has earned the first point and at least 1 of the 2 antiderivative points.
- An eligible response earns the fourth point by correctly including the constant of integration in an equation and substituting 0 for t and 4 for H .
- A response is eligible for the fifth point only if it has earned the first 4 points.
- A response earns the fifth point only for an answer of $H(t) = 1 + 3e^{\sin(t/2)}$ or a mathematically equivalent expression for $H(t)$ such as $H(t) = 1 + e^{\sin(t/2)+\ln 3}$.
- A response does not need to argue that $|H-1| = H-1$ in order to earn the fifth point.
- Special case: A response that presents an incorrect separation of variables of $\frac{1}{2} \cdot \frac{dH}{H-1} = \cos\left(\frac{t}{2}\right)dt$ does not earn the first point or the fifth point but is eligible for the 2 antiderivative points. If the response earns at least 1 of the 2 antiderivative points, then the response is eligible for the fourth point.

Total for part (c) 5 points

Total for question 3 9 points

Part B (AB or BC): Graphing calculator not allowed


Question 4

9 points

General Scoring Notes

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.



The graph of the differentiable function f , shown for $-6 \leq x \leq 7$, has a horizontal tangent at $x = -2$ and is linear for $0 \leq x \leq 7$. Let R be the region in the second quadrant bounded by the graph of f , the vertical line $x = -6$, and the x - and y -axes. Region R has area 12.

Model Solution	Scoring
(a) The function g is defined by $g(x) = \int_0^x f(t) dt$. Find the values of $g(-6)$, $g(4)$, and $g(6)$.	
$g(-6) = \int_0^{-6} f(t) dt = -\int_{-6}^0 f(t) dt = -12$	$g(-6)$ 1 point
$g(4) = \int_0^4 f(t) dt = \frac{1}{2} \cdot 4 \cdot 2 = 4$	$g(4)$ 1 point
$g(6) = \int_0^6 f(t) dt = \frac{1}{2} \cdot 4 \cdot 2 - \frac{1}{2} \cdot 2 \cdot 1 = 3$	$g(6)$ 1 point

Scoring notes:

- Supporting work is not required for any of these values. However, any supporting work that is shown must be correct to earn the corresponding point.
- Special case: A response that explicitly presents $g(x) = \int_{-6}^x f(t) dt$ does not earn the first point it would have otherwise earned. The response is eligible for all subsequent points for correct answers, or for consistent answers with supporting work.
 - Note: $\int_{-6}^{-6} f(t) dt = 0$, $\int_{-6}^4 f(t) dt = 16$, $\int_{-6}^6 f(t) dt = 15$
- Labeled values may be presented in any order. Unlabeled values are read from left to right and from top to bottom as $g(-6)$, $g(4)$, and $g(6)$, respectively. A response that presents only 1 or 2 values must label them to earn any points.

Total for part (a) 3 points

- (b) For the function g defined in part (a), find all values of x in the interval $0 \leq x \leq 6$ at which the graph of g has a critical point. Give a reason for your answer.

$g'(x) = f(x)$	Fundamental Theorem of Calculus	1 point
$g'(x) = f(x) = 0 \Rightarrow x = 4$	Answer with reason	1 point
Therefore, the graph of g has a critical point at $x = 4$.		

Scoring notes:

- The first point is earned for explicitly making the connection $g' = f$ in this part.
 - A response that writes $g'' = f'$ earns the first point but can only earn the second point by reasoning from $f = 0$.
- A response that does not earn the first point is eligible to earn the second point with an implied application of the FTC (e.g., “Because $g'(4) = 0$, $x = 4$ is a critical point”).
- A response that reports any additional critical points in $0 < x < 6$ does not earn the second point.
 - Any presented critical point outside the interval $0 < x < 6$ will not affect scoring.

Total for part (b) 2 points

- (c) The function h is defined by $h(x) = \int_{-6}^x f'(t) dt$. Find the values of $h(6)$, $h'(6)$, and $h''(6)$. Show the work that leads to your answers.

$h(6) = \int_{-6}^6 f'(t) dt = f(6) - f(-6) = -1 - 0.5 = -1.5$	Uses Fundamental Theorem of Calculus	1 point
	$h(6)$ with supporting work	1 point
$h'(x) = f'(x)$, so $h'(6) = f'(6) = -\frac{1}{2}$.	$h'(6)$	1 point
$h''(x) = f''(x)$, so $h''(6) = f''(6) = 0$.	$h''(6)$	1 point

Scoring notes:


- Labeled values may be presented in any order.
- Unlabeled values are read from left to right and from top to bottom as $h(6)$, $h'(6)$, and $h''(6)$, respectively. A response that presents only 1 or 2 values must label them in order to earn any points.
- A response of $h(6) = -1.5$ does not earn either of the first 2 points. A response of $h(6) = f(6) - f(-6)$ earns the first point but not yet the second point.
- A response of $h(6) = -1 - 0.5$ is the minimum work required to earn both of the first 2 points.
- To earn the third point a response must state either $h'(x) = f'(x)$ or $h'(6) = f'(6)$, and provide an answer of $-\frac{1}{2}$.
- The fourth point is earned for a response of $h''(6) = 0$, with or without supporting work.
- A response that has one or more linkage errors does not earn the first point it would have otherwise earned. For example, $h'(x) = f'(6) = -\frac{1}{2}$ does not earn the third point but is eligible for the fourth point even in the presence of another linkage error, such as $h''(x) = f''(6) = 0$.

Total for part (c) 4 points**Total for question 4 9 points**

Part B (AB or BC): Graphing calculator not allowed**Question 3****9 points****General Scoring Notes**

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Scoring guidelines and notes contain examples of the most common approaches seen in student responses. These guidelines can be applied to alternate approaches to ensure that these alternate approaches are scored appropriately.



A company designs spinning toys using the family of functions $y = cx\sqrt{4-x^2}$, where c is a positive constant.

The figure above shows the region in the first quadrant bounded by the x -axis and the graph of $y = cx\sqrt{4-x^2}$, for some c . Each spinning toy is in the shape of the solid generated when such a region is revolved about the x -axis. Both x and y are measured in inches.

Model Solution**Scoring**

- (a) Find the area of the region in the first quadrant bounded by the x -axis and the graph of $y = cx\sqrt{4-x^2}$ for $c = 6$.

$$6x\sqrt{4-x^2} = 0 \Rightarrow x = 0, x = 2$$

Integrand

1 point

$$\text{Area} = \int_0^2 6x\sqrt{4-x^2} \, dx$$

$$\text{Let } u = 4 - x^2.$$

Antiderivative

1 point

$$du = -2x \, dx \Rightarrow -\frac{1}{2} du = x \, dx$$

$$x = 0 \Rightarrow u = 4 - 0^2 = 4$$

$$x = 2 \Rightarrow u = 4 - 2^2 = 0$$

$$\int_0^2 6x\sqrt{4-x^2} \, dx = \int_4^0 6\left(-\frac{1}{2}\right)\sqrt{u} \, du = -3 \int_4^0 u^{1/2} \, du = 3 \int_0^4 u^{1/2} \, du$$

$$= 2u^{3/2} \Big|_{u=0}^{u=4} = 2 \cdot 8 = 16$$

The area of the region is 16 square inches.

Answer

1 point**Scoring notes:**

- Units are not required for any points in this question and are not read if presented (correctly or incorrectly) in any part of the response.
- The first point is earned for presenting $cx\sqrt{4-x^2}$ or $6x\sqrt{4-x^2}$ as the integrand in a definite integral. Limits of integration (numeric or alphanumeric) must be presented (as part of the definite integral) but do not need to be correct in order to earn the first point.
- If an indefinite integral is presented with an integrand of the correct form, the first point can be earned if the antiderivative (correct or incorrect) is eventually evaluated using the correct limits of integration.
- The second point can be earned without the first point. The second point is earned for the presentation of a correct antiderivative of a function of the form $Ax\sqrt{4-x^2}$, for any nonzero constant A . If the response has subsequent errors in simplification of the antiderivative or sign errors, the response will earn the second point but will not earn the third point.
- Responses that use u -substitution and have incorrect limits of integration or do not change the limits of integration from x - to u -values are eligible for the second point.
- The response is eligible for the third point only if it has earned the second point.
- The third point is earned only for the answer 16 or equivalent. In the case where a response only presents an indefinite integral, the use of the correct limits of integration to evaluate the antiderivative must be shown to earn the third point.
- The response cannot correct -16 to $+16$ in order to earn the third point; there is no possible reversal here.

Total for part (a) 3 points

(b)

It is known that, for $y = cx\sqrt{4-x^2}$, $\frac{dy}{dx} = \frac{c(4-2x^2)}{\sqrt{4-x^2}}$. For a particular spinning toy, the radius of the largest cross-sectional circular slice is 1.2 inches. What is the value of c for this spinning toy?

The cross-sectional circular slice with the largest radius occurs where $cx\sqrt{4-x^2}$ has its maximum on the interval $0 < x < 2$.

$$\frac{dy}{dx} = \frac{c(4-2x^2)}{\sqrt{4-x^2}} = 0 \Rightarrow x = \sqrt{2}$$

$$x = \sqrt{2} \Rightarrow y = c\sqrt{2}\sqrt{4-(\sqrt{2})^2} = 2c$$

$$2c = 1.2 \Rightarrow c = 0.6$$

Sets $\frac{dy}{dx} = 0$ **1 point**

Answer **1 point**

Scoring notes:

- The first point is earned for setting $\frac{dy}{dx} = 0$, $\frac{c(4-2x^2)}{\sqrt{4-x^2}} = 0$, or $c(4-2x^2) = 0$.
- An unsupported $x = \sqrt{2}$ does not earn the first point.
- The second point can be earned without the first point but is earned only for the answer $c = 0.6$ with supporting work.

Total for part (b) 2 points

(c) For another spinning toy, the volume is 2π cubic inches. What is the value of c for this spinning toy?

Volume $= \int_0^2 \pi (cx\sqrt{4-x^2})^2 dx = \pi c^2 \int_0^2 x^2 (4-x^2) dx$	Form of the integrand	1 point
$= \pi c^2 \int_0^2 (4x^2 - x^4) dx = \pi c^2 \left(\frac{4}{3}x^3 - \frac{1}{5}x^5 \right) \Big _0^2$	Limits and constant	1 point
$= \pi c^2 \left(\frac{32}{3} - \frac{32}{5} \right) = \frac{64\pi c^2}{15}$	Antiderivative	1 point
$\frac{64\pi c^2}{15} = 2\pi \Rightarrow c^2 = \frac{15}{32} \Rightarrow c = \sqrt{\frac{15}{32}}$	Answer	1 point

Scoring notes:

- The first point is earned for presenting an integrand of the form $A(x\sqrt{4-x^2})^2$ in a definite integral with any limits of integration (numeric or alphanumeric) and any nonzero constant A . Mishandling the c will result in the response being ineligible for the fourth point.
- The second point can be earned without the first point. The second point is earned for the limits of integration, $x = 0$ and $x = 2$, and the constant π (but not for 2π) as part of an integral with a correct or incorrect integrand.
- If an indefinite integral is presented with the correct constant π , the second point can be earned if the antiderivative (correct or incorrect) is evaluated using the correct limits of integration.
- A response that presents $2 = \int_0^2 (cx\sqrt{4-x^2})^2 dx$ earns the first and second points.
- The third point is earned for presenting a correct antiderivative of the presented integrand of the form $A(x\sqrt{4-x^2})^2$ for any nonzero A . If there are subsequent errors in simplification of the antiderivative, linkage errors, or sign errors, the response will not earn the fourth point.
- The fourth point cannot be earned without the third point. The fourth point is earned only for the correct answer. The expression does not need to be simplified to earn the fourth point.

Total for part (c) 4 points

Total for question 3 9 points

Part B (AB): Graphing calculator not allowed**Question 5****9 points****General Scoring Notes**

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Scoring guidelines and notes contain examples of the most common approaches seen in student responses. These guidelines can be applied to alternate approaches to ensure that these alternate approaches are scored appropriately.

Consider the function $y = f(x)$ whose curve is given by the equation $2y^2 - 6 = y \sin x$ for $y > 0$.

Model Solution	Scoring
(a) Show that $\frac{dy}{dx} = \frac{y \cos x}{4y - \sin x}$.	
$\frac{d}{dx}(2y^2 - 6) = \frac{d}{dx}(y \sin x) \Rightarrow 4y \frac{dy}{dx} = \frac{dy}{dx} \sin x + y \cos x$	Implicit differentiation 1 point
$\Rightarrow 4y \frac{dy}{dx} - \frac{dy}{dx} \sin x = y \cos x \Rightarrow \frac{dy}{dx}(4y - \sin x) = y \cos x$	Verification 1 point
$\Rightarrow \frac{dy}{dx} = \frac{y \cos x}{4y - \sin x}$	

Scoring notes:

- The first point is earned only for correctly implicitly differentiating $2y^2 - 6 = y \sin x$. Responses may use alternative notations for $\frac{dy}{dx}$, such as y' .
- The second point may not be earned without the first point.
- It is sufficient to present $\frac{dy}{dx}(4y - \sin x) = y \cos x$ to earn the second point, provided that there are no subsequent errors.

Total for part (a) 2 points

- (b) Write an equation for the line tangent to the curve at the point $(0, \sqrt{3})$.

At the point $(0, \sqrt{3})$, $\frac{dy}{dx} = \frac{\sqrt{3} \cos 0}{4\sqrt{3} - \sin 0} = \frac{1}{4}$.

Answer

1 point

An equation for the tangent line is $y = \sqrt{3} + \frac{1}{4}x$.

Scoring notes:

- Any correct tangent line equation will earn the point. No supporting work is required. Simplification of the slope value is not required.

Total for part (b) 1 point

- (c) For $0 \leq x \leq \pi$ and $y > 0$, find the coordinates of the point where the line tangent to the curve is horizontal.

$\frac{dy}{dx} = \frac{y \cos x}{4y - \sin x} = 0 \Rightarrow y \cos x = 0$ and $4y - \sin x \neq 0$

Sets $\frac{dy}{dx} = 0$

1 point

$y \cos x = 0$ and $y > 0 \Rightarrow x = \frac{\pi}{2}$

$x = \frac{\pi}{2}$

1 point

When $x = \frac{\pi}{2}$, $y \sin x = 2y^2 - 6 \Rightarrow y \sin \frac{\pi}{2} = 2y^2 - 6$

$y = 2$

1 point

$\Rightarrow y = 2y^2 - 6 \Rightarrow 2y^2 - y - 6 = 0$
 $\Rightarrow (2y + 3)(y - 2) = 0 \Rightarrow y = 2$

When $x = \frac{\pi}{2}$ and $y = 2$, $4y - \sin x = 8 - 1 \neq 0$. Therefore, the

line tangent to the curve is horizontal at the point $\left(\frac{\pi}{2}, 2\right)$.

Scoring notes:

- The first point is earned by any of $\frac{dy}{dx} = 0$, $\frac{y \cos x}{4y - \sin x} = 0$, $y \cos x = 0$, or $\cos x = 0$.
- If additional “correct” x -values are considered outside of the given domain, the response must commit to only $x = \frac{\pi}{2}$ to earn the second point. Any presented y -values, correct or incorrect, are not considered for the second point.
- Entering with $x = \frac{\pi}{2}$ does not earn the first point, earns the second point, and is eligible for the third point. The third point is earned for finding $y = 2$. The coordinates do not have to be presented as an ordered pair.
- The third point is not earned with additional points present unless the response commits to the correct point.

Total for part (c) 3 points

- (d) Determine whether f has a relative minimum, a relative maximum, or neither at the point found in part (c). Justify your answer.

$$\frac{d^2y}{dx^2} = \frac{(4y - \sin x)\left(\frac{dy}{dx}\cos x - y\sin x\right) - (y\cos x)\left(4\frac{dy}{dx} - \cos x\right)}{(4y - \sin x)^2}$$

Considers $\frac{d^2y}{dx^2}$ **1 point**

When $x = \frac{\pi}{2}$ and $y = 2$,

$$\frac{d^2y}{dx^2} = \frac{\left(4 \cdot 2 - \sin \frac{\pi}{2}\right)\left(0 \cdot \cos \frac{\pi}{2} - 2 \cdot \sin \frac{\pi}{2}\right) - \left(2 \cos \frac{\pi}{2}\right)\left(4 \cdot 0 - \cos \frac{\pi}{2}\right)}{\left(4 \cdot 2 - \sin \frac{\pi}{2}\right)^2}$$

$\frac{d^2y}{dx^2}$ at $\left(\frac{\pi}{2}, 2\right)$ **1 point**

$$= \frac{(7)(-2) - (0)(0)}{(7)^2} = \frac{-2}{7} < 0.$$

f has a relative maximum at the point $\left(\frac{\pi}{2}, 2\right)$ because $\frac{dy}{dx} = 0$ and $\frac{d^2y}{dx^2} < 0$. Answer with justification **1 point**

Scoring notes:

- The first point is earned for an attempt to use the quotient rule (or product rule) to find $\frac{d^2y}{dx^2}$.
- The second point is earned for correctly finding $\frac{d^2y}{dx^2}$ and evaluating to find that $\frac{d^2y}{dx^2} < 0$ at $\left(\frac{\pi}{2}, 2\right)$. The explicit value of $-\frac{2}{7}$ or the equivalent does not need to be reported, but any reported values must be correct in order to earn this point.
- The third point can be earned without the second point by reaching a consistent conclusion based on the reported sign of a nonzero value of $\frac{d^2y}{dx^2}$ obtained utilizing $\frac{dy}{dx} = 0$.
- Imports: A response is eligible to earn all 3 points in part (d) with a point of the form $\left(\frac{\pi}{2}, k\right)$ with $k > 0$, imported from part (c).

Alternate Solution for part (d)	Scoring for Alternate Solution
For the function $y = f(x)$ near the point $\left(\frac{\pi}{2}, 2\right)$, $4y - \sin x > 0$ and $y > 0$.	Considers sign of $4y - \sin x$ 1 point
Thus, $\frac{dy}{dx} = \frac{y \cos x}{4y - \sin x}$ changes from positive to negative at $x = \frac{\pi}{2}$.	$\frac{dy}{dx}$ changes from positive to negative at $x = \frac{\pi}{2}$ 1 point
By the First Derivative Test, f has a relative maximum at the point $\left(\frac{\pi}{2}, 2\right)$.	Conclusion 1 point

Scoring notes:

- The first point for considering the sign of $4y - \sin x$ may also be earned by stating that $4y - \sin x$ is not equal to zero.
- The second and third points can be earned without the first point.
- To earn the second point a response must state that $\frac{dy}{dx}$ (or $\cos x$) changes from positive to negative at $x = \frac{\pi}{2}$.
- The third point cannot be earned without the second point.
- A response that concludes there is a minimum at this point does not earn the third point.

Total for part (d) 3 points

Total for question 5 9 points