


Part A (AB or BC): Graphing calculator not allowed**Question 4****9 points****General Scoring Notes**

- The model solution is presented using standard mathematical notation.
- Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be accurate to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

The continuous function f is defined on the closed interval $-6 \leq x \leq 12$. The graph of f , consisting of two semicircles and one line segment, is shown in the figure.

Graph of f

Let g be the function defined by $g(x) = \int_6^x f(t) dt$.

	Model Solution	Scoring
A	Find $g'(8)$. Give a reason for your answer.	
	$g'(x) = f(x)$	Considers $g'(x) = f(x)$ Point 1 (P1)
	$g'(8) = f(8) = 1$	Answer Point 2 (P2)

Scoring Notes for Part A

- P1** is earned for $g' = f$, $g'(x) = f(x)$, or $g'(8) = f(8)$ in part A.
- A response of $g'(8) = f(8) = 1$ earns both **P1** and **P2**.
- A response that does not earn **P1** can earn **P2** with an implied application of the Fundamental Theorem of Calculus (e.g., $g'(8) = 1$ or $f(8) = 1$).
- A response of $g'(8) = f(8) - f(6) = 1$ earns **P2** but not **P1**.

- B** Find all values of x in the open interval $-6 < x < 12$ at which the graph of g has a point of inflection. Give a reason for your answer.

The graph of g has a point of inflection where $g'' = f'$ changes sign, which is where $g' = f$ changes from decreasing to increasing or vice versa.

The graph of g has points of inflection at $x = -3$ and $x = 6$ because f changes from decreasing to increasing there.

The graph of g also has a point of inflection at $x = 3$ because f changes from increasing to decreasing there.

Answer

Point 3 (P3)

Reason

Point 4 (P4)

Scoring Notes for Part B

- **P3** is earned only for an answer of $x = -3$, $x = 3$, and $x = 6$. If any other/additional values of x in $-6 < x < 12$ are declared to be points of inflection, the response does not earn either **P3** or **P4**. Consideration of $x = -6$ or of $x = 12$ does not impact scoring.
- To earn **P4**, a response must tie the reason to the given graph of f .
 - A response of “ g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because f changes from increasing to decreasing or decreasing to increasing there” earns both **P3** and **P4**.
 - A response of “ g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because the slope of f changes sign there” earns both **P3** and **P4**.
 - A response of “ g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because f attains relative extrema there” earns both **P3** and **P4**.
 - A response of “ g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because g changes concavity there” earns **P3** but not **P4**.
 - A response of “ g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because $g'' = f'$ changes sign there” earns **P3** but not **P4**.
 - A response that relies upon an ambiguous term such as “the function” or “the graph” does not earn **P4**.
- **Special case:** A response with two of the three correct x -values with correct reasoning and no other/additional values of x declared to be points of inflection earns **P4** but not **P3**.

C Find $g(12)$ and $g(0)$. Label your answers.

$$g(12) = \int_6^{12} f(t) \, dt = \frac{1}{2} \cdot 6 \cdot 3 = 9$$

$g(12)$

Point 5 (P5)

$$g(0) = \int_6^0 f(x) \, dx = -\int_0^6 f(x) \, dx = -\frac{\pi}{2} 3^2 = -\frac{9\pi}{2}$$

$g(0)$

Point 6 (P6)

Scoring Notes for Part C

- Unlabeled values do not earn either **P5** or **P6**.
- **P5** is earned for a response of $g(12) = 9$, with or without supporting work.
- **P6** is earned for a response of $g(0) = -\frac{9\pi}{2}$, with or without supporting work.

Note: Incorrect communication between the label “ $g(0)$ ” and the answer will be treated as scratch work and will not impact scoring. For example, $g(0) = \int_0^6 f(x) \, dx = -\frac{9\pi}{2}$ earns **P6**.

D Find the value of x at which g attains an absolute minimum on the closed interval $-6 \leq x \leq 12$. Justify your answer.											
For $-6 \leq x \leq 12$, g attains a minimum either when $g'(x) = f(x) = 0$ or at an endpoint. $g'(x) = f(x) = 0$	Considers $g'(x) = 0$ Point 7 (P7)										
$\Rightarrow x = 0, x = 6$ <table border="1" data-bbox="183 560 343 795"> <thead> <tr> <th>x</th> <th>$g(x)$</th> </tr> </thead> <tbody> <tr> <td>-6</td> <td>0</td> </tr> <tr> <td>0</td> <td>$-\frac{9\pi}{2}$</td> </tr> <tr> <td>6</td> <td>0</td> </tr> <tr> <td>12</td> <td>9</td> </tr> </tbody> </table>	x	$g(x)$	-6	0	0	$-\frac{9\pi}{2}$	6	0	12	9	Justification Point 8 (P8)
x	$g(x)$										
-6	0										
0	$-\frac{9\pi}{2}$										
6	0										
12	9										
Therefore, on the closed interval $-6 \leq x \leq 12$, g attains an absolute minimum value at $x = 0$.	Answer Point 9 (P9)										

Scoring Notes for Part D

- **P7** is earned for considering $g'(x) = 0$ or $f(x) = 0$. **P7** is not earned by just presenting $x = 0$ and $x = 6$.
A response that discusses the sign of $g'(x)$ or $f(x)$ changing OR uses the phrase “critical points of g ” also earns **P7**.
- To earn **P8** using a candidates test, a response must make a global argument by providing evaluations or reasoning for each of $g(-6)$, $g(0)$, $g(6)$, and $g(12)$ (and no other x -values).
- Alternate justification and answer:
Because $g'(x) \leq 0$ (or $f(x) \leq 0$) for $-6 \leq x < 0$ and $g'(x) \geq 0$ (or $f(x) \geq 0$) for $0 < x \leq 12$, the absolute minimum of g occurs at $x = 0$.
- A response that presents a local argument (such as a First Derivative Test or a Second Derivative Test) or an incorrect global argument does not earn **P8** but is eligible for **P9** with the correct answer of $x = 0$.
- For **P8**, values of $g(0)$ and $g(12)$ can be imported from part C. A response can earn **P9** with an answer that is consistent with the imported values.

Part A (AB or BC): Graphing calculator required**Question 1****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

t (seconds)	0	60	90	120	135	150
$f(t)$ (gallons per second)	0	0.1	0.15	0.1	0.05	0

A customer at a gas station is pumping gasoline into a gas tank. The rate of flow of gasoline is modeled by a differentiable function f , where $f(t)$ is measured in gallons per second and t is measured in seconds since pumping began. Selected values of $f(t)$ are given in the table.

Model Solution	Scoring
(a) Using correct units, interpret the meaning of $\int_{60}^{135} f(t) dt$ in the context of the problem. Use a right Riemann sum with the three subintervals $[60, 90]$, $[90, 120]$, and $[120, 135]$ to approximate the value of $\int_{60}^{135} f(t) dt$.	
$\int_{60}^{135} f(t) dt$ represents the total number of gallons of gasoline pumped into the gas tank from time $t = 60$ seconds to time $t = 135$ seconds.	Interpretation with units 1 point
$\int_{60}^{135} f(t) dt$ $\approx f(90)(90 - 60) + f(120)(120 - 90) + f(135)(135 - 120)$ $= (0.15)(30) + (0.1)(30) + (0.05)(15) = 8.25$	Form of Riemann sum 1 point
	Answer 1 point

Scoring notes:

- To earn the first point the response must reference gallons of gasoline added/pumped and the time interval $t = 60$ to $t = 135$.
- To earn the second point at least five of the six factors in the Riemann sum must be correct.
- If there is any error in the Riemann sum, the response does not earn the third point.
- A response of $(0.15)(30) + (0.1)(30) + (0.05)(15)$ earns both the second and third points, unless there is a subsequent error in simplification, in which case the response would earn only the second point.

- A response that presents a correct value with accompanying work that shows the three products in the Riemann sum but does not show all six of the factors and/or the sum process, does not earn the second point but does earn the third point. For example, responses of either $4.5 + 3.0 + 0.75$ or $(0.15)(30)$, $0.1(30)$, $0.05(15) \rightarrow 8.25$ earn the third point but not the second.
- A response of $f(90)(90 - 60) + f(120)(120 - 90) + f(135)(135 - 120) = 8.25$ earns both the second and the third points.
- A response that presents an answer of only 8.25 does not earn either the second or third point.
- A response that provides a completely correct left Riemann sum with accompanying work, $f(60)(30) + f(90)(30) + f(120)(15) = 9$, or $(0.1)(30) + (0.15)(30) + (0.1)(15)$ earns 1 of the last 2 points. A response with any errors or missing factors in a left Riemann sum earns neither of the last 2 points.

Total for part (a) 3 points**(b)** Must there exist a value of c , for $60 < c < 120$, such that $f'(c) = 0$? Justify your answer. f is differentiable. $\Rightarrow f$ is continuous on $[60, 120]$. $f(120) - f(60) = 0$ **1 point**

$$\frac{f(120) - f(60)}{120 - 60} = \frac{0.1 - 0.1}{60} = 0$$

Answer with justification **1 point**

By the Mean Value Theorem, there must exist a c , for $60 < c < 120$, such that $f'(c) = 0$.

Scoring notes:

- To earn the first point a response must present either $f(120) - f(60) = 0$, $0.1 - 0.1 = 0$ (perhaps as the numerator of a quotient), or $f(60) = f(120)$.
- To earn the second point a response must:
 - have earned the first point,
 - state that f is continuous because f is differentiable (or equivalent), and
 - answer “yes” in some way.
- A response may reference either the Mean Value Theorem or Rolle’s Theorem.
- A response that references the Intermediate Value Theorem cannot earn the second point.

Total for part (b) 2 points**(c)** The rate of flow of gasoline, in gallons per second, can also be modeled by

$$g(t) = \left(\frac{t}{500}\right) \cos\left(\left(\frac{t}{120}\right)^2\right) \text{ for } 0 \leq t \leq 150. \text{ Using this model, find the average rate of flow of}$$

gasoline over the time interval $0 \leq t \leq 150$. Show the setup for your calculations.

$$\frac{1}{150 - 0} \int_0^{150} g(t) dt$$

Average value formula **1 point**

$$= 0.0959967$$

Answer **1 point**

The average rate of flow of gasoline, in gallons per second, is 0.096 (or 0.095).

Scoring notes:

- The exact value of $\frac{1}{150} \int_0^{150} g(t) dt$ is $\frac{12}{125} \sin\left(\frac{25}{16}\right)$.
- A response may present the average value formula in single or multiple steps. For example, the following response earns both points: $\int_0^{150} g(t) dt = 14.399504$ so the average rate is 0.0959967.
- A response that presents the average value formula in multiple steps but provides incorrect or incomplete communication (e.g., $\int_0^{150} g(t) dt = \frac{14.399504}{150} = 0.0959967$) earns 1 out of 2 points.
- A response of $\int_0^{150} g(t) dt = 0.0959967$ does not earn either point.
- Degree mode: A response that presents answers obtained by using a calculator in degree mode does not earn the first point it would have otherwise earned. The response is generally eligible for all subsequent points (unless no answer is possible in degree mode or the question is made simpler by using degree mode). In degree mode, $\frac{1}{150} \int_0^{150} g(t) dt = 0.149981$ or 0.002618.

Total for part (c) 2 points

- (d) Using the model g defined in part (c), find the value of $g'(140)$. Interpret the meaning of your answer in the context of the problem.

$g'(140) \approx -0.004908$ $g'(140) = -0.005$ (or -0.004)	$g'(140)$ 1 point
The rate at which gasoline is flowing into the tank is decreasing at a rate of 0.005 (or 0.004) gallon per second per second at time $t = 140$ seconds.	Interpretation 1 point

Scoring notes:


- The exact value of $g'(140)$ is $\frac{1}{500} \cos\left(\frac{49}{36}\right) - \frac{49}{9000} \sin\left(\frac{49}{36}\right)$.
- The value of $g'(140)$ may appear only in the interpretation.
- To be eligible for the second point a response must present some numerical value for $g'(140)$.
- To earn the second point the interpretation must include “the rate of flow of gasoline is changing at a rate of [the declared value of $g'(140)$]” and “at $t = 140$ ” (or equivalent).
- An interpretation of “decreasing at a rate of -0.005 ” or “increasing at a rate of 0.005” does not earn the second point.
- Degree mode: In degree mode, $g'(140) = 0.001997$ or 0.00187.

Total for part (d) 2 points**Total for question 1 9 points**

Part B (AB or BC): Graphing calculator not allowed**Question 4****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Graph of f'

The function f is defined on the closed interval $[-2, 8]$ and satisfies $f(2) = 1$. The graph of f' , the derivative of f , consists of two line segments and a semicircle, as shown in the figure.

Model Solution**Scoring**

- (a) Does f have a relative minimum, a relative maximum, or neither at $x = 6$? Give a reason for your answer.

$f'(x) > 0$ on $(2, 6)$ and $f'(x) > 0$ on $(6, 8)$.

Answer with reason **1 point**

$f'(x)$ does not change sign at $x = 6$, so there is neither a relative maximum nor a relative minimum at this location.

Scoring notes:

- A response that declares $f'(x)$ does not change sign at $x = 6$, so neither, is sufficient to earn the point.
- A response does not have to present intervals on which $f'(x)$ is positive or negative, but if any are given, they must be correct. Any presented intervals may include none, one, or both endpoints.
- A response that declares $f'(x) > 0$ before and after $x = 6$ does not earn the point.

Total for part (a) 1 point

- (b) On what open intervals, if any, is the graph of f concave down? Give a reason for your answer.

The graph of f is concave down on $(-2, 0)$ and $(4, 6)$ because f' is decreasing on these intervals.	Intervals	1 point
	Reason	1 point

Scoring notes:

- The first point is earned only by an answer of $(-2, 0)$ and $(4, 6)$, or these intervals including one or both endpoints.
- A response must earn the first point to be eligible for second point.
- To earn the second point a response must correctly discuss the behavior of f' or the slopes of f' .
- Special case: A response that presents exactly one of the two correct intervals with a correct reason earns 1 out of 2 points.

Total for part (b) 2 points

- (c) Find the value of $\lim_{x \rightarrow 2} \frac{6f(x) - 3x}{x^2 - 5x + 6}$, or show that it does not exist. Justify your answer.

Because f is differentiable at $x = 2$, f is continuous at $x = 2$, so $\lim_{x \rightarrow 2} f(x) = f(2) = 1$. $\lim_{x \rightarrow 2} (6f(x) - 3x) = 6 \cdot 1 - 3 \cdot 2 = 0$ $\lim_{x \rightarrow 2} (x^2 - 5x + 6) = 0$	Limits of numerator and denominator	1 point
Because $\lim_{x \rightarrow 2} \frac{6f(x) - 3x}{x^2 - 5x + 6}$ is of indeterminate form $\frac{0}{0}$, L'Hospital's Rule can be applied. Using L'Hospital's Rule, $\lim_{x \rightarrow 2} \frac{6f(x) - 3x}{x^2 - 5x + 6} = \lim_{x \rightarrow 2} \frac{6f'(x) - 3}{2x - 5} = \frac{6 \cdot 0 - 3}{2 \cdot 2 - 5} = 3$.	Uses L'Hospital's Rule Answer	1 point 1 point

Scoring notes:

- The first point is earned by the presentation of two separate limits for the numerator and denominator.
- A response that presents a limit explicitly equal to $\frac{0}{0}$ does not earn the first point.
- The second point is earned by applying L'Hospital's Rule, that is, by presenting at least one correct derivative in the limit of a ratio of derivatives.
- The third point is earned for the correct answer with supporting work.

Total for part (c) 3 points

(d) Find the absolute minimum value of f on the closed interval $[-2, 8]$. Justify your answer.

$f'(x) = 0 \Rightarrow x = -1, x = 2, x = 6$	Considers $f'(x) = 0$	1 point												
The function f is continuous on $[-2, 8]$, so the candidates for the location of an absolute minimum for f are $x = -2, x = -1, x = 2, x = 6$, and $x = 8$.	Justification	1 point												
	Answer	1 point												
<table><tr><th>x</th><th>$f(x)$</th></tr><tr><td>-2</td><td>3</td></tr><tr><td>-1</td><td>4</td></tr><tr><td>2</td><td>1</td></tr><tr><td>6</td><td>$7 - \pi$</td></tr><tr><td>8</td><td>$11 - 2\pi$</td></tr></table>	x	$f(x)$	-2	3	-1	4	2	1	6	$7 - \pi$	8	$11 - 2\pi$		
x	$f(x)$													
-2	3													
-1	4													
2	1													
6	$7 - \pi$													
8	$11 - 2\pi$													
The absolute minimum value of f is $f(2) = 1$.														

Scoring notes:

- To earn the first point a response must state $f' = 0$ or equivalent. Listing the zeros of f' is not sufficient.
- A response that presents any error in evaluating f at any critical point or endpoint will not earn the justification point.
- A response need not present the value of $f(-1)$ provided $x = -1$ is eliminated because it is the location of a local maximum. A response need not present the value of $f(6)$ provided $x = 6$ is eliminated by reference to part (a) or eliminated through analysis.
- A response need not present the value of $f(8)$ provided there is a presentation that argues $f'(x) \geq 0$ for $x > 2$ and, therefore, $f(8) > f(2)$.
- A response that does not consider both endpoints does not earn the justification point.
- The answer point is earned only for indicating that the minimum value is 1. It is not earned for noting that the minimum occurs at $x = 2$.


Total for part (d) 3 points

Total for question 4 9 points

Part B (AB or BC): Graphing calculator not allowed**Question 4****9 points****General Scoring Notes**

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Scoring guidelines and notes contain examples of the most common approaches seen in student responses. These guidelines can be applied to alternate approaches to ensure that these alternate approaches are scored appropriately.

Graph of f

Let f be a continuous function defined on the closed interval $-4 \leq x \leq 6$. The graph of f , consisting of four line segments, is shown above. Let G be the function defined by $G(x) = \int_0^x f(t) dt$.

Model Solution	Scoring
$G'(x) = f(x)$ in any part of the response.	$G'(x) = f(x)$ 1 point

Scoring notes:

- This “global point” can be earned in any one part. Expressions that show this connection and therefore earn this point include: $G' = f$, $G'(x) = f(x)$, $G''(x) = f'(x)$ in part (a), $G'(3) = f(3)$ in part (b), or $G'(2) = f(2)$ in part (c).

Total 1 point

- (a) On what open intervals is the graph of G concave up? Give a reason for your answer.

$$G'(x) = f(x)$$

The graph of G is concave up for $-4 < x < -2$ and $2 < x < 6$, because $G' = f$ is increasing on these intervals.

Answer with reason

1 point

Scoring notes:

- Intervals may also include one or both endpoints.

Total for part (a)

1 point

- (b) Let P be the function defined by $P(x) = G(x) \cdot f(x)$. Find $P'(3)$.

$$P'(x) = G(x) \cdot f'(x) + f(x) \cdot G'(x)$$

$$P'(3) = G(3) \cdot f'(3) + f(3) \cdot G'(3)$$

Product rule

1 point

Substituting $G(3) = \int_0^3 f(t) dt = -3.5$ and $G'(3) = f(3) = -3$ into the above expression for $P'(3)$ gives the following:

$G(3)$ or $G'(3)$

1 point

$$P'(3) = -3.5 \cdot 1 + (-3) \cdot (-3) = 5.5$$

Answer

1 point

Scoring notes:

- The first point is earned for the correct application of the product rule in terms of x or in the evaluation of $P'(3)$. Once earned, this point cannot be lost.
- The second point is earned by correctly evaluating $G(3) = -3.5$, $G'(3) = -3$, or $f(3) = -3$.
- To be eligible to earn the third point a response must have earned the first two points.
- Simplification of the numerical value is not required to earn the third point.

Total for part (b)

3 points

(c) Find $\lim_{x \rightarrow 2} \frac{G(x)}{x^2 - 2x}$.

$$\lim_{x \rightarrow 2} (x^2 - 2x) = 0$$

Because G is continuous for $-4 \leq x \leq 6$,

$$\lim_{x \rightarrow 2} G(x) = \int_0^2 f(t) dt = 0.$$

Therefore, the limit $\lim_{x \rightarrow 2} \frac{G(x)}{x^2 - 2x}$ is an indeterminate form of type $\frac{0}{0}$.

Uses L'Hospital's Rule

1 point

Using L'Hospital's Rule,

$$\begin{aligned} \lim_{x \rightarrow 2} \frac{G(x)}{x^2 - 2x} &= \lim_{x \rightarrow 2} \frac{G'(x)}{2x - 2} \\ &= \lim_{x \rightarrow 2} \frac{f(x)}{2x - 2} = \frac{f(2)}{2} = \frac{-4}{2} = -2 \end{aligned}$$

Answer with justification

1 point

Scoring notes:

- To earn the first point the response must show $\lim_{x \rightarrow 2} (x^2 - 2x) = 0$ and $\lim_{x \rightarrow 2} G(x) = 0$ and must show a ratio of the two derivatives, $G'(x)$ and $2x - 2$. The ratio may be shown as evaluations of the derivatives at $x = 2$, such as $\frac{G'(2)}{2}$.
- To earn the second point the response must evaluate correctly with appropriate limit notation. In particular the response must include $\lim_{x \rightarrow 2} \frac{G'(x)}{2x - 2}$ or $\lim_{x \rightarrow 2} \frac{f(x)}{2x - 2}$.
- With any linkage errors (such as $\frac{G'(x)}{2x - 2} = \frac{f(2)}{2}$), the response does not earn the second point.

Total for part (c) 2 points

- (d) Find the average rate of change of G on the interval $[-4, 2]$. Does the Mean Value Theorem guarantee a value c , $-4 < c < 2$, for which $G'(c)$ is equal to this average rate of change? Justify your answer.

$$G(2) = \int_0^2 f(t) dt = 0 \text{ and } G(-4) = \int_0^{-4} f(t) dt = -16$$

Average rate of change **1 point**

$$\text{Average rate of change} = \frac{G(2) - G(-4)}{2 - (-4)} = \frac{0 - (-16)}{6} = \frac{8}{3}$$

Yes, $G'(x) = f(x)$ so G is differentiable on $(-4, 2)$ and continuous on $[-4, 2]$. Therefore, the Mean Value Theorem applies and guarantees a value c , $-4 < c < 2$, such that

$$G'(c) = \frac{8}{3}.$$

Answer with justification **1 point**

Scoring notes:

- To earn the first point a response must present at least a difference and a quotient and a correct evaluation. For example, $\frac{0 + 16}{6}$ or $\frac{G(2) - G(-4)}{6} = \frac{16}{6}$.
- Simplification of the numerical value is not required to earn the first point, but any simplification must be correct.
- The second point can be earned without the first point if the response has the correct setup but an incorrect or no evaluation of the average rate of change. The Mean Value Theorem need not be explicitly stated provided the conditions and conclusion are stated.

Total for part (d) 2 points

Total for question 4 9 points

2018

AP

CollegeBoard

AP Calculus AB

Scoring Guidelines

2018 SCORING GUIDELINES

Question 1

(a) $\int_0^{300} r(t) dt = 270$

According to the model, 270 people enter the line for the escalator during the time interval $0 \leq t \leq 300$.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) $20 + \int_0^{300} (r(t) - 0.7) dt = 20 + \int_0^{300} r(t) dt - 0.7 \cdot 300 = 80$

According to the model, 80 people are in line at time $t = 300$.

2 : $\begin{cases} 1 : \text{considers rate out} \\ 1 : \text{answer} \end{cases}$

(c) Based on part (b), the number of people in line at time $t = 300$ is 80.

The first time t that there are no people in line is

$$300 + \frac{80}{0.7} = 414.286 \text{ (or 414.285) seconds.}$$

1 : answer

(d) The total number of people in line at time t , $0 \leq t \leq 300$, is modeled by

$$20 + \int_0^t r(x) dx - 0.7t.$$

$$r(t) - 0.7 = 0 \Rightarrow t_1 = 33.013298, t_2 = 166.574719$$

4 : $\begin{cases} 1 : \text{considers } r(t) - 0.7 = 0 \\ 1 : \text{identifies } t = 33.013 \\ 1 : \text{answers} \\ 1 : \text{justification} \end{cases}$

t	People in line for escalator
0	20
t_1	3.803
t_2	158.070
300	80

The number of people in line is a minimum at time $t = 33.013$ seconds, when there are 4 people in line.

2018 SCORING GUIDELINES

Question 2

(a) $v'(3) = -2.118$

The acceleration of the particle at time $t = 3$ is -2.118 .

1 : answer

(b) $x(3) = x(0) + \int_0^3 v(t) dt = -5 + \int_0^3 v(t) dt = -1.760213$

The position of the particle at time $t = 3$ is -1.760 .

3 : $\begin{cases} 1 : \int_0^3 v(t) dt \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

(c) $\int_0^{3.5} v(t) dt = 2.844$ (or 2.843)

$$\int_0^{3.5} |v(t)| dt = 3.737$$

The integral $\int_0^{3.5} v(t) dt$ is the displacement of the particle over the time interval $0 \leq t \leq 3.5$.

The integral $\int_0^{3.5} |v(t)| dt$ is the total distance traveled by the particle over the time interval $0 \leq t \leq 3.5$.

3 : $\begin{cases} 1 : \text{answers} \\ 2 : \text{interpretations of } \int_0^{3.5} v(t) dt \\ \text{and } \int_0^{3.5} |v(t)| dt \end{cases}$

(d) $v(t) = x_2'(t)$

$$v(t) = 2t - 1 \Rightarrow t = 1.57054$$

The two particles are moving with the same velocity at time $t = 1.571$ (or 1.570).

2 : $\begin{cases} 1 : \text{sets } v(t) = x_2'(t) \\ 1 : \text{answer} \end{cases}$

2018 SCORING GUIDELINES

Question 3

$$\begin{aligned} \text{(a)} \quad f(-5) &= f(1) + \int_1^{-5} g(x) \, dx = f(1) - \int_{-5}^1 g(x) \, dx \\ &= 3 - \left(-9 - \frac{3}{2} + 1 \right) = 3 - \left(-\frac{19}{2} \right) = \frac{25}{2} \end{aligned}$$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b)} \quad \int_1^6 g(x) \, dx &= \int_1^3 g(x) \, dx + \int_3^6 g(x) \, dx \\ &= \int_1^3 2 \, dx + \int_3^6 2(x-4)^2 \, dx \\ &= 4 + \left[\frac{2}{3}(x-4)^3 \right]_{x=3}^{x=6} = 4 + \frac{16}{3} - \left(-\frac{2}{3} \right) = 10 \end{aligned}$$

3 : $\begin{cases} 1 : \text{split at } x = 3 \\ 1 : \text{antiderivative of } 2(x-4)^2 \\ 1 : \text{answer} \end{cases}$

(c) The graph of f is increasing and concave up on $0 < x < 1$ and $4 < x < 6$ because $f'(x) = g(x) > 0$ and $f''(x) = g'(x)$ is increasing on those intervals.

2 : $\begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$

(d) The graph of f has a point of inflection at $x = 4$ because $f'(x) = g(x)$ changes from decreasing to increasing at $x = 4$.

2 : $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

2018 SCORING GUIDELINES

Question 4

$$(a) \quad H'(6) \approx \frac{H(7) - H(5)}{7 - 5} = \frac{11 - 6}{2} = \frac{5}{2}$$

$H'(6)$ is the rate at which the height of the tree is changing, in meters per year, at time $t = 6$ years.

$$(b) \quad \frac{H(5) - H(3)}{5 - 3} = \frac{6 - 2}{2} = 2$$

Because H is differentiable on $3 \leq t \leq 5$, H is continuous on $3 \leq t \leq 5$.

By the Mean Value Theorem, there exists a value c , $3 < c < 5$, such that $H'(c) = 2$.

$$(c) \quad \text{The average height of the tree over the time interval } 2 \leq t \leq 10 \text{ is given by } \frac{1}{10 - 2} \int_2^{10} H(t) \, dt.$$

$$\begin{aligned} \frac{1}{8} \int_2^{10} H(t) \, dt &\approx \frac{1}{8} \left(\frac{1.5 + 2}{2} \cdot 1 + \frac{2 + 6}{2} \cdot 2 + \frac{6 + 11}{2} \cdot 2 + \frac{11 + 15}{2} \cdot 3 \right) \\ &= \frac{1}{8} (65.75) = \frac{263}{32} \end{aligned}$$

The average height of the tree over the time interval $2 \leq t \leq 10$ is $\frac{263}{32}$ meters.

$$(d) \quad G(x) = 50 \Rightarrow x = 1$$

$$\frac{d}{dt}(G(x)) = \frac{d}{dx}(G(x)) \cdot \frac{dx}{dt} = \frac{(1+x)100 - 100x \cdot 1}{(1+x)^2} \cdot \frac{dx}{dt} = \frac{100}{(1+x)^2} \cdot \frac{dx}{dt}$$

$$\left. \frac{d}{dt}(G(x)) \right|_{x=1} = \frac{100}{(1+1)^2} \cdot 0.03 = \frac{3}{4}$$

According to the model, the rate of change of the height of the tree with respect to time when the tree is 50 meters tall is $\frac{3}{4}$ meter per year.

$$2 : \begin{cases} 1 : \text{estimate} \\ 1 : \text{interpretation with units} \end{cases}$$

$$2 : \begin{cases} 1 : \frac{H(5) - H(3)}{5 - 3} \\ 1 : \text{conclusion using Mean Value Theorem} \end{cases}$$

$$2 : \begin{cases} 1 : \text{trapezoidal sum} \\ 1 : \text{approximation} \end{cases}$$

$$3 : \begin{cases} 2 : \frac{d}{dt}(G(x)) \\ 1 : \text{answer} \end{cases}$$

Note: max 1/3 [1-0] if no chain rule

2018 SCORING GUIDELINES

Question 5

- (a) The average rate of change of
- f
- on the interval
- $0 \leq x \leq \pi$
- is

$$\frac{f(\pi) - f(0)}{\pi - 0} = \frac{-e^\pi - 1}{\pi}.$$

1 : answer

- (b)
- $f'(x) = e^x \cos x - e^x \sin x$

$$f'\left(\frac{3\pi}{2}\right) = e^{3\pi/2} \cos\left(\frac{3\pi}{2}\right) - e^{3\pi/2} \sin\left(\frac{3\pi}{2}\right) = e^{3\pi/2}$$

2 : $\begin{cases} 1 : f'(x) \\ 1 : \text{slope} \end{cases}$

The slope of the line tangent to the graph of f at $x = \frac{3\pi}{2}$ is $e^{3\pi/2}$.

- (c)
- $f'(x) = 0 \Rightarrow \cos x - \sin x = 0 \Rightarrow x = \frac{\pi}{4}, x = \frac{5\pi}{4}$

x	$f(x)$
0	1
$\frac{\pi}{4}$	$\frac{1}{\sqrt{2}}e^{\pi/4}$
$\frac{5\pi}{4}$	$-\frac{1}{\sqrt{2}}e^{5\pi/4}$
2π	$e^{2\pi}$

3 : $\begin{cases} 1 : \text{sets } f'(x) = 0 \\ 1 : \text{identifies } x = \frac{\pi}{4}, x = \frac{5\pi}{4} \\ \text{as candidates} \\ 1 : \text{answer with justification} \end{cases}$

The absolute minimum value of f on $0 \leq x \leq 2\pi$ is $-\frac{1}{\sqrt{2}}e^{5\pi/4}$.

- (d)
- $\lim_{x \rightarrow \pi/2} f(x) = 0$

Because g is differentiable, g is continuous.

$$\lim_{x \rightarrow \pi/2} g(x) = g\left(\frac{\pi}{2}\right) = 0$$

By L'Hospital's Rule,

$$\lim_{x \rightarrow \pi/2} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \pi/2} \frac{f'(x)}{g'(x)} = \frac{-e^{\pi/2}}{2}.$$


3 : $\begin{cases} 1 : g \text{ is continuous at } x = \frac{\pi}{2} \\ \text{and limits equal 0} \\ 1 : \text{applies L'Hospital's Rule} \\ 1 : \text{answer} \end{cases}$

Note: max 1/3 [1-0-0] if no limit notation attached to a ratio of derivatives

2018 SCORING GUIDELINES

Question 6

(a)



$$2 : \begin{cases} 1 : \text{solution curve through } (0, 2) \\ 1 : \text{solution curve through } (1, 0) \end{cases}$$

Curves must go through the indicated points, follow the given slope lines, and extend to the boundary of the slope field.

$$(b) \left. \frac{dy}{dx} \right|_{(x, y)=(1, 0)} = \frac{4}{3}$$

An equation for the line tangent to the graph of $y = f(x)$ at

$$x = 1 \text{ is } y = \frac{4}{3}(x - 1).$$

$$f(0.7) \approx \frac{4}{3}(0.7 - 1) = -0.4$$

$$(c) \frac{dy}{dx} = \frac{1}{3}x(y - 2)^2$$

$$\int \frac{dy}{(y - 2)^2} = \int \frac{1}{3}x \, dx$$

$$\frac{-1}{y - 2} = \frac{1}{6}x^2 + C$$

$$\frac{1}{2} = \frac{1}{6} + C \Rightarrow C = \frac{1}{3}$$

$$\frac{-1}{y - 2} = \frac{1}{6}x^2 + \frac{1}{3} = \frac{x^2 + 2}{6}$$

$$y = 2 - \frac{6}{x^2 + 2}$$

Note: this solution is valid for $-\infty < x < \infty$.

$$2 : \begin{cases} 1 : \text{equation of tangent line} \\ 1 : \text{approximation} \end{cases}$$

$$5 : \begin{cases} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ \text{and uses initial condition} \\ 1 : \text{solves for } y \end{cases}$$

Note: 0/5 if no separation of variables

Note: max 3/5 [1-2-0-0] if no constant of integration

AP[®] Calculus AB

2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

0

2016 SCORING GUIDELINES

Question 1

t (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.

- (a) Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.
- (b) Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
- (c) Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
- (d) For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

$$(a) \quad R'(2) \approx \frac{R(3) - R(1)}{3 - 1} = \frac{950 - 1190}{3 - 1} = -120 \text{ liters/hr}^2$$

$$2 : \begin{cases} 1 : \text{estimate} \\ 1 : \text{units} \end{cases}$$

(b) The total amount of water removed is given by $\int_0^8 R(t) dt$.

$$\begin{aligned} \int_0^8 R(t) dt &\approx 1 \cdot R(0) + 2 \cdot R(1) + 3 \cdot R(3) + 2 \cdot R(6) \\ &= 1(1340) + 2(1190) + 3(950) + 2(740) \\ &= 8050 \text{ liters} \end{aligned}$$

This is an overestimate since R is a decreasing function.

$$3 : \begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{estimate} \\ 1 : \text{overestimate with reason} \end{cases}$$

$$\begin{aligned} (c) \quad \text{Total} &\approx 50000 + \int_0^8 W(t) dt - 8050 \\ &= 50000 + 7836.195325 - 8050 \approx 49786 \text{ liters} \end{aligned}$$

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{estimate} \end{cases}$$

(d) $W(0) - R(0) > 0$, $W(8) - R(8) < 0$, and $W(t) - R(t)$ is continuous.

$$2 : \begin{cases} 1 : \text{considers } W(t) - R(t) \\ 1 : \text{answer with explanation} \end{cases}$$

Therefore, the Intermediate Value Theorem guarantees at least one time t , $0 < t < 8$, for which $W(t) - R(t) = 0$, or $W(t) = R(t)$.

For this value of t , the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank.

2016 SCORING GUIDELINES

Question 2

For $t \geq 0$, a particle moves along the x -axis. The velocity of the particle at time t is given by

$v(t) = 1 + 2\sin\left(\frac{t^2}{2}\right)$. The particle is at position $x = 2$ at time $t = 4$.

- At time $t = 4$, is the particle speeding up or slowing down?
- Find all times t in the interval $0 < t < 3$ when the particle changes direction. Justify your answer.
- Find the position of the particle at time $t = 0$.
- Find the total distance the particle travels from time $t = 0$ to time $t = 3$.

(a) $v(4) = 2.978716 > 0$

$v'(4) = -1.164000 < 0$

The particle is slowing down since the velocity and acceleration have different signs.

(b) $v(t) = 0 \Rightarrow t = 2.707468$

$v(t)$ changes from positive to negative at $t = 2.707$.

Therefore, the particle changes direction at this time.

(c) $x(0) = x(4) + \int_4^0 v(t) dt$

$= 2 + (-5.815027) = -3.815$

(d) Distance $= \int_0^3 |v(t)| dt = 5.301$

2 : conclusion with reason

2 : $\begin{cases} 1 : t = 2.707 \\ 1 : \text{justification} \end{cases}$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$


2016 SCORING GUIDELINES

Question 3

The figure above shows the graph of the piecewise-linear function f . For $-4 \leq x \leq 12$, the function g is defined by

$$g(x) = \int_2^x f(t) \, dt.$$

- (a) Does g have a relative minimum, a relative maximum, or neither at $x = 10$? Justify your answer.
- (b) Does the graph of g have a point of inflection at $x = 4$? Justify your answer.
- (c) Find the absolute minimum value and the absolute maximum value of g on the interval $-4 \leq x \leq 12$. Justify your answers.
- (d) For $-4 \leq x \leq 12$, find all intervals for which $g(x) \leq 0$.

Graph of f

- (a) The function g has neither a relative minimum nor a relative maximum at $x = 10$ since $g'(x) = f(x)$ and $f(x) \leq 0$ for $8 \leq x \leq 12$.
- (b) The graph of g has a point of inflection at $x = 4$ since $g'(x) = f(x)$ is increasing for $2 \leq x \leq 4$ and decreasing for $4 \leq x \leq 8$.
- (c) $g'(x) = f(x)$ changes sign only at $x = -2$ and $x = 6$.

x	$g(x)$
-4	-4
-2	-8
6	8
12	-4

On the interval $-4 \leq x \leq 12$, the absolute minimum value is $g(-2) = -8$ and the absolute maximum value is $g(6) = 8$.

- (d) $g(x) \leq 0$ for $-4 \leq x \leq 2$ and $10 \leq x \leq 12$.

1 : $g'(x) = f(x)$ in (a), (b), (c), or (d)

1 : answer with justification

1 : answer with justification

4 : $\left\{ \begin{array}{l} 1 : \text{considers } x = -2 \text{ and } x = 6 \\ \quad \text{as candidates} \\ 1 : \text{considers } x = -4 \text{ and } x = 12 \\ 2 : \text{answers with justification} \end{array} \right.$


2 : intervals

2016 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = \frac{y^2}{x-1}$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.
- (b) Let $y = f(x)$ be the particular solution to the given differential equation with the initial condition $f(2) = 3$. Write an equation for the line tangent to the graph of $y = f(x)$ at $x = 2$.
Use your equation to approximate $f(2.1)$.
- (c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(2) = 3$.



2 : $\begin{cases} 1 : \text{zero slopes} \\ 1 : \text{nonzero slopes} \end{cases}$

(b) $\left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} = \frac{3^2}{2-1} = 9$

An equation for the tangent line is $y = 9(x - 2) + 3$.

$$f(2.1) \approx 9(2.1 - 2) + 3 = 3.9$$

2 : $\begin{cases} 1 : \text{tangent line equation} \\ 1 : \text{approximation} \end{cases}$

(c) $\frac{1}{y^2} dy = \frac{1}{x-1} dx$
 $\int \frac{1}{y^2} dy = \int \frac{1}{x-1} dx$
 $-\frac{1}{y} = \ln|x-1| + C$
 $-\frac{1}{3} = \ln|2-1| + C \Rightarrow C = -\frac{1}{3}$
 $-\frac{1}{y} = \ln|x-1| - \frac{1}{3}$
 $y = \frac{1}{\frac{1}{3} - \ln(x-1)}$

5 : $\begin{cases} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration and} \\ \quad \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$


Note: max 3/5 [1-2-0-0] if no constant of integration

Note: 0/5 if no separation of variables

Note: This solution is valid for $1 < x < 1 + e^{1/3}$.

2016 SCORING GUIDELINES

Question 5



The inside of a funnel of height 10 inches has circular cross sections, as shown in the figure above. At height h , the radius of the funnel is given by $r = \frac{1}{20}(3 + h^2)$, where $0 \leq h \leq 10$. The units of r and h are inches.

- (a) Find the average value of the radius of the funnel.
- (b) Find the volume of the funnel.
- (c) The funnel contains liquid that is draining from the bottom. At the instant when the height of the liquid is $h = 3$ inches, the radius of the surface of the liquid is decreasing at a rate of $\frac{1}{5}$ inch per second. At this instant, what is the rate of change of the height of the liquid with respect to time?

$$\begin{aligned} \text{(a) Average radius} &= \frac{1}{10} \int_0^{10} \frac{1}{20}(3 + h^2) dh = \frac{1}{200} \left[3h + \frac{h^3}{3} \right]_0^{10} \\ &= \frac{1}{200} \left(\left(30 + \frac{1000}{3} \right) - 0 \right) = \frac{109}{60} \text{ in} \end{aligned}$$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b) Volume} &= \pi \int_0^{10} \left(\left(\frac{1}{20} \right) (3 + h^2) \right)^2 dh = \frac{\pi}{400} \int_0^{10} (9 + 6h^2 + h^4) dh \\ &= \frac{\pi}{400} \left[9h + 2h^3 + \frac{h^5}{5} \right]_0^{10} \\ &= \frac{\pi}{400} \left(\left(90 + 2000 + \frac{100000}{5} \right) - 0 \right) = \frac{2209\pi}{40} \text{ in}^3 \end{aligned}$$

3 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(c) } \frac{dr}{dt} &= \frac{1}{20}(2h) \frac{dh}{dt} \\ -\frac{1}{5} &= \frac{3}{10} \frac{dh}{dt} \\ \frac{dh}{dt} &= -\frac{1}{5} \cdot \frac{10}{3} = -\frac{2}{3} \text{ in/sec} \end{aligned}$$

3 : $\begin{cases} 2 : \text{chain rule} \\ 1 : \text{answer} \end{cases}$

2016 SCORING GUIDELINES

Question 6

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	-6	3	2	8
2	2	-2	-3	0
3	8	7	6	2
6	4	5	3	-1

The functions f and g have continuous second derivatives. The table above gives values of the functions and their derivatives at selected values of x .

(a) Let $k(x) = f(g(x))$. Write an equation for the line tangent to the graph of k at $x = 3$.

(b) Let $h(x) = \frac{g(x)}{f(x)}$. Find $h'(1)$.

(c) Evaluate $\int_1^3 f''(2x) dx$.

$$\begin{aligned} \text{(a)} \quad k(3) &= f(g(3)) = f(6) = 4 \\ k'(3) &= f'(g(3)) \cdot g'(3) = f'(6) \cdot 2 = 5 \cdot 2 = 10 \end{aligned}$$

An equation for the tangent line is $y = 10(x - 3) + 4$.

$$\begin{aligned} \text{(b)} \quad h'(1) &= \frac{f(1) \cdot g'(1) - g(1) \cdot f'(1)}{(f(1))^2} \\ &= \frac{(-6) \cdot 8 - 2 \cdot 3}{(-6)^2} = \frac{-54}{36} = -\frac{3}{2} \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad \int_1^3 f''(2x) dx &= \frac{1}{2} [f'(2x)]_1^3 = \frac{1}{2} [f'(6) - f'(2)] \\ &= \frac{1}{2} [5 - (-2)] = \frac{7}{2} \end{aligned}$$

3 : $\begin{cases} 2 : \text{slope at } x = 3 \\ 1 : \text{equation for tangent line} \end{cases}$

3 : $\begin{cases} 2 : \text{expression for } h'(1) \\ 1 : \text{answer} \end{cases}$

3 : $\begin{cases} 2 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

AP[®] Calculus AB

2014 Scoring Guidelines

© 2014 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

0

2014 SCORING GUIDELINES

Question 1

Grass clippings are placed in a bin, where they decompose. For $0 \leq t \leq 30$, the amount of grass clippings remaining in the bin is modeled by $A(t) = 6.687(0.931)^t$, where $A(t)$ is measured in pounds and t is measured in days.

- Find the average rate of change of $A(t)$ over the interval $0 \leq t \leq 30$. Indicate units of measure.
- Find the value of $A'(15)$. Using correct units, interpret the meaning of the value in the context of the problem.
- Find the time t for which the amount of grass clippings in the bin is equal to the average amount of grass clippings in the bin over the interval $0 \leq t \leq 30$.
- For $t > 30$, $L(t)$, the linear approximation to A at $t = 30$, is a better model for the amount of grass clippings remaining in the bin. Use $L(t)$ to predict the time at which there will be 0.5 pound of grass clippings remaining in the bin. Show the work that leads to your answer.

$$(a) \frac{A(30) - A(0)}{30 - 0} = -0.197 \text{ (or } -0.196) \text{ lbs/day}$$

1 : answer with units

$$(b) A'(15) = -0.164 \text{ (or } -0.163)$$

The amount of grass clippings in the bin is decreasing at a rate of 0.164 (or 0.163) lbs/day at time $t = 15$ days.

2 : $\begin{cases} 1 : A'(15) \\ 1 : \text{interpretation} \end{cases}$

$$(c) A(t) = \frac{1}{30} \int_0^{30} A(t) dt \Rightarrow t = 12.415 \text{ (or } 12.414)$$

2 : $\begin{cases} 1 : \frac{1}{30} \int_0^{30} A(t) dt \\ 1 : \text{answer} \end{cases}$

$$(d) L(t) = A(30) + A'(30) \cdot (t - 30)$$

$$A'(30) = -0.055976$$

$$A(30) = 0.782928$$

$$L(t) = 0.5 \Rightarrow t = 35.054$$


4 : $\begin{cases} 2 : \text{expression for } L(t) \\ 1 : L(t) = 0.5 \\ 1 : \text{answer} \end{cases}$

2014 SCORING GUIDELINES

Question 2

Let R be the region enclosed by the graph of $f(x) = x^4 - 2.3x^3 + 4$ and the horizontal line $y = 4$, as shown in the figure above.

- (a) Find the volume of the solid generated when R is rotated about the horizontal line $y = -2$.
- (b) Region R is the base of a solid. For this solid, each cross section perpendicular to the x -axis is an isosceles right triangle with a leg in R . Find the volume of the solid.
- (c) The vertical line $x = k$ divides R into two regions with equal areas. Write, but do not solve, an equation involving integral expressions whose solution gives the value k .



(a) $f(x) = 4 \Rightarrow x = 0, 2.3$

$$\begin{aligned} \text{Volume} &= \pi \int_0^{2.3} \left[(4 + 2)^2 - (f(x) + 2)^2 \right] dx \\ &= 98.868 \text{ (or } 98.867) \end{aligned}$$

$$4 : \begin{cases} 2 : \text{integrand} \\ 1 : \text{limits} \\ 1 : \text{answer} \end{cases}$$

(b) $\text{Volume} = \int_0^{2.3} \frac{1}{2} (4 - f(x))^2 dx$
 $= 3.574 \text{ (or } 3.573)$

$$3 : \begin{cases} 2 : \text{integrand} \\ 1 : \text{answer} \end{cases}$$

(c) $\int_0^k (4 - f(x)) dx = \int_k^{2.3} (4 - f(x)) dx$

$$2 : \begin{cases} 1 : \text{area of one region} \\ 1 : \text{equation} \end{cases}$$


2014 SCORING GUIDELINES

Question 3

The function f is defined on the closed interval $[-5, 4]$. The graph of f consists of three line segments and is shown in the figure above.

Let g be the function defined by $g(x) = \int_{-3}^x f(t) dt$.

- (a) Find $g(3)$.
- (b) On what open intervals contained in $-5 < x < 4$ is the graph of g both increasing and concave down? Give a reason for your answer.
- (c) The function h is defined by $h(x) = \frac{g(x)}{5x}$. Find $h'(3)$.
- (d) The function p is defined by $p(x) = f(x^2 - x)$. Find the slope of the line tangent to the graph of p at the point where $x = -1$.

Graph of f

(a) $g(3) = \int_{-3}^3 f(t) dt = 6 + 4 - 1 = 9$

1 : answer

(b) $g'(x) = f(x)$

The graph of g is increasing and concave down on the intervals $-5 < x < -3$ and $0 < x < 2$ because $g' = f$ is positive and decreasing on these intervals.

2 : $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

(c) $h'(x) = \frac{5xg'(x) - g(x)5}{(5x)^2} = \frac{5xg'(x) - 5g(x)}{25x^2}$

3 : $\begin{cases} 2 : h'(x) \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} h'(3) &= \frac{(5)(3)g'(3) - 5g(3)}{25 \cdot 3^2} \\ &= \frac{15(-2) - 5(9)}{225} = \frac{-75}{225} = -\frac{1}{3} \end{aligned}$$

(d) $p'(x) = f'(x^2 - x)(2x - 1)$

3 : $\begin{cases} 2 : p'(x) \\ 1 : \text{answer} \end{cases}$

$$p'(-1) = f'(2)(-3) = (-2)(-3) = 6$$

2014 SCORING GUIDELINES

Question 4

Train A runs back and forth on an east-west section of railroad track. Train A 's velocity, measured in meters per minute, is given by a differentiable function $v_A(t)$, where time t is measured in minutes. Selected values for $v_A(t)$ are given in the table above.

t (minutes)	0	2	5	8	12
$v_A(t)$ (meters/minute)	0	100	40	-120	-150

- (a) Find the average acceleration of train A over the interval $2 \leq t \leq 8$.
- (b) Do the data in the table support the conclusion that train A 's velocity is -100 meters per minute at some time t with $5 < t < 8$? Give a reason for your answer.
- (c) At time $t = 2$, train A 's position is 300 meters east of the Origin Station, and the train is moving to the east. Write an expression involving an integral that gives the position of train A , in meters from the Origin Station, at time $t = 12$. Use a trapezoidal sum with three subintervals indicated by the table to approximate the position of the train at time $t = 12$.
- (d) A second train, train B , travels north from the Origin Station. At time t the velocity of train B is given by $v_B(t) = -5t^2 + 60t + 25$, and at time $t = 2$ the train is 400 meters north of the station. Find the rate, in meters per minute, at which the distance between train A and train B is changing at time $t = 2$.

(a) average accel = $\frac{v_A(8) - v_A(2)}{8 - 2} = \frac{-120 - 100}{6} = -\frac{110}{3} \text{ m/min}^2$

(b) v_A is differentiable $\Rightarrow v_A$ is continuous
 $v_A(8) = -120 < -100 < 40 = v_A(5)$

Therefore, by the Intermediate Value Theorem, there is a time t , $5 < t < 8$, such that $v_A(t) = -100$.

(c) $s_A(12) = s_A(2) + \int_2^{12} v_A(t) dt = 300 + \int_2^{12} v_A(t) dt$
 $\int_2^{12} v_A(t) dt \approx 3 \cdot \frac{100 + 40}{2} + 3 \cdot \frac{40 - 120}{2} + 4 \cdot \frac{-120 - 150}{2}$
 $= -450$

$$s_A(12) \approx 300 - 450 = -150$$

The position of Train A at time $t = 12$ minutes is approximately 150 meters west of Origin Station.

- (d) Let x be train A 's position, y train B 's position, and z the distance between train A and train B .

$$z^2 = x^2 + y^2 \Rightarrow 2z \frac{dz}{dt} = 2x \frac{dx}{dt} + 2y \frac{dy}{dt}$$

$$x = 300, y = 400 \Rightarrow z = 500$$

$$v_B(2) = -20 + 120 + 25 = 125$$

$$500 \frac{dz}{dt} = (300)(100) + (400)(125)$$

$$\frac{dz}{dt} = \frac{80000}{500} = 160 \text{ meters per minute}$$

1 : average acceleration

2 : $\begin{cases} 1 : v_A(8) < -100 < v_A(5) \\ 1 : \text{conclusion, using IVT} \end{cases}$

3 : $\begin{cases} 1 : \text{position expression} \\ 1 : \text{trapezoidal sum} \\ 1 : \text{position at time } t = 12 \end{cases}$

3 : $\begin{cases} 2 : \text{implicit differentiation of} \\ \quad \text{distance relationship} \\ 1 : \text{answer} \end{cases}$

2014 SCORING GUIDELINES

Question 5

x	-2	$-2 < x < -1$	-1	$-1 < x < 1$	1	$1 < x < 3$	3
$f(x)$	12	Positive	8	Positive	2	Positive	7
$f'(x)$	-5	Negative	0	Negative	0	Positive	$\frac{1}{2}$
$g(x)$	-1	Negative	0	Positive	3	Positive	1
$g'(x)$	2	Positive	$\frac{3}{2}$	Positive	0	Negative	-2

The twice-differentiable functions f and g are defined for all real numbers x . Values of f , f' , g , and g' for various values of x are given in the table above.

- (a) Find the x -coordinate of each relative minimum of f on the interval $[-2, 3]$. Justify your answers.
- (b) Explain why there must be a value c , for $-1 < c < 1$, such that $f''(c) = 0$.
- (c) The function h is defined by $h(x) = \ln(f(x))$. Find $h'(3)$. Show the computations that lead to your answer.
- (d) Evaluate $\int_{-2}^3 f'(g(x))g'(x) dx$.

- (a) $x = 1$ is the only critical point at which f' changes sign from negative to positive. Therefore, f has a relative minimum at $x = 1$.

- (b) f' is differentiable $\Rightarrow f'$ is continuous on the interval $-1 \leq x \leq 1$

$$\frac{f'(1) - f'(-1)}{1 - (-1)} = \frac{0 - 0}{2} = 0$$

Therefore, by the Mean Value Theorem, there is at least one value c , $-1 < c < 1$, such that $f''(c) = 0$.

- (c) $h'(x) = \frac{1}{f(x)} \cdot f'(x)$

$$h'(3) = \frac{1}{f(3)} \cdot f'(3) = \frac{1}{7} \cdot \frac{1}{2} = \frac{1}{14}$$

- (d) $\int_{-2}^3 f'(g(x))g'(x) dx = \left[f(g(x)) \right]_{x=-2}^{x=3}$
 $= f(g(3)) - f(g(-2))$
 $= f(1) - f(-1)$
 $= 2 - 8 = -6$

1 : answer with justification

$$2 : \begin{cases} 1 : f'(1) - f'(-1) = 0 \\ 1 : \text{explanation, using Mean Value Theorem} \end{cases}$$

$$3 : \begin{cases} 2 : h'(x) \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 2 : \text{Fundamental Theorem of Calculus} \\ 1 : \text{answer} \end{cases}$$


2014 SCORING GUIDELINES

Question 6

Consider the differential equation $\frac{dy}{dx} = (3 - y)\cos x$. Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(0) = 1$. The function f is defined for all real numbers.

- (a) A portion of the slope field of the differential equation is given below. Sketch the solution curve through the point $(0, 1)$.
- (b) Write an equation for the line tangent to the solution curve in part (a) at the point $(0, 1)$. Use the equation to approximate $f(0.2)$.
- (c) Find $y = f(x)$, the particular solution to the differential equation with the initial condition $f(0) = 1$.

(a) 1 : solution curve



(b) $\left. \frac{dy}{dx} \right|_{(x, y) = (0, 1)} = 2 \cos 0 = 2$

An equation for the tangent line is $y = 2x + 1$.

$$f(0.2) \approx 2(0.2) + 1 = 1.4$$

2 : $\begin{cases} 1 : \text{tangent line equation} \\ 1 : \text{approximation} \end{cases}$

(c) $\frac{dy}{dx} = (3 - y)\cos x$

$$\int \frac{dy}{3 - y} = \int \cos x \, dx$$

$$-\ln|3 - y| = \sin x + C$$

$$-\ln 2 = \sin 0 + C \Rightarrow C = -\ln 2$$

$$-\ln|3 - y| = \sin x - \ln 2$$

Because $y(0) = 1$, $y < 3$, so $|3 - y| = 3 - y$

$$3 - y = 2e^{-\sin x}$$

$$y = 3 - 2e^{-\sin x}$$

6 : $\begin{cases} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$

Note: max 3/6 [1-2-0-0-0] if no constant of integration

Note: 0/6 if no separation of variables