

Part B (AB or BC): Graphing calculator not allowed**Question 3****9 points****General Scoring Notes**


The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

The depth of seawater at a location can be modeled by the function H that satisfies the differential equation $\frac{dH}{dt} = \frac{1}{2}(H - 1)\cos\left(\frac{t}{2}\right)$, where $H(t)$ is measured in feet and t is measured in hours after noon ($t = 0$). It is known that $H(0) = 4$.

Model Solution**Scoring**

- (a) A portion of the slope field for the differential equation is provided. Sketch the solution curve, $y = H(t)$, through the point $(0, 4)$.



Solution curve

1 point**Scoring notes:**

- The solution curve must pass through the point $(0, 4)$, extend to at least $t = 4.5$, and have no obvious conflicts with the given slope lines.
- Only portions of the solution curve within the given slope field are considered.

Total for part (a) 1 point

- (b) For $0 < t < 5$, it can be shown that $H(t) > 1$. Find the value of t , for $0 < t < 5$, at which H has a critical point. Determine whether the critical point corresponds to a relative minimum, a relative maximum, or neither a relative minimum nor a relative maximum of the depth of seawater at the location. Justify your answer.

Because $H(t) > 1$, then $\frac{dH}{dt} = 0$ implies $\cos\left(\frac{t}{2}\right) = 0$.

Considers sign of $\frac{dH}{dt}$ **1 point**

This implies that $t = \pi$ is a critical point.

Identifies $t = \pi$ **1 point**

For $0 < t < \pi$, $\frac{dH}{dt} > 0$ and for $\pi < t < 5$, $\frac{dH}{dt} < 0$. Therefore, $t = \pi$ is the location of a relative maximum value of H .

Answer with justification **1 point**

Scoring notes:

- The first point is earned for considering $\frac{dH}{dt} = 0$, $\frac{dH}{dt} > 0$, $\frac{dH}{dt} < 0$, $\cos\left(\frac{t}{2}\right) = 0$, $\cos\left(\frac{t}{2}\right) > 0$, or $\cos\left(\frac{t}{2}\right) < 0$.
- The second point is earned for identifying $t = \pi$, with or without supporting work. A response may consider $H = 1$ or $t = 1$ as potential critical points without penalty.
- The third point cannot be earned without the first point. The third point is earned only for a correct justification and a correct answer of "relative maximum."
- The justification can be shown by determining the sign of $\frac{dH}{dt}$ (or $\cos\left(\frac{t}{2}\right)$) at a single value in $0 < t < \pi$ and at a single value in $\pi < t < 5$. It is not necessary to state that $\frac{dH}{dt}$ does not change sign on these intervals.
- The third point can also be earned by using the Second Derivative Test. For example:

$$\frac{d^2H}{dt^2} = \frac{1}{2}(H - 1)\left(-\frac{1}{2}\sin\left(\frac{t}{2}\right)\right) + \cos\left(\frac{t}{2}\right) \cdot \frac{1}{2} \cdot \frac{dH}{dt}$$

$$\frac{d^2H}{dt^2}\bigg|_{t=\pi} < 0$$
 Therefore, $t = \pi$ is the location of a relative maximum value of H .

Total for part (b) 3 points

- (c) Use separation of variables to find $y = H(t)$, the particular solution to the differential equation

$$\frac{dH}{dt} = \frac{1}{2}(H-1)\cos\left(\frac{t}{2}\right) \text{ with initial condition } H(0) = 4.$$

$\frac{dH}{H-1} = \frac{1}{2}\cos\left(\frac{t}{2}\right)dt$	Separation of variables	1 point
$\int \frac{dH}{H-1} = \int \frac{1}{2}\cos\left(\frac{t}{2}\right)dt$	One antiderivative	1 point
$\Rightarrow \ln H-1 = \sin\left(\frac{t}{2}\right) + C$	Second antiderivative	1 point
$\ln 4-1 = \sin\left(\frac{0}{2}\right) + C \Rightarrow C = \ln 3$ Because $H(0) = 4$, $H > 1$, so $ H-1 = H-1$. $\ln(H-1) = \sin\left(\frac{t}{2}\right) + \ln 3$	Constant of integration and uses initial condition	1 point
$H-1 = e^{\sin(t/2)+\ln 3} = 3e^{\sin(t/2)}$ $H(t) = 1 + 3e^{\sin(t/2)}$	Solves for H	1 point

Scoring notes:

- A response with no separation of variables earns 0 out of 5 points.
- A response that presents $\int \frac{dH}{H-1} = \ln(H-1)$ without absolute value symbols earns that antiderivative point.
- A response with no constant of integration can earn at most the first 3 points.
- A response is eligible for the fourth point only if it has earned the first point and at least 1 of the 2 antiderivative points.
- An eligible response earns the fourth point by correctly including the constant of integration in an equation and substituting 0 for t and 4 for H .
- A response is eligible for the fifth point only if it has earned the first 4 points.
- A response earns the fifth point only for an answer of $H(t) = 1 + 3e^{\sin(t/2)}$ or a mathematically equivalent expression for $H(t)$ such as $H(t) = 1 + e^{\sin(t/2)+\ln 3}$.
- A response does not need to argue that $|H-1| = H-1$ in order to earn the fifth point.
- Special case: A response that presents an incorrect separation of variables of $\frac{1}{2} \cdot \frac{dH}{H-1} = \cos\left(\frac{t}{2}\right)dt$ does not earn the first point or the fifth point but is eligible for the 2 antiderivative points. If the response earns at least 1 of the 2 antiderivative points, then the response is eligible for the fourth point.

Total for part (c) 5 points

Total for question 3 9 points

Part B (AB or BC): Graphing calculator not allowed

Question 3

9 points

General Scoring Notes


The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

A bottle of milk is taken out of a refrigerator and placed in a pan of hot water to be warmed. The increasing function M models the temperature of the milk at time t , where $M(t)$ is measured in degrees Celsius ($^{\circ}\text{C}$) and t is the number of minutes since the bottle was placed in the pan. M satisfies the differential equation $\frac{dM}{dt} = \frac{1}{4}(40 - M)$. At time $t = 0$, the temperature of the milk is 5°C . It can be shown that $M(t) < 40$ for all values of t .

Model Solution	Scoring
----------------	---------

- (a) A slope field for the differential equation $\frac{dM}{dt} = \frac{1}{4}(40 - M)$ is shown. Sketch the solution curve through the point $(0, 5)$.



Scoring notes:

- The solution curve must pass through the point $(0, 5)$, extend reasonably close to the left and right edges of the rectangle, and have no obvious conflicts with the given slope lines.
- Only portions of the solution curve within the given slope field are considered.
- The solution curve must lie entirely below the horizontal line segments at $M = 40$.

Total for part (a) 1 point

- (b) Use the line tangent to the graph of M at $t = 0$ to approximate $M(2)$, the temperature of the milk at time $t = 2$ minutes.

$\left. \frac{dM}{dt} \right _{t=0} = \frac{1}{4}(40 - 5) = \frac{35}{4}$	$\left. \frac{dM}{dt} \right _{t=0}$	1 point
---	--------------------------------------	----------------

The tangent line equation is $y = 5 + \frac{35}{4}(t - 0)$.

$$M(2) \approx 5 + \frac{35}{4} \cdot 2 = 22.5$$

The temperature of the milk at time $t = 2$ minutes is approximately 22.5° Celsius.

Scoring notes:

- The value of the slope may appear in a tangent line equation or approximation.
- A response of $5 + \frac{35}{4} \cdot 2$ is the minimal response to earn both points.
- A response of $\frac{1}{4}(40 - 5)$ earns the first point. If there are any subsequent errors in simplification, the response does not earn the second point.
- In order to earn the second point the response must present an approximation found by using a tangent line that:
 - passes through the point $(0, 5)$ and
 - has slope $\frac{35}{4}$ or a nonzero slope that is declared to be the value of $\frac{dM}{dt}$.
- An unsupported approximation does not earn the second point.
- The approximation need not be simplified, but the response does not earn the second point if the approximation is simplified incorrectly.

Total for part (b) 2 points

- (c) Write an expression for $\frac{d^2M}{dt^2}$ in terms of M . Use $\frac{d^2M}{dt^2}$ to determine whether the approximation from part (b) is an underestimate or an overestimate for the actual value of $M(2)$. Give a reason for your answer.

$\frac{d^2M}{dt^2} = -\frac{1}{4} \frac{dM}{dt} = -\frac{1}{4} \left(\frac{1}{4}(40 - M) \right) = -\frac{1}{16}(40 - M)$	$\frac{d^2M}{dt^2}$	1 point
--	---------------------	----------------

Because $M(t) < 40$, $\frac{d^2M}{dt^2} < 0$, so the graph of M is concave down. Therefore, the tangent line approximation of $M(2)$ is an overestimate.

Overestimate with reason **1 point**

Scoring notes:

- The first point is earned for either $\frac{d^2M}{dt^2} = -\frac{1}{4} \left(\frac{1}{4}(40 - M) \right)$ or $\frac{d^2M}{dt^2} = -\frac{1}{16}(40 - M)$ (or equivalent). A response that presents any subsequent simplification error does not earn the second point.
- A response that presents an expression for $\frac{d^2M}{dt^2}$ in terms of $\frac{dM}{dt}$ but fails to continue to an expression in terms of M (i.e., $\frac{d^2M}{dt^2} = -\frac{1}{4} \frac{dM}{dt}$) does not earn the first point but is eligible for the second point.
- If the response presents an expression for $\frac{d^2M}{dt^2}$ that is incorrect, the response is eligible for the second point only if the expression is a nonconstant linear function that is negative for $5 < M < 40$.
 - Special case: A response that presents $\frac{d^2M}{dt^2} = \frac{1}{16}(40 - M)$ does not earn the first point but is eligible to earn the second point for a consistent answer and reason.
- To earn the second point a response must include $\frac{d^2M}{dt^2} < 0$, or $\frac{dM}{dt}$ is decreasing, or the graph of M is concave down, as well as the conclusion that the approximation is an overestimate.
- A response that presents an argument based on $\frac{d^2M}{dt^2}$ or concavity at a single point does not earn the second point.

Total for part (c) 2 points

- (d) Use separation of variables to find an expression for $M(t)$, the particular solution to the differential equation $\frac{dM}{dt} = \frac{1}{4}(40 - M)$ with initial condition $M(0) = 5$.

$\frac{dM}{40 - M} = \frac{1}{4} dt$	Separates variables	1 point
--------------------------------------	---------------------	----------------

$$\int \frac{dM}{40 - M} = \int \frac{1}{4} dt$$

$-\ln 40 - M = \frac{1}{4}t + C$	Finds antiderivatives	1 point
-----------------------------------	-----------------------	----------------

$-\ln 40 - 5 = 0 + C \Rightarrow C = -\ln 35$ $M(t) < 40 \Rightarrow 40 - M > 0 \Rightarrow 40 - M = 40 - M$	Constant of integration and uses initial condition	1 point
--	--	----------------

$$-\ln(40 - M) = \frac{1}{4}t - \ln 35$$

$$\ln(40 - M) = -\frac{1}{4}t + \ln 35$$

$40 - M = 35e^{-t/4}$	Solves for M	1 point
$M = 40 - 35e^{-t/4}$		

Scoring notes:

- A response with no separation of variables earns 0 out of 4 points.
- A response that presents an antiderivative of $-\ln(40 - M)$ without absolute value symbols is eligible for all 4 points.
- A response with no constant of integration can earn at most the first 2 points.
- A response is eligible for the third point only if it has earned the first 2 points.
 - Special Case: A response that presents $+\ln(40 - M) = \frac{t}{4} + C$ (or equivalent) does not earn the second point, is eligible for the third point, but not eligible for the fourth.
- An eligible response earns the third point by correctly including the constant of integration in an equation and substituting 0 for t and 5 for M .
- A response is eligible for the fourth point only if it has earned the first 3 points.
- A response earns the fourth point only for an answer of $M = 40 - 35e^{-t/4}$ or equivalent.

Total for part (d) 4 points


Total for question 3 9 points

Part B (AB): Graphing calculator not allowed**Question 5****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Consider the differential equation $\frac{dy}{dx} = \frac{1}{2} \sin\left(\frac{\pi}{2}x\right)\sqrt{y+7}$. Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(1) = 2$. The function f is defined for all real numbers.

Model Solution	Scoring
(a) A portion of the slope field for the differential equation is given below. Sketch the solution curve through the point $(1, 2)$. 	Solution curve 1 point

Scoring notes:

- The solution curve must pass through the point $(1, 2)$, extend reasonably close to the left and right edges of the square and have no obvious conflicts with the given slope lines.
- Only portions of the solution curve within the given slope field are considered.
- The solution curve must indicate $f(x) > 0$ for all points on the curve.
- All local maximum/minimum points on the solution curve must occur at horizontal line segments in the slope field.

Total for part (a) 1 point

- (b) Write an equation for the line tangent to the solution curve in part (a) at the point (1, 2). Use the equation to approximate $f(0.8)$.

$$\left. \frac{dy}{dx} \right|_{(x,y)=(1,2)} = \frac{1}{2} \cdot 3 \cdot \sin\left(\frac{\pi}{2}\right) = \frac{3}{2}$$

An equation for the tangent line is $y = 2 + \frac{3}{2}(x - 1)$.

$$f(0.8) \approx 2 + \frac{3}{2}(0.8 - 1) = 1.7$$

Tangent line equation **1 point**

Approximation **1 point**

Scoring notes:

- The tangent line equation can be presented in any equivalent form.
- An incorrect tangent line equation with a slope of $\frac{3}{2}$ is eligible to earn the second point for a consistent answer.
- A response of only $2 + \frac{3}{2}(0.8 - 1)$ earns the second point but not the first.

Total for part (b) 2 points

- (c) It is known that $f''(x) > 0$ for $-1 \leq x \leq 1$. Is the approximation found in part (b) an overestimate or an underestimate for $f(0.8)$? Give a reason for your answer.

Because $f''(x) > 0$, f is concave up on $-1 \leq x \leq 1$, the tangent line lies below the graph of $y = f(x)$ at $x = 0.8$, and the approximation for $f(0.8)$ is an underestimate.

Answer with reason **1 point**

Scoring notes:

- The reason must include $f''(x) > 0$, $f'(x)$ is increasing, or $f(x)$ is concave up.

Total for part (c) 1 point

- (d) Use separation of variables to find $y = f(x)$, the particular solution to the differential equation

$$\frac{dy}{dx} = \frac{1}{2} \sin\left(\frac{\pi}{2}x\right) \sqrt{y+7} \text{ with the initial condition } f(1) = 2.$$

$$\int \frac{dy}{\sqrt{y+7}} = \int \frac{1}{2} \sin\left(\frac{\pi}{2}x\right) dx$$

Separation of variables **1 point**

$$2\sqrt{y+7} = -\frac{1}{\pi} \cos\left(\frac{\pi}{2}x\right) + C$$

One correct antiderivative **1 point**

The other correct antiderivative **1 point**

$$f(1) = 2 \Rightarrow 2\sqrt{2+7} = -\frac{1}{\pi} \cos\left(\frac{\pi}{2} \cdot 1\right) + C$$

$$\Rightarrow 6 = -\frac{1}{\pi} \cos\left(\frac{\pi}{2}\right) + C \Rightarrow C = 6$$

Constant of integration and uses initial condition **1 point**

$$\sqrt{y+7} = 3 - \frac{1}{2\pi} \cos\left(\frac{\pi}{2}x\right)$$

$$y = \left(3 - \frac{1}{2\pi} \cos\left(\frac{\pi}{2}x\right)\right)^2 - 7$$

Solves for y **1 point**

Scoring notes:

- A response with no separation of variables earns 0 out of 5 points.
- A response with no constant of integration can earn at most the first 3 points.
- A response is eligible for the fourth point only if it has earned the first point and at least 1 of the 2 antiderivative points.
 - Special case: The incorrect separation of $\sqrt{y+7} \, dy = \frac{1}{2} \sin\left(\frac{\pi}{2}x\right) dx$ does not earn the first point, is only eligible for the antiderivative point for $-\frac{1}{\pi} \cos\left(\frac{\pi}{2}x\right)$, and is eligible for the fourth point.
- An eligible response earns the fourth point by correctly including the constant of integration in an equation and substituting 1 for x and 2 for y .
- A response is eligible for the fifth point only if it has earned the first 4 points.

Total for part (d) 5 points

Total for question 5 9 points

Part B (AB): Graphing calculator not allowed**Question 6****9 points****General Scoring Notes**


Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Scoring guidelines and notes contain examples of the most common approaches seen in student responses. These guidelines can be applied to alternate approaches to ensure that these alternate approaches are scored appropriately.

A medication is administered to a patient. The amount, in milligrams, of the medication in the patient at time t hours is modeled by a function $y = A(t)$ that satisfies the differential equation $\frac{dy}{dt} = \frac{12 - y}{3}$. At time $t = 0$ hours, there are 0 milligrams of the medication in the patient.

Model Solution**Scoring**

- (a) A portion of the slope field for the differential equation $\frac{dy}{dt} = \frac{12 - y}{3}$ is given below. Sketch the solution curve through the point $(0, 0)$.

Solution curve **1 point****Scoring notes:**

- To earn the point the solution curve must pass through the point $(0, 0)$, be generally increasing and concave down, and approach the horizontal asymptote from below as t increases. The point is not earned if two or more solution curves are presented.

Total for part (a) 1 point

- (b) Using correct units, interpret the statement $\lim_{t \rightarrow \infty} A(t) = 12$ in the context of this problem.

Over time the amount of medication in the patient approaches 12 milligrams.	Interpretation	1 point
---	----------------	----------------

Scoring notes:

- To earn the point the interpretation must include “medication in the patient,” “approaches 12,” and units (milligrams), or their equivalents.

Total for part (b) 1 point

- (c) Use separation of variables to find $y = A(t)$, the particular solution to the differential equation

$$\frac{dy}{dt} = \frac{12 - y}{3} \text{ with initial condition } A(0) = 0.$$

$\frac{dy}{dt} = \frac{12 - y}{3} \Rightarrow \frac{dy}{12 - y} = \frac{dt}{3}$	Separation of variables	1 point
---	-------------------------	----------------

$\int \frac{dy}{12 - y} = \int \frac{dt}{3} \Rightarrow -\ln 12 - y = \frac{t}{3} + C$	Antiderivatives	1 point
---	-----------------	----------------

$\ln 12 - y = -\frac{t}{3} - C \Rightarrow 12 - y = e^{-t/3 - C}$ $\Rightarrow y = 12 + Ke^{-t/3}$	Constant of integration and uses initial condition	1 point
--	--	----------------

$$0 = 12 + K \Rightarrow K = -12$$

$y = A(t) = 12 - 12e^{-t/3}$	Solves for y	1 point
------------------------------	----------------	----------------

Scoring notes:

- A response of $\frac{dy}{12 - y} = 3 dt$ is a bad separation and does not earn the first point. However, this response is eligible for the second and third points. It cannot earn the fourth point.
- Absolute value bars are not required in this part.
- A response that correctly separates to $\frac{3 dy}{12 - y} = dt$ but then incorrectly simplifies to $\frac{dy}{4 - y} = dt$ earns the first point (for the initial correct separation), is eligible for the second point (for $-\ln|4 - y| = t$, with or without $+C$), but is not eligible for the third or fourth points.
- $+\ln|12 - y| = \frac{t}{3}$ (with or without $+C$) does not earn the second point and is not eligible for the fourth point; $+\ln|12 - y| = \frac{t}{3} + C$ is eligible for the third point.
- In all other cases, the points are earned consecutively—the second point cannot be earned without the first, the third without the second, etc.

Total for part (c) 4 points

- (d) A different procedure is used to administer the medication to a second patient. The amount, in milligrams, of the medication in the second patient at time t hours is modeled by a function $y = B(t)$ that satisfies the differential equation $\frac{dy}{dt} = 3 - \frac{y}{t+2}$. At time $t = 1$ hour, there are 2.5 milligrams of the medication in the second patient. Is the rate of change of the amount of medication in the second patient increasing or decreasing at time $t = 1$? Give a reason for your answer.

$\frac{dy}{dt} = 3 - \frac{y}{t+2} \Rightarrow \frac{d^2y}{dt^2} = (-1) \frac{\frac{dy}{dt}(t+2) - y}{(t+2)^2}$	Quotient rule	1 point
$B'(1) = 3 - \frac{B(1)}{3} = 3 - \frac{2.5}{3} = \frac{6.5}{3}$ $B''(1) = -\frac{B'(1) \cdot 3 - B(1)}{3^2} = -\frac{6.5 - 2.5}{9} = -\frac{4}{9} < 0$	$B''(1) < 0$	1 point
The rate of change of the amount of medication is decreasing at time $t = 1$ because $B''(1) < 0$ and $\frac{d^2y}{dt^2}$ is continuous in an interval containing $t = 1$.	Answer with reason	1 point

Scoring notes:

- The first point is for correctly applying the quotient rule to $\frac{y}{t+2}$ or applying the product rule to $y(t+2)^{-1}$. Errors in differentiating the constant, 3, or handling the sign of the second term of $\frac{dy}{dt}$ will result in not earning the second point.
- The second point cannot be earned unless the second derivative $\frac{d^2y}{dt^2}$ is correct.
- For the second point it is sufficient to state the sign of $B''(1)$ is negative with supporting work. If a value is declared for $B''(1)$, it must be correct in order to earn the second point.
- Eligibility for the third point: An attempt at using the quotient rule (or product rule) to find $B''(1)$. In this case the third point will be earned for a consistent conclusion based on the declared value (or sign) of $B''(1)$.

Total for part (d) 3 points**Total for question 6 9 points**

2018

AP Calculus AB

Scoring Guidelines

2018 SCORING GUIDELINES

Question 1

(a) $\int_0^{300} r(t) dt = 270$

According to the model, 270 people enter the line for the escalator during the time interval $0 \leq t \leq 300$.

(b) $20 + \int_0^{300} (r(t) - 0.7) dt = 20 + \int_0^{300} r(t) dt - 0.7 \cdot 300 = 80$

According to the model, 80 people are in line at time $t = 300$.

(c) Based on part (b), the number of people in line at time $t = 300$ is 80.

The first time t that there are no people in line is

$$300 + \frac{80}{0.7} = 414.286 \text{ (or } 414.285 \text{) seconds.}$$

(d) The total number of people in line at time t , $0 \leq t \leq 300$, is modeled by

$$20 + \int_0^t r(x) dx - 0.7t.$$

$$r(t) - 0.7 = 0 \Rightarrow t_1 = 33.013298, t_2 = 166.574719$$

t	People in line for escalator
0	20
t_1	3.803
t_2	158.070
300	80

The number of people in line is a minimum at time $t = 33.013$ seconds, when there are 4 people in line.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{considers rate out} \\ 1 : \text{answer} \end{cases}$

1 : answer

4 : $\begin{cases} 1 : \text{considers } r(t) - 0.7 = 0 \\ 1 : \text{identifies } t = 33.013 \\ 1 : \text{answers} \\ 1 : \text{justification} \end{cases}$

2018 SCORING GUIDELINES

Question 2

(a) $v'(3) = -2.118$

The acceleration of the particle at time $t = 3$ is -2.118 .

(b) $x(3) = x(0) + \int_0^3 v(t) dt = -5 + \int_0^3 v(t) dt = -1.760213$

The position of the particle at time $t = 3$ is -1.760 .

(c) $\int_0^{3.5} v(t) dt = 2.844$ (or 2.843)

$$\int_0^{3.5} |v(t)| dt = 3.737$$

The integral $\int_0^{3.5} v(t) dt$ is the displacement of the particle over the time interval $0 \leq t \leq 3.5$.

The integral $\int_0^{3.5} |v(t)| dt$ is the total distance traveled by the particle over the time interval $0 \leq t \leq 3.5$.

(d) $v(t) = x_2'(t)$

$$v(t) = 2t - 1 \Rightarrow t = 1.57054$$

The two particles are moving with the same velocity at time $t = 1.571$ (or 1.570).

1 : answer

3 : $\begin{cases} 1 : \int_0^3 v(t) dt \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

3 : $\begin{cases} 1 : \text{answers} \\ 2 : \text{interpretations of } \int_0^{3.5} v(t) dt \\ \text{and } \int_0^{3.5} |v(t)| dt \end{cases}$

2 : $\begin{cases} 1 : \text{sets } v(t) = x_2'(t) \\ 1 : \text{answer} \end{cases}$

2018 SCORING GUIDELINES

Question 3

$$\begin{aligned} \text{(a)} \quad f(-5) &= f(1) + \int_1^{-5} g(x) \, dx = f(1) - \int_{-5}^1 g(x) \, dx \\ &= 3 - \left(-9 - \frac{3}{2} + 1\right) = 3 - \left(-\frac{19}{2}\right) = \frac{25}{2} \end{aligned}$$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b)} \quad \int_1^6 g(x) \, dx &= \int_1^3 g(x) \, dx + \int_3^6 g(x) \, dx \\ &= \int_1^3 2 \, dx + \int_3^6 2(x-4)^2 \, dx \\ &= 4 + \left[\frac{2}{3}(x-4)^3\right]_{x=3}^{x=6} = 4 + \frac{16}{3} - \left(-\frac{2}{3}\right) = 10 \end{aligned}$$

3 : $\begin{cases} 1 : \text{split at } x = 3 \\ 1 : \text{antiderivative of } 2(x-4)^2 \\ 1 : \text{answer} \end{cases}$

(c) The graph of f is increasing and concave up on $0 < x < 1$ and $4 < x < 6$ because $f'(x) = g(x) > 0$ and $f'(x) = g(x)$ is increasing on those intervals.

2 : $\begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$

(d) The graph of f has a point of inflection at $x = 4$ because $f'(x) = g(x)$ changes from decreasing to increasing at $x = 4$.

2 : $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

2018 SCORING GUIDELINES

Question 4

$$\text{(a)} \quad H'(6) \approx \frac{H(7) - H(5)}{7 - 5} = \frac{11 - 6}{2} = \frac{5}{2}$$

$H'(6)$ is the rate at which the height of the tree is changing, in meters per year, at time $t = 6$ years.

2 : $\begin{cases} 1 : \text{estimate} \\ 1 : \text{interpretation with units} \end{cases}$

$$\text{(b)} \quad \frac{H(5) - H(3)}{5 - 3} = \frac{6 - 2}{2} = 2$$

Because H is differentiable on $3 \leq t \leq 5$, H is continuous on $3 \leq t \leq 5$.

By the Mean Value Theorem, there exists a value c , $3 < c < 5$, such that $H'(c) = 2$.

2 : $\begin{cases} 1 : \frac{H(5) - H(3)}{5 - 3} \\ 1 : \text{conclusion using} \\ \text{Mean Value Theorem} \end{cases}$

(c) The average height of the tree over the time interval $2 \leq t \leq 10$ is given by $\frac{1}{10 - 2} \int_2^{10} H(t) \, dt$.

2 : $\begin{cases} 1 : \text{trapezoidal sum} \\ 1 : \text{approximation} \end{cases}$

$$\begin{aligned} \frac{1}{8} \int_2^{10} H(t) \, dt &\approx \frac{1}{8} \left(\frac{1.5 + 2}{2} \cdot 1 + \frac{2 + 6}{2} \cdot 2 + \frac{6 + 11}{2} \cdot 2 + \frac{11 + 15}{2} \cdot 3 \right) \\ &= \frac{1}{8} (65.75) = \frac{263}{32} \end{aligned}$$

The average height of the tree over the time interval $2 \leq t \leq 10$ is $\frac{263}{32}$ meters.

$$\text{(d)} \quad G(x) = 50 \Rightarrow x = 1$$

3 : $\begin{cases} 2 : \frac{d}{dt}(G(x)) \\ 1 : \text{answer} \end{cases}$

$$\frac{d}{dt}(G(x)) = \frac{d}{dx}(G(x)) \cdot \frac{dx}{dt} = \frac{(1+x)100 - 100x \cdot 1}{(1+x)^2} \cdot \frac{dx}{dt} = \frac{100}{(1+x)^2} \cdot \frac{dx}{dt}$$

$$\left. \frac{d}{dt}(G(x)) \right|_{x=1} = \frac{100}{(1+1)^2} \cdot 0.03 = \frac{3}{4}$$

According to the model, the rate of change of the height of the tree with respect to time when the tree is 50 meters tall is $\frac{3}{4}$ meter per year.

Note: max 1/3 [1-0] if no chain rule

2018 SCORING GUIDELINES

Question 5

- (a) The average rate of change of
- f
- on the interval
- $0 \leq x \leq \pi$
- is

$$\frac{f(\pi) - f(0)}{\pi - 0} = \frac{-e^\pi - 1}{\pi}.$$

- (b)
- $f'(x) = e^x \cos x - e^x \sin x$

$$f'\left(\frac{3\pi}{2}\right) = e^{3\pi/2} \cos\left(\frac{3\pi}{2}\right) - e^{3\pi/2} \sin\left(\frac{3\pi}{2}\right) = e^{3\pi/2}$$

The slope of the line tangent to the graph of f at $x = \frac{3\pi}{2}$ is $e^{3\pi/2}$.

- (c)
- $f'(x) = 0 \Rightarrow \cos x - \sin x = 0 \Rightarrow x = \frac{\pi}{4}, x = \frac{5\pi}{4}$

x	$f(x)$
0	1
$\frac{\pi}{4}$	$\frac{1}{\sqrt{2}}e^{\pi/4}$
$\frac{5\pi}{4}$	$-\frac{1}{\sqrt{2}}e^{5\pi/4}$
2π	$e^{2\pi}$

The absolute minimum value of f on $0 \leq x \leq 2\pi$ is $-\frac{1}{\sqrt{2}}e^{5\pi/4}$.

- (d)
- $\lim_{x \rightarrow \pi/2} f(x) = 0$

Because g is differentiable, g is continuous.

$$\lim_{x \rightarrow \pi/2} g(x) = g\left(\frac{\pi}{2}\right) = 0$$

By L'Hospital's Rule,

$$\lim_{x \rightarrow \pi/2} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \pi/2} \frac{f'(x)}{g'(x)} = \frac{-e^{\pi/2}}{2}.$$

1 : answer


2 : $\begin{cases} 1 : f'(x) \\ 1 : \text{slope} \end{cases}$ 3 : $\begin{cases} 1 : \text{sets } f'(x) = 0 \\ 1 : \text{identifies } x = \frac{\pi}{4}, x = \frac{5\pi}{4} \\ \text{as candidates} \\ 1 : \text{answer with justification} \end{cases}$ 3 : $\begin{cases} 1 : g \text{ is continuous at } x = \frac{\pi}{2} \\ \text{and limits equal 0} \\ 1 : \text{applies L'Hospital's Rule} \\ 1 : \text{answer} \end{cases}$

Note: max 1/3 [1-0-0] if no limit notation attached to a ratio of derivatives

2018 SCORING GUIDELINES

Question 6

- (a)



- 2 :
- $\begin{cases} 1 : \text{solution curve through } (0, 2) \\ 1 : \text{solution curve through } (1, 0) \end{cases}$

Curves must go through the indicated points, follow the given slope lines, and extend to the boundary of the slope field.

- (b)
- $\frac{dy}{dx}\bigg|_{(x,y)=(1,0)} = \frac{4}{3}$

An equation for the line tangent to the graph of $y = f(x)$ at

$$x = 1 \text{ is } y = \frac{4}{3}(x - 1).$$

$$f(0.7) \approx \frac{4}{3}(0.7 - 1) = -0.4$$

- 2 :
- $\begin{cases} 1 : \text{equation of tangent line} \\ 1 : \text{approximation} \end{cases}$

- (c)
- $\frac{dy}{dx} = \frac{1}{3}x(y - 2)^2$

$$\int \frac{dy}{(y - 2)^2} = \int \frac{1}{3}x \, dx$$

$$\frac{-1}{y - 2} = \frac{1}{6}x^2 + C$$

$$\frac{1}{2} = \frac{1}{6} + C \Rightarrow C = \frac{1}{3}$$

$$\frac{-1}{y - 2} = \frac{1}{6}x^2 + \frac{1}{3} = \frac{x^2 + 2}{6}$$

$$y = 2 - \frac{6}{x^2 + 2}$$

Note: this solution is valid for $-\infty < x < \infty$.

- 5 :
- $\begin{cases} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ \text{and uses initial condition} \\ 1 : \text{solves for } y \end{cases}$

Note: 0/5 if no separation of variables

Note: max 3/5 [1-2-0-0] if no constant of integration

AP[®] Calculus AB

2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2016 SCORING GUIDELINES

Question 1

t (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.

- Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.
- Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
- Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
- For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

$$(a) \quad R'(2) \approx \frac{R(3) - R(1)}{3 - 1} = \frac{950 - 1190}{3 - 1} = -120 \text{ liters/hr}^2$$

$$2 : \begin{cases} 1 : \text{estimate} \\ 1 : \text{units} \end{cases}$$

$$(b) \quad \text{The total amount of water removed is given by } \int_0^8 R(t) \, dt.$$

$$\begin{aligned} \int_0^8 R(t) \, dt &\approx 1 \cdot R(0) + 2 \cdot R(1) + 3 \cdot R(3) + 2 \cdot R(6) \\ &= 1(1340) + 2(1190) + 3(950) + 2(740) \\ &= 8050 \text{ liters} \end{aligned}$$

This is an overestimate since R is a decreasing function.

$$\begin{aligned} (c) \quad \text{Total} &\approx 50000 + \int_0^8 W(t) \, dt - 8050 \\ &= 50000 + 7836.195325 - 8050 \approx 49786 \text{ liters} \end{aligned}$$

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{estimate} \end{cases}$$

$$(d) \quad W(0) - R(0) > 0, \quad W(8) - R(8) < 0, \quad \text{and } W(t) - R(t) \text{ is continuous.}$$

$$2 : \begin{cases} 1 : \text{considers } W(t) - R(t) \\ 1 : \text{answer with explanation} \end{cases}$$

Therefore, the Intermediate Value Theorem guarantees at least one time t , $0 < t < 8$, for which $W(t) - R(t) = 0$, or $W(t) = R(t)$.

For this value of t , the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank.

2016 SCORING GUIDELINES

Question 2

For $t \geq 0$, a particle moves along the x -axis. The velocity of the particle at time t is given by

$v(t) = 1 + 2\sin\left(\frac{t^2}{2}\right)$. The particle is at position $x = 2$ at time $t = 4$.

- (a) At time $t = 4$, is the particle speeding up or slowing down?
 (b) Find all times t in the interval $0 < t < 3$ when the particle changes direction. Justify your answer.
 (c) Find the position of the particle at time $t = 0$.
 (d) Find the total distance the particle travels from time $t = 0$ to time $t = 3$.

(a) $v(4) = 2.978716 > 0$
 $v'(4) = -1.164000 < 0$

The particle is slowing down since the velocity and acceleration have different signs.

(b) $v(t) = 0 \Rightarrow t = 2.707468$

$v(t)$ changes from positive to negative at $t = 2.707$.
 Therefore, the particle changes direction at this time.

(c) $x(0) = x(4) + \int_4^0 v(t) dt$
 $= 2 + (-5.815027) = -3.815$

(d) Distance $= \int_0^3 |v(t)| dt = 5.301$

2 : conclusion with reason

2 : $\begin{cases} 1 : t = 2.707 \\ 1 : \text{justification} \end{cases}$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$


2016 SCORING GUIDELINES

Question 3

The figure above shows the graph of the piecewise-linear function f . For $-4 \leq x \leq 12$, the function g is defined by

$$g(x) = \int_2^x f(t) dt.$$

- (a) Does g have a relative minimum, a relative maximum, or neither at $x = 10$? Justify your answer.
 (b) Does the graph of g have a point of inflection at $x = 4$? Justify your answer.
 (c) Find the absolute minimum value and the absolute maximum value of g on the interval $-4 \leq x \leq 12$. Justify your answers.



Graph of f

- (d) For $-4 \leq x \leq 12$, find all intervals for which $g(x) \leq 0$.

- (a) The function g has neither a relative minimum nor a relative maximum at $x = 10$ since $g'(x) = f(x)$ and $f(x) \leq 0$ for $8 \leq x \leq 12$.

- (b) The graph of g has a point of inflection at $x = 4$ since $g'(x) = f(x)$ is increasing for $2 \leq x \leq 4$ and decreasing for $4 \leq x \leq 6$.

- (c) $g'(x) = f(x)$ changes sign only at $x = -2$ and $x = 6$.

x	$g(x)$
-4	-4
-2	-8
6	8
12	-4

On the interval $-4 \leq x \leq 12$, the absolute minimum value is $g(-2) = -8$ and the absolute maximum value is $g(6) = 8$.

- (d) $g(x) \leq 0$ for $-4 \leq x \leq 2$ and $10 \leq x \leq 12$.

1 : $g'(x) = f(x)$ in (a), (b), (c), or (d)

1 : answer with justification

1 : answer with justification

4 : $\begin{cases} 1 : \text{considers } x = -2 \text{ and } x = 6 \\ \quad \text{as candidates} \\ 1 : \text{considers } x = -4 \text{ and } x = 12 \\ 2 : \text{answers with justification} \end{cases}$


2 : intervals

2016 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = \frac{y^2}{x-1}$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.
- (b) Let $y = f(x)$ be the particular solution to the given differential equation with the initial condition $f(2) = 3$. Write an equation for the line tangent to the graph of $y = f(x)$ at $x = 2$. Use your equation to approximate $f(2.1)$.
- (c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(2) = 3$.



(b) $\left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} = \frac{3^2}{2-1} = 9$

An equation for the tangent line is $y = 9(x - 2) + 3$.

$$f(2.1) \approx 9(2.1 - 2) + 3 = 3.9$$

(c) $\frac{1}{y^2} dy = \frac{1}{x-1} dx$

$$\int \frac{1}{y^2} dy = \int \frac{1}{x-1} dx$$

$$-\frac{1}{y} = \ln|x-1| + C$$

$$-\frac{1}{3} = \ln|2-1| + C \Rightarrow C = -\frac{1}{3}$$

$$-\frac{1}{y} = \ln|x-1| - \frac{1}{3}$$

$$y = \frac{1}{\frac{1}{3} - \ln(x-1)}$$

Note: This solution is valid for $1 < x < 1 + e^{1/3}$.

2 : $\begin{cases} 1 : \text{zero slopes} \\ 1 : \text{nonzero slopes} \end{cases}$

2 : $\begin{cases} 1 : \text{tangent line equation} \\ 1 : \text{approximation} \end{cases}$


5 : $\begin{cases} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration and uses initial condition} \\ 1 : \text{solves for } y \end{cases}$

Note: max 3/5 [1-2-0-0] if no constant of integration

Note: 0/5 if no separation of variables

2016 SCORING GUIDELINES

Question 5



The inside of a funnel of height 10 inches has circular cross sections, as shown in the figure above. At height h , the radius of the funnel is given by $r = \frac{1}{20}(3 + h^2)$, where $0 \leq h \leq 10$. The units of r and h are inches.

- (a) Find the average value of the radius of the funnel.
- (b) Find the volume of the funnel.
- (c) The funnel contains liquid that is draining from the bottom. At the instant when the height of the liquid is $h = 3$ inches, the radius of the surface of the liquid is decreasing at a rate of $\frac{1}{5}$ inch per second. At this instant, what is the rate of change of the height of the liquid with respect to time?

(a) Average radius $= \frac{1}{10} \int_0^{10} \frac{1}{20}(3 + h^2) dh = \frac{1}{200} \left[3h + \frac{h^3}{3} \right]_0^{10}$

$$= \frac{1}{200} \left(\left(30 + \frac{1000}{3} \right) - 0 \right) = \frac{109}{60} \text{ in}$$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

(b) Volume $= \pi \int_0^{10} \left(\left(\frac{1}{20} \right) (3 + h^2) \right)^2 dh = \frac{\pi}{400} \int_0^{10} (9 + 6h^2 + h^4) dh$

$$= \frac{\pi}{400} \left[9h + 2h^3 + \frac{h^5}{5} \right]_0^{10}$$

$$= \frac{\pi}{400} \left(\left(90 + 2000 + \frac{100000}{5} \right) - 0 \right) = \frac{2209\pi}{40} \text{ in}^3$$

3 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

(c) $\frac{dr}{dt} = \frac{1}{20}(2h) \frac{dh}{dt}$

$$-\frac{1}{5} = \frac{3}{10} \frac{dh}{dt}$$

$$\frac{dh}{dt} = -\frac{1}{5} \cdot \frac{10}{3} = -\frac{2}{3} \text{ in/sec}$$

3 : $\begin{cases} 2 : \text{chain rule} \\ 1 : \text{answer} \end{cases}$

2016 SCORING GUIDELINES

Question 6

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	-6	3	2	8
2	2	-2	-3	0
3	8	7	6	2
6	4	5	3	-1

The functions f and g have continuous second derivatives. The table above gives values of the functions and their derivatives at selected values of x .

(a) Let $k(x) = f(g(x))$. Write an equation for the line tangent to the graph of k at $x = 3$.

(b) Let $h(x) = \frac{g(x)}{f(x)}$. Find $h'(1)$.

(c) Evaluate $\int_1^3 f''(2x) dx$.

$$\begin{aligned} \text{(a)} \quad k(3) &= f(g(3)) = f(6) = 4 \\ k'(3) &= f'(g(3)) \cdot g'(3) = f'(6) \cdot 2 = 5 \cdot 2 = 10 \end{aligned}$$

An equation for the tangent line is $y = 10(x - 3) + 4$.

$$\begin{aligned} \text{(b)} \quad h'(1) &= \frac{f(1) \cdot g'(1) - g(1) \cdot f'(1)}{(f(1))^2} \\ &= \frac{(-6) \cdot 8 - 2 \cdot 3}{(-6)^2} = \frac{-54}{36} = -\frac{3}{2} \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad \int_1^3 f''(2x) dx &= \frac{1}{2} [f'(2x)]_1^3 = \frac{1}{2} [f'(6) - f'(2)] \\ &= \frac{1}{2} [5 - (-2)] = \frac{7}{2} \end{aligned}$$

$$3 : \begin{cases} 2 : \text{slope at } x = 3 \\ 1 : \text{equation for tangent line} \end{cases}$$

$$3 : \begin{cases} 2 : \text{expression for } h'(1) \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 2 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$$

AP[®] Calculus AB

2015 Scoring Guidelines

© 2015 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2015 SCORING GUIDELINES

Question 1

The rate at which rainwater flows into a drainpipe is modeled by the function R , where $R(t) = 20\sin\left(\frac{t^2}{35}\right)$ cubic feet per hour, t is measured in hours, and $0 \leq t \leq 8$. The pipe is partially blocked, allowing water to drain out the other end of the pipe at a rate modeled by $D(t) = -0.04t^3 + 0.4t^2 + 0.96t$ cubic feet per hour, for $0 \leq t \leq 8$. There are 30 cubic feet of water in the pipe at time $t = 0$.

- (a) How many cubic feet of rainwater flow into the pipe during the 8-hour time interval $0 \leq t \leq 8$?
- (b) Is the amount of water in the pipe increasing or decreasing at time $t = 3$ hours? Give a reason for your answer.
- (c) At what time t , $0 \leq t \leq 8$, is the amount of water in the pipe at a minimum? Justify your answer.
- (d) The pipe can hold 50 cubic feet of water before overflowing. For $t > 8$, water continues to flow into and out of the pipe at the given rates until the pipe begins to overflow. Write, but do not solve, an equation involving one or more integrals that gives the time w when the pipe will begin to overflow.

(a) $\int_0^8 R(t) dt = 76.570$

2 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b) $R(3) - D(3) = -0.313632 < 0$
 Since $R(3) < D(3)$, the amount of water in the pipe is decreasing at time $t = 3$ hours.

2 : $\begin{cases} 1 : \text{considers } R(3) \text{ and } D(3) \\ 1 : \text{answer and reason} \end{cases}$

(c) The amount of water in the pipe at time t , $0 \leq t \leq 8$, is $30 + \int_0^t [R(x) - D(x)] dx$.

3 : $\begin{cases} 1 : \text{considers } R(t) - D(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

$$R(t) - D(t) = 0 \Rightarrow t = 0, 3.271658$$

t	Amount of water in the pipe
0	30
3.271658	27.964561
8	48.543686

The amount of water in the pipe is a minimum at time $t = 3.272$ (or 3.271) hours.


(d) $30 + \int_0^w [R(t) - D(t)] dt = 50$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{equation} \end{cases}$

2015 SCORING GUIDELINES

Question 2

Let f and g be the functions defined by $f(x) = 1 + x + e^{x^2-2x}$ and $g(x) = x^4 - 6.5x^2 + 6x + 2$. Let R and S be the two regions enclosed by the graphs of f and g shown in the figure above.



- (a) Find the sum of the areas of regions R and S .
- (b) Region S is the base of a solid whose cross sections perpendicular to the x -axis are squares. Find the volume of the solid.
- (c) Let h be the vertical distance between the graphs of f and g in region S . Find the rate at which h changes with respect to x when $x = 1.8$.

- (a) The graphs of $y = f(x)$ and $y = g(x)$ intersect in the first quadrant at the points $(0, 2)$, $(2, 4)$, and $(A, B) = (1.032832, 2.401108)$.

$$\begin{aligned} \text{Area} &= \int_0^A [g(x) - f(x)] dx + \int_A^2 [f(x) - g(x)] dx \\ &= 0.997427 + 1.006919 = 2.004 \end{aligned}$$

(b) $\text{Volume} = \int_A^2 [f(x) - g(x)]^2 dx = 1.283$

(c) $h(x) = f(x) - g(x)$
 $h'(x) = f'(x) - g'(x)$
 $h'(1.8) = f'(1.8) - g'(1.8) = -3.812$ (or -3.811)

4 : $\begin{cases} 1 : \text{limits} \\ 2 : \text{integrands} \\ 1 : \text{answer} \end{cases}$

3 : $\begin{cases} 2 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{considers } h' \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 3

t (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	-220	150

Johanna jogs along a straight path. For $0 \leq t \leq 40$, Johanna's velocity is given by a differentiable function v . Selected values of $v(t)$, where t is measured in minutes and $v(t)$ is measured in meters per minute, are given in the table above.

- (a) Use the data in the table to estimate the value of $v'(16)$.
- (b) Using correct units, explain the meaning of the definite integral $\int_0^{40} |v(t)| dt$ in the context of the problem.

Approximate the value of $\int_0^{40} |v(t)| dt$ using a right Riemann sum with the four subintervals indicated in the table.

- (c) Bob is riding his bicycle along the same path. For $0 \leq t \leq 10$, Bob's velocity is modeled by $B(t) = t^3 - 6t^2 + 300$, where t is measured in minutes and $B(t)$ is measured in meters per minute.

Find Bob's acceleration at time $t = 5$.

- (d) Based on the model B from part (c), find Bob's average velocity during the interval $0 \leq t \leq 10$.

(a) $v'(16) \approx \frac{240 - 200}{20 - 12} = 5$ meters/min²

1 : approximation

- (b) $\int_0^{40} |v(t)| dt$ is the total distance Johanna jogs, in meters, over the time interval $0 \leq t \leq 40$ minutes.

3 : $\begin{cases} 1 : \text{explanation} \\ 1 : \text{right Riemann sum} \\ 1 : \text{approximation} \end{cases}$

$$\begin{aligned} \int_0^{40} |v(t)| dt &\approx 12 \cdot |v(12)| + 8 \cdot |v(20)| + 4 \cdot |v(24)| + 16 \cdot |v(40)| \\ &= 12 \cdot 200 + 8 \cdot 240 + 4 \cdot 220 + 16 \cdot 150 \\ &= 2400 + 1920 + 880 + 2400 \\ &= 7600 \text{ meters} \end{aligned}$$

- (c) Bob's acceleration is $B'(t) = 3t^2 - 12t$.

$$B'(5) = 3(25) - 12(5) = 15 \text{ meters/min}^2$$

2 : $\begin{cases} 1 : \text{uses } B'(t) \\ 1 : \text{answer} \end{cases}$

- (d) Avg vel = $\frac{1}{10} \int_0^{10} (t^3 - 6t^2 + 300) dt$

$$\begin{aligned} &= \frac{1}{10} \left[\frac{t^4}{4} - 2t^3 + 300t \right]_0^{10} \\ &= \frac{1}{10} \left[\frac{10000}{4} - 2000 + 3000 \right] = 350 \text{ meters/min} \end{aligned}$$


3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = 2x - y$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.
- (b) Find $\frac{d^2y}{dx^2}$ in terms of x and y . Determine the concavity of all solution curves for the given differential equation in Quadrant II. Give a reason for your answer.
- (c) Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(2) = 3$. Does f have a relative minimum, a relative maximum, or neither at $x = 2$? Justify your answer.
- (d) Find the values of the constants m and b for which $y = mx + b$ is a solution to the differential equation.



2 : $\begin{cases} 1 : \text{slopes where } x = 0 \\ 1 : \text{slopes where } x = 1 \end{cases}$

(b) $\frac{d^2y}{dx^2} = 2 - \frac{dy}{dx} = 2 - (2x - y) = 2 - 2x + y$

In Quadrant II, $x < 0$ and $y > 0$, so $2 - 2x + y > 0$.

Therefore, all solution curves are concave up in Quadrant II.

2 : $\begin{cases} 1 : \frac{d^2y}{dx^2} \\ 1 : \text{concave up with reason} \end{cases}$

(c) $\left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} = 2(2) - 3 = 1 \neq 0$

Therefore, f has neither a relative minimum nor a relative maximum at $x = 2$.

2 : $\begin{cases} 1 : \text{considers } \left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} \\ 1 : \text{conclusion with justification} \end{cases}$

(d) $y = mx + b \Rightarrow \frac{dy}{dx} = \frac{d}{dx}(mx + b) = m$

$$2x - y = m$$

$$2x - (mx + b) = m$$

$$(2 - m)x - (m + b) = 0$$

$$2 - m = 0 \Rightarrow m = 2$$


$$b = -m \Rightarrow b = -2$$

3 : $\begin{cases} 1 : \frac{d}{dx}(mx + b) = m \\ 1 : 2x - y = m \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 5

The figure above shows the graph of f' , the derivative of a twice-differentiable function f , on the interval $[-3, 4]$. The graph of f' has horizontal tangents at $x = -1$, $x = 1$, and $x = 3$. The areas of the regions bounded by the x -axis and the graph of f' on the intervals $[-2, 1]$ and $[1, 4]$ are 9 and 12, respectively.



- (a) Find all x -coordinates at which f has a relative maximum. Give a reason for your answer.
- (b) On what open intervals contained in $-3 < x < 4$ is the graph of f both concave down and decreasing? Give a reason for your answer.
- (c) Find the x -coordinates of all points of inflection for the graph of f . Give a reason for your answer.
- (d) Given that $f(1) = 3$, write an expression for $f(x)$ that involves an integral. Find $f(4)$ and $f(-2)$.

- (a) $f'(x) = 0$ at $x = -2$, $x = 1$, and $x = 4$.
 $f'(x)$ changes from positive to negative at $x = -2$.
 Therefore, f has a relative maximum at $x = -2$.

2 : $\begin{cases} 1 : \text{identifies } x = -2 \\ 1 : \text{answer with reason} \end{cases}$

- (b) The graph of f is concave down and decreasing on the intervals $-2 < x < -1$ and $1 < x < 3$ because f' is decreasing and negative on these intervals.

2 : $\begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$

- (c) The graph of f has a point of inflection at $x = -1$ and $x = 3$ because f' changes from decreasing to increasing at these points.

2 : $\begin{cases} 1 : \text{identifies } x = -1, 1, \text{ and } 3 \\ 1 : \text{reason} \end{cases}$

The graph of f has a point of inflection at $x = 1$ because f' changes from increasing to decreasing at this point.

- (d) $f(x) = 3 + \int_1^x f'(t) dt$

$$f(4) = 3 + \int_1^4 f'(t) dt = 3 + (-12) = -9$$

$$f(-2) = 3 + \int_1^{-2} f'(t) dt = 3 - \int_{-2}^1 f'(t) dt = 3 - (-9) = 12$$

3 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{expression for } f(x) \\ 1 : f(4) \text{ and } f(-2) \end{cases}$

2015 SCORING GUIDELINES

Question 6

Consider the curve given by the equation $y^3 - xy = 2$. It can be shown that $\frac{dy}{dx} = \frac{y}{3y^2 - x}$.

- (a) Write an equation for the line tangent to the curve at the point $(-1, 1)$.
- (b) Find the coordinates of all points on the curve at which the line tangent to the curve at that point is vertical.
- (c) Evaluate $\frac{d^2y}{dx^2}$ at the point on the curve where $x = -1$ and $y = 1$.

(a) $\frac{dy}{dx}\bigg|_{(x,y)=(-1,1)} = \frac{1}{3(1)^2 - (-1)} = \frac{1}{4}$

2 : $\begin{cases} 1 : \text{slope} \\ 1 : \text{equation for tangent line} \end{cases}$

An equation for the tangent line is $y = \frac{1}{4}(x + 1) + 1$.

(b) $3y^2 - x = 0 \Rightarrow x = 3y^2$

So, $y^3 - xy = 2 \Rightarrow y^3 - (3y^2)(y) = 2 \Rightarrow y = -1$

$(-1)^3 - x(-1) = 2 \Rightarrow x = 3$

The tangent line to the curve is vertical at the point $(3, -1)$.

3 : $\begin{cases} 1 : \text{sets } 3y^2 - x = 0 \\ 1 : \text{equation in one variable} \\ 1 : \text{coordinates} \end{cases}$

(c) $\frac{d^2y}{dx^2} = \frac{(3y^2 - x)\frac{dy}{dx} - y\left(6y\frac{dy}{dx} - 1\right)}{(3y^2 - x)^2}$

$$\frac{d^2y}{dx^2}\bigg|_{(x,y)=(-1,1)} = \frac{(3 \cdot 1^2 - (-1)) \cdot \frac{1}{4} - 1 \cdot \left(6 \cdot 1 \cdot \frac{1}{4} - 1\right)}{(3 \cdot 1^2 - (-1))^2} = \frac{1 - \frac{1}{2}}{16} = \frac{1}{32}$$

4 : $\begin{cases} 2 : \text{implicit differentiation} \\ 1 : \text{substitution for } \frac{dy}{dx} \\ 1 : \text{answer} \end{cases}$

AP[®] Calculus AB

2014 Scoring Guidelines

© 2014 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.
Visit the College Board on the Web: www.collegeboard.org.
AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2014 SCORING GUIDELINES

Question 1

Grass clippings are placed in a bin, where they decompose. For $0 \leq t \leq 30$, the amount of grass clippings remaining in the bin is modeled by $A(t) = 6.687(0.931)^t$, where $A(t)$ is measured in pounds and t is measured in days.

- Find the average rate of change of $A(t)$ over the interval $0 \leq t \leq 30$. Indicate units of measure.
- Find the value of $A'(15)$. Using correct units, interpret the meaning of the value in the context of the problem.
- Find the time t for which the amount of grass clippings in the bin is equal to the average amount of grass clippings in the bin over the interval $0 \leq t \leq 30$.
- For $t > 30$, $L(t)$, the linear approximation to A at $t = 30$, is a better model for the amount of grass clippings remaining in the bin. Use $L(t)$ to predict the time at which there will be 0.5 pound of grass clippings remaining in the bin. Show the work that leads to your answer.

$$(a) \quad \frac{A(30) - A(0)}{30 - 0} = -0.197 \text{ (or } -0.196) \text{ lbs/day}$$

1 : answer with units

$$(b) \quad A'(15) = -0.164 \text{ (or } -0.163)$$

The amount of grass clippings in the bin is decreasing at a rate of 0.164 (or 0.163) lbs/day at time $t = 15$ days.

2 : $\begin{cases} 1 : A'(15) \\ 1 : \text{interpretation} \end{cases}$

$$(c) \quad A(t) = \frac{1}{30} \int_0^{30} A(t) \, dt \Rightarrow t = 12.415 \text{ (or } 12.414)$$

2 : $\begin{cases} 1 : \frac{1}{30} \int_0^{30} A(t) \, dt \\ 1 : \text{answer} \end{cases}$

$$(d) \quad L(t) = A(30) + A'(30) \cdot (t - 30)$$

$$A'(30) = -0.055976$$

$$A(30) = 0.782928$$

$$L(t) = 0.5 \Rightarrow t = 35.054$$


4 : $\begin{cases} 2 : \text{expression for } L(t) \\ 1 : L(t) = 0.5 \\ 1 : \text{answer} \end{cases}$

2014 SCORING GUIDELINES

Question 2

Let R be the region enclosed by the graph of $f(x) = x^4 - 2.3x^3 + 4$ and the horizontal line $y = 4$, as shown in the figure above.

- (a) Find the volume of the solid generated when R is rotated about the horizontal line $y = -2$.
- (b) Region R is the base of a solid. For this solid, each cross section perpendicular to the x -axis is an isosceles right triangle with a leg in R . Find the volume of the solid.
- (c) The vertical line $x = k$ divides R into two regions with equal areas. Write, but do not solve, an equation involving integral expressions whose solution gives the value k .



(a) $f(x) = 4 \Rightarrow x = 0, 2.3$

$$\text{Volume} = \pi \int_0^{2.3} [(4+2)^2 - (f(x)+2)^2] dx$$

$$= 98.868 \text{ (or } 98.867)$$

4 : $\begin{cases} 2 : \text{integrand} \\ 1 : \text{limits} \\ 1 : \text{answer} \end{cases}$

(b) $\text{Volume} = \int_0^{2.3} \frac{1}{2} (4 - f(x))^2 dx$

$$= 3.574 \text{ (or } 3.573)$$

3 : $\begin{cases} 2 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(c) $\int_0^k (4 - f(x)) dx = \int_k^{2.3} (4 - f(x)) dx$

2 : $\begin{cases} 1 : \text{area of one region} \\ 1 : \text{equation} \end{cases}$


2014 SCORING GUIDELINES

Question 3

The function f is defined on the closed interval $[-5, 4]$. The graph of f consists of three line segments and is shown in the figure above.

Let g be the function defined by $g(x) = \int_{-3}^x f(t) dt$.

- (a) Find $g(3)$.
- (b) On what open intervals contained in $-5 < x < 4$ is the graph of g both increasing and concave down? Give a reason for your answer.
- (c) The function h is defined by $h(x) = \frac{g(x)}{5x}$. Find $h'(3)$.
- (d) The function p is defined by $p(x) = f(x^2 - x)$. Find the slope of the line tangent to the graph of p at the point where $x = -1$.

Graph of f

(a) $g(3) = \int_{-3}^3 f(t) dt = 6 + 4 - 1 = 9$

1 : answer

(b) $g'(x) = f(x)$

The graph of g is increasing and concave down on the intervals $-5 < x < -3$ and $0 < x < 2$ because $g' = f$ is positive and decreasing on these intervals.

2 : $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

(c) $h'(x) = \frac{5xg'(x) - g(x)5}{(5x)^2} = \frac{5xg'(x) - 5g(x)}{25x^2}$

3 : $\begin{cases} 2 : h'(x) \\ 1 : \text{answer} \end{cases}$

$$h'(3) = \frac{(5)(3)g'(3) - 5g(3)}{25 \cdot 3^2}$$

$$= \frac{15(-2) - 5(9)}{225} = \frac{-75}{225} = -\frac{1}{3}$$

(d) $p'(x) = f'(x^2 - x)(2x - 1)$

$$p'(-1) = f'(-2)(-3) = (-2)(-3) = 6$$

3 : $\begin{cases} 2 : p'(x) \\ 1 : \text{answer} \end{cases}$

2014 SCORING GUIDELINES

Question 4

Train A runs back and forth on an east-west section of railroad track. Train A 's velocity, measured in meters per minute, is given by a differentiable function $v_A(t)$, where time t is measured in minutes. Selected values for $v_A(t)$ are given in the table above.

t (minutes)	0	2	5	8	12
$v_A(t)$ (meters/minute)	0	100	40	-120	-150

- (a) Find the average acceleration of train A over the interval $2 \leq t \leq 8$.
- (b) Do the data in the table support the conclusion that train A 's velocity is -100 meters per minute at some time t with $5 < t < 8$? Give a reason for your answer.
- (c) At time $t = 2$, train A 's position is 300 meters east of the Origin Station, and the train is moving to the east. Write an expression involving an integral that gives the position of train A , in meters from the Origin Station, at time $t = 12$. Use a trapezoidal sum with three subintervals indicated by the table to approximate the position of the train at time $t = 12$.
- (d) A second train, train B , travels north from the Origin Station. At time t the velocity of train B is given by $v_B(t) = -5t^2 + 60t + 25$, and at time $t = 2$ the train is 400 meters north of the station. Find the rate, in meters per minute, at which the distance between train A and train B is changing at time $t = 2$.

(a) average accel = $\frac{v_A(8) - v_A(2)}{8 - 2} = \frac{-120 - 100}{6} = -\frac{110}{3}$ m/min²

1 : average acceleration

(b) v_A is differentiable $\Rightarrow v_A$ is continuous
 $v_A(8) = -120 < -100 < 40 = v_A(5)$

2 : $\begin{cases} 1 : v_A(8) < -100 < v_A(5) \\ 1 : \text{conclusion, using IVT} \end{cases}$

Therefore, by the Intermediate Value Theorem, there is a time t , $5 < t < 8$, such that $v_A(t) = -100$.

(c) $s_A(12) = s_A(2) + \int_2^{12} v_A(t) dt = 300 + \int_2^{12} v_A(t) dt$
 $\int_2^{12} v_A(t) dt \approx 3 \cdot \frac{100 + 40}{2} + 3 \cdot \frac{40 - 120}{2} + 4 \cdot \frac{-120 - 150}{2}$
 $= -450$

3 : $\begin{cases} 1 : \text{position expression} \\ 1 : \text{trapezoidal sum} \\ 1 : \text{position at time } t = 12 \end{cases}$

$s_A(12) \approx 300 - 450 = -150$

The position of Train A at time $t = 12$ minutes is approximately 150 meters west of Origin Station.

- (d) Let x be train A 's position, y train B 's position, and z the distance between train A and train B .

$z^2 = x^2 + y^2 \Rightarrow 2z \frac{dz}{dt} = 2x \frac{dx}{dt} + 2y \frac{dy}{dt}$

$x = 300, y = 400 \Rightarrow z = 500$

$v_B(2) = -20 + 120 + 25 = 125$

$500 \frac{dz}{dt} = (300)(100) + (400)(125)$

$\frac{dz}{dt} = \frac{80000}{500} = 160$ meters per minute

3 : $\begin{cases} 2 : \text{implicit differentiation of} \\ \text{distance relationship} \\ 1 : \text{answer} \end{cases}$

2014 SCORING GUIDELINES

Question 5

x	-2	$-2 < x < -1$	-1	$-1 < x < 1$	1	$1 < x < 3$	3
$f(x)$	12	Positive	8	Positive	2	Positive	7
$f'(x)$	-5	Negative	0	Negative	0	Positive	$\frac{1}{2}$
$g(x)$	-1	Negative	0	Positive	3	Positive	1
$g'(x)$	2	Positive	$\frac{3}{2}$	Positive	0	Negative	-2

The twice-differentiable functions f and g are defined for all real numbers x . Values of f , f' , g , and g' for various values of x are given in the table above.

- (a) Find the x -coordinate of each relative minimum of f on the interval $[-2, 3]$. Justify your answers.
- (b) Explain why there must be a value c , for $-1 < c < 1$, such that $f''(c) = 0$.
- (c) The function h is defined by $h(x) = \ln(f(x))$. Find $h'(3)$. Show the computations that lead to your answer.
- (d) Evaluate $\int_{-2}^3 f'(g(x))g'(x) dx$.

- (a) $x = 1$ is the only critical point at which f' changes sign from negative to positive. Therefore, f has a relative minimum at $x = 1$.

1 : answer with justification

- (b) f' is differentiable $\Rightarrow f'$ is continuous on the interval $-1 \leq x \leq 1$
 $\frac{f'(1) - f'(-1)}{1 - (-1)} = \frac{0 - 0}{2} = 0$

2 : $\begin{cases} 1 : f'(1) - f'(-1) = 0 \\ 1 : \text{explanation, using Mean Value Theorem} \end{cases}$

Therefore, by the Mean Value Theorem, there is at least one value c , $-1 < c < 1$, such that $f''(c) = 0$.

(c) $h'(x) = \frac{1}{f(x)} \cdot f'(x)$

$h'(3) = \frac{1}{f(3)} \cdot f'(3) = \frac{1}{7} \cdot \frac{1}{2} = \frac{1}{14}$

3 : $\begin{cases} 2 : h'(x) \\ 1 : \text{answer} \end{cases}$

(d) $\int_{-2}^3 f'(g(x))g'(x) dx = [f(g(x))]_{x=-2}^{x=3}$
 $= f(g(3)) - f(g(-2))$
 $= f(1) - f(-1)$
 $= 2 - 8 = -6$

3 : $\begin{cases} 2 : \text{Fundamental Theorem of Calculus} \\ 1 : \text{answer} \end{cases}$


2014 SCORING GUIDELINES

Question 6

Consider the differential equation $\frac{dy}{dx} = (3 - y)\cos x$. Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(0) = 1$. The function f is defined for all real numbers.

- (a) A portion of the slope field of the differential equation is given below. Sketch the solution curve through the point $(0, 1)$.
- (b) Write an equation for the line tangent to the solution curve in part (a) at the point $(0, 1)$. Use the equation to approximate $f(0.2)$.
- (c) Find $y = f(x)$, the particular solution to the differential equation with the initial condition $f(0) = 1$.

(a)



1 : solution curve

$$(b) \left. \frac{dy}{dx} \right|_{(x,y)=(0,1)} = 2 \cos 0 = 2$$

An equation for the tangent line is $y = 2x + 1$.

$$f(0.2) \approx 2(0.2) + 1 = 1.4$$

$$2 : \begin{cases} 1 : \text{tangent line equation} \\ 1 : \text{approximation} \end{cases}$$

$$(c) \frac{dy}{dx} = (3 - y)\cos x$$

$$\int \frac{dy}{3 - y} = \int \cos x \, dx$$

$$-\ln|3 - y| = \sin x + C$$

$$-\ln 2 = \sin 0 + C \Rightarrow C = -\ln 2$$

$$-\ln|3 - y| = \sin x - \ln 2$$

Because $y(0) = 1$, $y < 3$, so $|3 - y| = 3 - y$

$$3 - y = 2e^{-\sin x}$$

$$y = 3 - 2e^{-\sin x}$$

$$6 : \begin{cases} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$$

Note: max 3/6 [1-2-0-0-0] if no constant of integration

Note: 0/6 if no separation of variables