


Part B (AB or BC): Graphing calculator not allowed**Question 3****9 points****General Scoring Notes**

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Scoring guidelines and notes contain examples of the most common approaches seen in student responses. These guidelines can be applied to alternate approaches to ensure that these alternate approaches are scored appropriately.



A company designs spinning toys using the family of functions $y = cx\sqrt{4 - x^2}$, where c is a positive constant.

The figure above shows the region in the first quadrant bounded by the x -axis and the graph of $y = cx\sqrt{4 - x^2}$, for some c . Each spinning toy is in the shape of the solid generated when such a region is revolved about the x -axis. Both x and y are measured in inches.

Model Solution	Scoring
(a) Find the area of the region in the first quadrant bounded by the x -axis and the graph of $y = cx\sqrt{4 - x^2}$ for $c = 6$.	
$6x\sqrt{4 - x^2} = 0 \Rightarrow x = 0, x = 2$ $\text{Area} = \int_0^2 6x\sqrt{4 - x^2} \, dx$	Integrand 1 point
Let $u = 4 - x^2$. $du = -2x \, dx \Rightarrow -\frac{1}{2} du = x \, dx$ $x = 0 \Rightarrow u = 4 - 0^2 = 4$ $x = 2 \Rightarrow u = 4 - 2^2 = 0$ $\int_0^2 6x\sqrt{4 - x^2} \, dx = \int_4^0 6\left(-\frac{1}{2}\right)\sqrt{u} \, du = -3\int_4^0 u^{1/2} \, du = 3\int_0^4 u^{1/2} \, du$ $= 2u^{3/2} \Big _{u=0}^{u=4} = 2 \cdot 8 = 16$	Antiderivative 1 point
The area of the region is 16 square inches.	Answer 1 point

Scoring notes:

- Units are not required for any points in this question and are not read if presented (correctly or incorrectly) in any part of the response.
- The first point is earned for presenting $cx\sqrt{4 - x^2}$ or $6x\sqrt{4 - x^2}$ as the integrand in a definite integral. Limits of integration (numeric or alphanumeric) must be presented (as part of the definite integral) but do not need to be correct in order to earn the first point.
- If an indefinite integral is presented with an integrand of the correct form, the first point can be earned if the antiderivative (correct or incorrect) is eventually evaluated using the correct limits of integration.
- The second point can be earned without the first point. The second point is earned for the presentation of a correct antiderivative of a function of the form $Ax\sqrt{4 - x^2}$, for any nonzero constant A . If the response has subsequent errors in simplification of the antiderivative or sign errors, the response will earn the second point but will not earn the third point.
- Responses that use u -substitution and have incorrect limits of integration or do not change the limits of integration from x - to u -values are eligible for the second point.
- The response is eligible for the third point only if it has earned the second point.
- The third point is earned only for the answer 16 or equivalent. In the case where a response only presents an indefinite integral, the use of the correct limits of integration to evaluate the antiderivative must be shown to earn the third point.
- The response cannot correct -16 to $+16$ in order to earn the third point; there is no possible reversal here.

Total for part (a) 3 points

(b)

It is known that, for $y = cx\sqrt{4 - x^2}$, $\frac{dy}{dx} = \frac{c(4 - 2x^2)}{\sqrt{4 - x^2}}$. For a particular spinning toy, the radius of the largest cross-sectional circular slice is 1.2 inches. What is the value of c for this spinning toy?

The cross-sectional circular slice with the largest radius occurs where $cx\sqrt{4 - x^2}$ has its maximum on the interval $0 < x < 2$.

Sets $\frac{dy}{dx} = 0$ **1 point**

$$\frac{dy}{dx} = \frac{c(4 - 2x^2)}{\sqrt{4 - x^2}} = 0 \Rightarrow x = \sqrt{2}$$

$$x = \sqrt{2} \Rightarrow y = c\sqrt{2}\sqrt{4 - (\sqrt{2})^2} = 2c$$

$$2c = 1.2 \Rightarrow c = 0.6$$

Answer **1 point**

Scoring notes:

- The first point is earned for setting $\frac{dy}{dx} = 0$, $\frac{c(4 - 2x^2)}{\sqrt{4 - x^2}} = 0$, or $c(4 - 2x^2) = 0$.
- An unsupported $x = \sqrt{2}$ does not earn the first point.
- The second point can be earned without the first point but is earned only for the answer $c = 0.6$ with supporting work.

Total for part (b) 2 points

- (c) For another spinning toy, the volume is 2π cubic inches. What is the value of c for this spinning toy?

$\text{Volume} = \int_0^2 \pi \left(cx\sqrt{4-x^2} \right)^2 dx = \pi c^2 \int_0^2 x^2 (4-x^2) dx$	Form of the integrand	1 point
	Limits and constant	1 point
$= \pi c^2 \int_0^2 (4x^2 - x^4) dx = \pi c^2 \left(\frac{4}{3}x^3 - \frac{1}{5}x^5 \right) \Big _0^2$	Antiderivative	1 point
$= \pi c^2 \left(\frac{32}{3} - \frac{32}{5} \right) = \frac{64\pi c^2}{15}$ $\frac{64\pi c^2}{15} = 2\pi \Rightarrow c^2 = \frac{15}{32} \Rightarrow c = \sqrt{\frac{15}{32}}$	Answer	1 point

Scoring notes:

- The first point is earned for presenting an integrand of the form $A(x\sqrt{4-x^2})^2$ in a definite integral with any limits of integration (numeric or alphanumeric) and any nonzero constant A . Mishandling the c will result in the response being ineligible for the fourth point.
- The second point can be earned without the first point. The second point is earned for the limits of integration, $x = 0$ and $x = 2$, and the constant π (but not for 2π) as part of an integral with a correct or incorrect integrand.
- If an indefinite integral is presented with the correct constant π , the second point can be earned if the antiderivative (correct or incorrect) is evaluated using the correct limits of integration.
- A response that presents $2 = \int_0^2 (cx\sqrt{4-x^2})^2 dx$ earns the first and second points.
- The third point is earned for presenting a correct antiderivative of the presented integrand of the form $A(x\sqrt{4-x^2})^2$ for any nonzero A . If there are subsequent errors in simplification of the antiderivative, linkage errors, or sign errors, the response will not earn the fourth point.
- The fourth point cannot be earned without the third point. The fourth point is earned only for the correct answer. The expression does not need to be simplified to earn the fourth point.

Total for part (c) 4 points

Total for question 3 9 points

AP[®] Calculus AB

2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2016 SCORING GUIDELINES

Question 1

t (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.

- (a) Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.
- (b) Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
- (c) Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
- (d) For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

$$(a) \quad R'(2) \approx \frac{R(3) - R(1)}{3 - 1} = \frac{950 - 1190}{3 - 1} = -120 \text{ liters/hr}^2$$

$$2 : \begin{cases} 1 : \text{estimate} \\ 1 : \text{units} \end{cases}$$

(b) The total amount of water removed is given by $\int_0^8 R(t) dt$.

$$\begin{aligned} \int_0^8 R(t) dt &\approx 1 \cdot R(0) + 2 \cdot R(1) + 3 \cdot R(3) + 2 \cdot R(6) \\ &= 1(1340) + 2(1190) + 3(950) + 2(740) \\ &= 8050 \text{ liters} \end{aligned}$$

This is an overestimate since R is a decreasing function.

$$3 : \begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{estimate} \\ 1 : \text{overestimate with reason} \end{cases}$$

$$\begin{aligned} (c) \quad \text{Total} &\approx 50000 + \int_0^8 W(t) dt - 8050 \\ &= 50000 + 7836.195325 - 8050 \approx 49786 \text{ liters} \end{aligned}$$

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{estimate} \end{cases}$$

(d) $W(0) - R(0) > 0$, $W(8) - R(8) < 0$, and $W(t) - R(t)$ is continuous.

$$2 : \begin{cases} 1 : \text{considers } W(t) - R(t) \\ 1 : \text{answer with explanation} \end{cases}$$

Therefore, the Intermediate Value Theorem guarantees at least one time t , $0 < t < 8$, for which $W(t) - R(t) = 0$, or $W(t) = R(t)$.

For this value of t , the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank.

2016 SCORING GUIDELINES

Question 2

For $t \geq 0$, a particle moves along the x -axis. The velocity of the particle at time t is given by

$v(t) = 1 + 2\sin\left(\frac{t^2}{2}\right)$. The particle is at position $x = 2$ at time $t = 4$.

- At time $t = 4$, is the particle speeding up or slowing down?
- Find all times t in the interval $0 < t < 3$ when the particle changes direction. Justify your answer.
- Find the position of the particle at time $t = 0$.
- Find the total distance the particle travels from time $t = 0$ to time $t = 3$.

(a) $v(4) = 2.978716 > 0$
 $v'(4) = -1.164000 < 0$

The particle is slowing down since the velocity and acceleration have different signs.

(b) $v(t) = 0 \Rightarrow t = 2.707468$

$v(t)$ changes from positive to negative at $t = 2.707$.
 Therefore, the particle changes direction at this time.

(c) $x(0) = x(4) + \int_4^0 v(t) dt$
 $= 2 + (-5.815027) = -3.815$

(d) Distance $= \int_0^3 |v(t)| dt = 5.301$

2 : conclusion with reason

2 : $\begin{cases} 1 : t = 2.707 \\ 1 : \text{justification} \end{cases}$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$


2016 SCORING GUIDELINES

Question 3

The figure above shows the graph of the piecewise-linear function f . For $-4 \leq x \leq 12$, the function g is defined by

$$g(x) = \int_2^x f(t) \, dt.$$

- (a) Does g have a relative minimum, a relative maximum, or neither at $x = 10$? Justify your answer.
- (b) Does the graph of g have a point of inflection at $x = 4$? Justify your answer.
- (c) Find the absolute minimum value and the absolute maximum value of g on the interval $-4 \leq x \leq 12$. Justify your answers.
- (d) For $-4 \leq x \leq 12$, find all intervals for which $g(x) \leq 0$.

Graph of f

- (a) The function g has neither a relative minimum nor a relative maximum at $x = 10$ since $g'(x) = f(x)$ and $f(x) \leq 0$ for $8 \leq x \leq 12$.
- (b) The graph of g has a point of inflection at $x = 4$ since $g'(x) = f(x)$ is increasing for $2 \leq x \leq 4$ and decreasing for $4 \leq x \leq 8$.
- (c) $g'(x) = f(x)$ changes sign only at $x = -2$ and $x = 6$.

x	$g(x)$
-4	-4
-2	-8
6	8
12	-4

On the interval $-4 \leq x \leq 12$, the absolute minimum value is $g(-2) = -8$ and the absolute maximum value is $g(6) = 8$.

- (d) $g(x) \leq 0$ for $-4 \leq x \leq 2$ and $10 \leq x \leq 12$.

1 : $g'(x) = f(x)$ in (a), (b), (c), or (d)

1 : answer with justification

1 : answer with justification

4 : $\left\{ \begin{array}{l} 1 : \text{considers } x = -2 \text{ and } x = 6 \\ \quad \text{as candidates} \\ 1 : \text{considers } x = -4 \text{ and } x = 12 \\ 2 : \text{answers with justification} \end{array} \right.$


2 : intervals

2016 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = \frac{y^2}{x-1}$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.
- (b) Let $y = f(x)$ be the particular solution to the given differential equation with the initial condition $f(2) = 3$. Write an equation for the line tangent to the graph of $y = f(x)$ at $x = 2$. Use your equation to approximate $f(2.1)$.
- (c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(2) = 3$.



2 : $\begin{cases} 1 : \text{zero slopes} \\ 1 : \text{nonzero slopes} \end{cases}$

(b) $\left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} = \frac{3^2}{2-1} = 9$

2 : $\begin{cases} 1 : \text{tangent line equation} \\ 1 : \text{approximation} \end{cases}$

An equation for the tangent line is $y = 9(x - 2) + 3$.

$$f(2.1) \approx 9(2.1 - 2) + 3 = 3.9$$

(c) $\frac{1}{y^2} dy = \frac{1}{x-1} dx$

$$\int \frac{1}{y^2} dy = \int \frac{1}{x-1} dx$$

$$-\frac{1}{y} = \ln|x-1| + C$$

$$-\frac{1}{3} = \ln|2-1| + C \Rightarrow C = -\frac{1}{3}$$

$$-\frac{1}{y} = \ln|x-1| - \frac{1}{3}$$

$$y = \frac{1}{\frac{1}{3} - \ln(x-1)}$$

5 : $\begin{cases} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration and} \\ \quad \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$


Note: max 3/5 [1-2-0-0] if no constant of integration

Note: 0/5 if no separation of variables

Note: This solution is valid for $1 < x < 1 + e^{1/3}$.

2016 SCORING GUIDELINES

Question 5



The inside of a funnel of height 10 inches has circular cross sections, as shown in the figure above. At height h , the radius of the funnel is given by $r = \frac{1}{20}(3 + h^2)$, where $0 \leq h \leq 10$. The units of r and h are inches.

- (a) Find the average value of the radius of the funnel.
- (b) Find the volume of the funnel.
- (c) The funnel contains liquid that is draining from the bottom. At the instant when the height of the liquid is $h = 3$ inches, the radius of the surface of the liquid is decreasing at a rate of $\frac{1}{5}$ inch per second. At this instant, what is the rate of change of the height of the liquid with respect to time?

$$\begin{aligned} \text{(a) Average radius} &= \frac{1}{10} \int_0^{10} \frac{1}{20}(3 + h^2) dh = \frac{1}{200} \left[3h + \frac{h^3}{3} \right]_0^{10} \\ &= \frac{1}{200} \left(\left(30 + \frac{1000}{3} \right) - 0 \right) = \frac{109}{60} \text{ in} \end{aligned}$$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b) Volume} &= \pi \int_0^{10} \left(\left(\frac{1}{20} \right) (3 + h^2) \right)^2 dh = \frac{\pi}{400} \int_0^{10} (9 + 6h^2 + h^4) dh \\ &= \frac{\pi}{400} \left[9h + 2h^3 + \frac{h^5}{5} \right]_0^{10} \\ &= \frac{\pi}{400} \left(\left(90 + 2000 + \frac{100000}{5} \right) - 0 \right) = \frac{2209\pi}{40} \text{ in}^3 \end{aligned}$$

3 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(c) } \frac{dr}{dt} &= \frac{1}{20}(2h) \frac{dh}{dt} \\ -\frac{1}{5} &= \frac{3}{10} \frac{dh}{dt} \\ \frac{dh}{dt} &= -\frac{1}{5} \cdot \frac{10}{3} = -\frac{2}{3} \text{ in/sec} \end{aligned}$$

3 : $\begin{cases} 2 : \text{chain rule} \\ 1 : \text{answer} \end{cases}$

2016 SCORING GUIDELINES

Question 6

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	-6	3	2	8
2	2	-2	-3	0
3	8	7	6	2
6	4	5	3	-1

The functions f and g have continuous second derivatives. The table above gives values of the functions and their derivatives at selected values of x .

(a) Let $k(x) = f(g(x))$. Write an equation for the line tangent to the graph of k at $x = 3$.

(b) Let $h(x) = \frac{g(x)}{f(x)}$. Find $h'(1)$.

(c) Evaluate $\int_1^3 f''(2x) dx$.

$$\begin{aligned} \text{(a)} \quad k(3) &= f(g(3)) = f(6) = 4 \\ k'(3) &= f'(g(3)) \cdot g'(3) = f'(6) \cdot 2 = 5 \cdot 2 = 10 \end{aligned}$$

An equation for the tangent line is $y = 10(x - 3) + 4$.

$$\begin{aligned} \text{(b)} \quad h'(1) &= \frac{f(1) \cdot g'(1) - g(1) \cdot f'(1)}{(f(1))^2} \\ &= \frac{(-6) \cdot 8 - 2 \cdot 3}{(-6)^2} = \frac{-54}{36} = -\frac{3}{2} \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad \int_1^3 f''(2x) dx &= \frac{1}{2} [f'(2x)]_1^3 = \frac{1}{2} [f'(6) - f'(2)] \\ &= \frac{1}{2} [5 - (-2)] = \frac{7}{2} \end{aligned}$$

3 : $\begin{cases} 2 : \text{slope at } x = 3 \\ 1 : \text{equation for tangent line} \end{cases}$

3 : $\begin{cases} 2 : \text{expression for } h'(1) \\ 1 : \text{answer} \end{cases}$

3 : $\begin{cases} 2 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$