

Chain Rule, Implicit & Inverse

AP Calculus AB • Unit 3


AP Calculus AB



Chain Rule, Implicit & Inverse

What we will cover

- 1 Composite functions
- 2 The chain rule
- 3 Implicit differentiation
- 4 Derivatives of inverse functions
- 5 Inverse trig derivatives
- 6 Higher-order derivatives & strategy



1 Chain rule

Chain Rule, Implicit & Inverse

Chain rule

What makes a function composite?

A **composite function** 复合函数 $f(g(x))$ nests one function **inside** another.

Always ask two questions: **what is the inside**, and **what is its derivative**?



Chain rule
matryoshka

Chain Rule, Implicit & Inverse

Chain rule

The chain rule

Differentiate **outer-then-inner**:

$$\frac{d}{dx}f(g(x)) = f'(g(x)) \cdot g'(x).$$

In Leibniz notation, with $y = f(u)$ and $u = g(x)$:

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}.$$

The inner derivative $g'(x)$ is the **most-forgotten** factor on the exam.

differentiate the **outer**,
leave the in-
multiply by the
derivative of
the **inside**
 $\cos(x^2) \cdot 2x$

Chain Rule, Implicit & Inverse


Chain rule

Worked example – a power of a polynomial

Differentiate $h(x) = (2x^2 + 1)^5$

Outer is $(\cdot)^5$, inner is $2x^2 + 1$:

$$\begin{aligned} h'(x) &= 5(2x^2 + 1)^4 \cdot \frac{d}{dx}(2x^2 + 1) \\ &= 5(2x^2 + 1)^4 \cdot 4x = 20x(2x^2 + 1)^4. \end{aligned}$$



Exam skill – the chain rule from a table

A table gives values; no formula is needed. With $h(x) = f(g(x))$:

$$h'(a) = f'(g(a)) \cdot g'(a).$$

Read $g(a)$ **first**, then look up f' at **that** value – not at a .

Find $h'(1)$

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	6	4	2	5
2	3	7	1	9

$$g(1) = 2, \text{ so } h'(1) = f'(2) \cdot g'(1) = 7 \times 5 = 35.$$

2 Implicit

Implicit differentiation

Some curves are defined **implicitly** 隐式地 – an equation in x and y not solved for y , e.g. $x^2 + y^2 = 25$.

- 1 differentiate **both sides** in x
- 2 each y -term picks up a $\frac{dy}{dx}$ (chain rule)
- 3 solve algebraically for $\frac{dy}{dx}$

$$x^2 + y^2 = 25$$

$$2x + 2y \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = -\frac{x}{y}$$

Worked example – tangent to a circle

Tangent to $x^2 + y^2 = 25$ at $(3, 4)$


From $\frac{dy}{dx} = -\frac{x}{y}$, at $(3, 4)$:

$$\frac{dy}{dx} = -\frac{3}{4}.$$

So the tangent is

$$y - 4 = -\frac{3}{4}(x - 3).$$

The tangent is **perpendicular** to the **radius** – a free sanity check.



Exam skill – “Show that $\frac{dy}{dx} = \dots$ ”

The target is **given**, so the marks are in the **algebra**, not the answer:

- 1 differentiate both sides
- 2 product/chain rule on mixed xy terms
- 3 collect every $\frac{dy}{dx}$ on one side
- 4 factor, then divide

A correct final line that **skips** the algebra earns little.

Horizontal tangent
 $\frac{dy}{dx} = 0$: **numerator** = 0

Vertical tangent
denominator = 0


3 Inverse functions

Derivatives of inverse functions

If g is the **inverse function** 反函数 of f (so $f(g(x)) = x$):

$$g'(x) = \frac{1}{f'(g(x))}.$$

Slopes at **matching points** are **reciprocals** 倒数 – the graphs are mirror images in $y = x$.




Worked example – inverse from a point

$f(2) = 5$ and $f'(2) = 3$; g is the inverse of f

$(2, 5)$ lies on f , so $(5, 2)$ lies on g . Then

$$g'(5) = \frac{1}{f'(2)} = \frac{1}{3}.$$

Swap the coordinates **first**, then take the reciprocal of f' at the **original** x .



Inverse trigonometric derivatives

The same idea gives the **inverse trigonometric functions** 反三角函数 you must know:

$$\frac{d}{dx} \arcsin x = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \arctan x = \frac{1}{1+x^2}$$

$$\frac{d}{dx} \operatorname{arcsec} x = \frac{1}{|x|\sqrt{x^2-1}}$$

Chain them when the input is a function: $\frac{d}{dx} \arctan(3x) = \frac{3}{1+9x^2}$.

4


Higher-order & strategy

Higher-order derivatives

Differentiating f' gives the **second derivative** 二阶导数 f'' :


$$f''(x) = \frac{d^2 y}{dx^2} = y'', \quad f^{(n)}(x) = \frac{d^n y}{dx^n}.$$

f'' measures how the **slope** is changing – it drives **concavity** 凹凸性 (Unit 5) and acceleration (Unit 4).



Choosing the right rule

Read the structure **from the outside in**: name the outermost operation, apply its rule, recurse inward.



Neatness prevents the sign and bookkeeping errors that cost the marks.

5 Recap

Chain Rule, Implicit & Inverse
↳ Recap

Unit 3 – key takeaways

1 Chain rule

$f(g(x)) \cdot g'(x)$ – never drop the inner derivative

2 Implicit

differentiate both sides; each y gives a dy/dx ; then solve

3 Inverse

$g'(x) = 1/f'(g(x))$ – reciprocal slopes at swapped points

4 Second derivative

differentiate twice; velocity $\rightarrow$ acceleration

Chain Rule, Implicit & Inverse
↳ Recap


Check yourself

1 Differentiate $\cos(3x)$.

2 For $x^2 + y^2 = 25$, what is $\frac{dy}{dx}$ at $(3, 4)$?

3 $f(2) = 5$, $f'(2) = 3$. If $g = f^{-1}$, what is $g'(5)$?

4 Which factor do students most often forget?



Answers 1. $-3 \sin(3x)$ 2. $-\frac{3}{4}$ 3. $\frac{1}{3}$ 4. the inner derivative $g'(x)$