

1. An invasive species of plant appears in a fruit grove at time $t = 0$ and begins to spread. The function C defined by $C(t) = 7.6 \arctan(0.2t)$ models the number of acres in the fruit grove affected by the species t weeks after the species appears. It can be shown that $C'(t) = \frac{38}{25 + t^2}$.

(Note: Your calculator should be in radian mode.)


- A. Find the average number of acres affected by the invasive species from time $t = 0$ to time $t = 4$ weeks. Show the setup for your calculations.
- B. Find the time t when the instantaneous rate of change of C equals the average rate of change of C over the time interval $0 \leq t \leq 4$. Show the setup for your calculations.
- C. Assume that the invasive species continues to spread according to the given model for all times $t > 0$. Write a limit expression that describes the end behavior of the rate of change in the number of acres affected by the species. Evaluate this limit expression.
- D. At time $t = 4$ weeks after the invasive species appears in the fruit grove, measures are taken to counter the spread of the species. The function A , defined by $A(t) = C(t) - \int_4^t 0.1 \cdot \ln(x) dx$, models the number of acres affected by the species over the time interval $4 \leq t \leq 36$. At what time t , for $4 \leq t \leq 36$, does A attain its maximum value? Justify your answer.

3. The depth of seawater at a location can be modeled by the function H that satisfies the differential equation

$$\frac{dH}{dt} = \frac{1}{2}(H - 1)\cos\left(\frac{t}{2}\right), \text{ where } H(t) \text{ is measured in feet and } t \text{ is measured in hours after noon } (t = 0). \text{ It is}$$

known that $H(0) = 4$.

- (a) A portion of the slope field for the differential equation is provided. Sketch the solution curve, $y = H(t)$, through the point $(0, 4)$.



- (b) For $0 < t < 5$, it can be shown that $H(t) > 1$. Find the value of t , for $0 < t < 5$, at which H has a critical point. Determine whether the critical point corresponds to a relative minimum, a relative maximum, or neither a relative minimum nor a relative maximum of the depth of seawater at the location. Justify your answer.
- (c) Use separation of variables to find $y = H(t)$, the particular solution to the differential equation

$$\frac{dH}{dt} = \frac{1}{2}(H - 1)\cos\left(\frac{t}{2}\right) \text{ with initial condition } H(0) = 4.$$

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

x	0	2	4	7
$f(x)$	10	7	4	5
$f'(x)$	$\frac{3}{2}$	-8	3	6
$g(x)$	1	2	-3	0
$g'(x)$	5	4	2	8

5. The functions f and g are twice differentiable. The table shown gives values of the functions and their first derivatives at selected values of x .

- (a) Let h be the function defined by $h(x) = f(g(x))$. Find $h'(7)$. Show the work that leads to your answer.
- (b) Let k be a differentiable function such that $k'(x) = (f(x))^2 \cdot g(x)$. Is the graph of k concave up or concave down at the point where $x = 4$? Give a reason for your answer.
- (c) Let m be the function defined by $m(x) = 5x^3 + \int_0^x f'(t) dt$. Find $m(2)$. Show the work that leads to your answer.
- (d) Is the function m defined in part (c) increasing, decreasing, or neither at $x = 2$? Justify your answer.

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.