# 8.11 Volume with Washer Method: Revolving Around the x- or y-Axis

Name: _____ Class: _____ Date: _____ **Total: 14 marks**

**KEY WORDS** washer method

## Objective

Build the skills to answer exam questions on the **washer method** – volumes of revolution with a hole.

**You must be able to:**

- identify the **outer** and **inner** radii of a washer (ring)
- integrate $\pi\left(R_{\text{outer}}^2 - R_{\text{inner}}^2\right)$ (the **washer method** )
- set each radius as a distance from a curve to the axis

## 1 Key knowledge and worked examples

Study these first. Each one shows the method for a question type used later – follow the steps and you can do the Practice and Exam-style questions yourself.

**When there is a hole**

If the region does **not** touch the axis, revolving it leaves a hole, so each slice is a **washer** (a ring). Its area is (outer disc) − (inner disc):

$$V = \pi \int_a^b \left(R_{\text{outer}}^2 - R_{\text{inner}}^2\right) dx.$$

**Identifying the radii**

Revolve the region between $y = \sqrt{x}$ (top) and $y = x$ (bottom) on $[0, 1]$ about the $x$-axis. Outer radius (further curve) $R = \sqrt{x}$; inner radius $r = x$:

$$V = \pi \int_0^1 \left((\sqrt{x})^2 - x^2\right) dx = \pi \int_0^1 (x - x^2)\, dx = \pi\left(\tfrac{1}{2} - \tfrac{1}{3}\right) = \tfrac{\pi}{6}.$$

**Do not square then subtract wrongly**

$R^2 - r^2$ is **not** $(R - r)^2$. Square each radius separately, then subtract.

**Choosing outer vs inner**

The **outer** radius comes from the curve **farther** from the axis; the **inner** from the curve **nearer** the axis. A quick sketch settles which is which.

## 2 Practice

Now apply the methods above.

**2.1** State the washer-method formula for the volume. [1]

**2.2** For $R = 4$ and $r = 2$, find $\pi(R^2 - r^2)$. [1]

**2.3** Evaluate $\pi \int_0^1 (x - x^2)\, dx$. [2]

## 3 Exam-style questions

**3.1** The washer method integrates [1]

- ☐ **A** $\pi \int (R - r)^2\, dx$
- **B** $\pi \int (R^2 - r^2)\, dx$
- **C** $\pi \int R^2\, dx$
- **D** $\int (R^2 - r^2)\, dx$

**3.2** The region between $y = x$ and $y = x^2$ (from $x = 0$ to $x = 1$) is revolved about the $x$-axis.

(a) Identify the outer and inner radii. [2]

(b) Set up and evaluate the volume. [4]

**3.3** The region bounded by $y = 4$ and $y = x^2$ (for $0 \leq x \leq 2$) is revolved about the $x$-axis. Set up the washer integral for the volume. [3]

## Solutions

Each answer shows the steps, then the **result**.

**2.1**

- $V = \pi \int_a^b \left(R_{\text{outer}}^2 - R_{\text{inner}}^2\right) dx$

**2.2**

- $\pi(16 - 4) = 12\pi$

**2.3**

- $\pi\left[\frac{x^2}{2} - \frac{x^3}{3}\right]_0^1 = \pi\left(\frac{1}{2} - \frac{1}{3}\right) = \frac{\pi}{6}$

**3.1 B**

- subtract the **squares** of the radii

**3.2**

(a)

- On $(0, 1)$, $y = x$ is above $y = x^2$; outer $R = x$, inner $r = x^2$

(b)

- $V = \pi \int_0^1 (x^2 - x^4)\, dx = \pi\left[\frac{x^3}{3} - \frac{x^5}{5}\right]_0^1 = \pi\left(\frac{1}{3} - \frac{1}{5}\right) = \frac{2\pi}{15}$

**3.3**

- Outer $R = 4$, inner $r = x^2$; $V = \pi \int_0^2 \left(4^2 - (x^2)^2\right) dx = \pi \int_0^2 (16 - x^4)\, dx$