9.9 Finding the Area of the Region Bounded by Two Polar Curves

Name: _____ Class: _____ Date: _____ **Total: 36 marks**

KEY WORDS meeting point • inside • outside • cardioid • pole • region

Objective

Build the skills to answer exam questions on **the area between two polar curves** : finding where the curves meet, $\frac{1}{2}\int(R^2-r^2)\,d\theta$, and regions inside one curve and outside another.

You must be able to:

- find the angles where two polar curves meet by solving $f(\theta)=g(\theta)$
- write the area inside $R=f(\theta)$ and outside $r=g(\theta)$ as $\frac{1}{2}\int(R^2-r^2)\,d\theta$
- evaluate such integrals with the double-angle identities
- build an area from simpler areas by adding or subtracting

1 Key knowledge and worked examples

■ Area between polar curves

idea	rule
meeting points	solve $f(\theta)=g(\theta)$; also check the pole, where $r=0$ on each curve
inside R, outside r	$A=\dfrac{1}{2}\displaystyle\int_{\alpha}^{\beta}\left(R^2-r^2\right)d\theta$, where $R\geq r$ on $[\alpha,\beta]$
not	$\frac{1}{2}\int(R-r)^2\,d\theta$
limits	the angles of the meeting points that bound the region
build up	inside both = one whole area − the part outside the other
identities	$\sin^2\theta=\frac{1}{2}(1-\cos 2\theta)$, $\cos^2\theta=\frac{1}{2}(1+\cos 2\theta)$

exercises.lol exercises.com.cn

Worked example 1: inside a circle, outside another. Inside $r=2\sin\theta$ and outside $r=1$.

1. Known: $2\sin\theta=1$ at $\theta=\frac{\pi}{6}$ and $\frac{5\pi}{6}$.
2. Rule: $\dfrac{1}{2}\displaystyle\int_{\pi/6}^{5\pi/6}(4\sin^2\theta-1)\,d\theta=\frac{1}{2}\int_{\pi/6}^{5\pi/6}(1-2\cos 2\theta)\,d\theta$.
3. Numbers: $\dfrac{1}{2}\Big[\theta-\sin 2\theta\Big]_{\pi/6}^{5\pi/6}=\dfrac{1}{2}\left(\dfrac{2\pi}{3}+\sqrt{3}\right)$.
4. Result: $\dfrac{\pi}{3}+\dfrac{\sqrt{3}}{2}$.


Worked example 2: meeting points. Where do $r=1$ and $r=2\cos\theta$ meet, $-\frac{\pi}{2}\leq\theta\leq\frac{\pi}{2}$?

1. Known: $2\cos\theta=1$.
2. Rule: $\cos\theta=\frac{1}{2}$.
3. Numbers: $\theta=\pm\frac{\pi}{3}$, with $r=1$.
4. Result: $\left(\frac{1}{2},\,\pm\frac{\sqrt{3}}{2}\right)$ in Cartesian form.


Answer shapes: sketch or read the figure to decide which curve is outside; write the limits next to the integral.

2 Practice


2.1 Find the area of the ring between the circles $r=3$ and $r=1$. [2]



2.2 Find the points where the circles $r=1$ and $r=2\cos\theta$ meet. Give them in polar and in Cartesian form. [3]




2.3 Find the area of the region inside $r = 2$ and outside $r = 2\sin\theta$, for $0 \le \theta \le \frac{\pi}{2}$. [3]



3 Exam-style questions


3.1 Find the exact area of the region inside $r = 4\cos\theta$ and outside $r = 2$. [4]



3.2 The cardioid $r = 1 + \sin\theta$ and the circle $r = 1$ are shown.

(a) Find the values of θ in $[0, 2\pi)$ where the curves meet. [2]

(b) Find the area inside the cardioid and outside the circle. [3]




(c) Find the area inside the circle and outside the cardioid. [1]

3.3 The circle $r = 3\sin\theta$ and the cardioid $r = 1 + \sin\theta$ are shown. They meet at two points and at the pole.

(a) Find the Cartesian coordinates of the two points where the curves meet, other than the pole. [3]

(b) Find the area of the region inside the circle and outside the cardioid. [4]



exercises.lol exercises.com.cn

(c) Find the area of the region inside both curves. Give the exact value and a decimal. [5]

(d) Find the area of the region inside the cardioid and outside the circle. [6]

exercises.lol exercises.com.cn