

AP Calculus BC

Name: _____ Score: _____

1. The slope of a **secant line** between two points on a curve gives the...
- ☐ average rate of change over that interval
 - ☐ exact value of the function
 - ☐ area under the curve
 - ☐ instantaneous rate of change
2. The notation $\lim_{x \rightarrow c} f(x) = L$ means that as $x \dots$
- ☐ is greater than c , $f(x)$ equals L
 - ☐ equals c , the output is exactly L
 - ☐ grows without bound, $f(x)$ reaches L
 - ☐ gets close to c (without equalling it), the output heads toward L
3. A table gives $f(2.9) = 6.9$, $f(2.99) = 6.99$, $f(3.01) = 7.01$, $f(3.1) = 7.1$. Estimate $\lim_{x \rightarrow 3} f(x)$.
- _____
4. Evaluate $\lim_{x \rightarrow 4} \frac{x^2 - 16}{x - 4}$.
- _____
5. Direct substitution into $\lim_{x \rightarrow 4} \frac{x - 4}{x^2 - 16}$ gives $\frac{0}{0}$. What next?
- ☐ factor and cancel the common $(x - 4)$
 - ☐ declare it does not exist
 - ☐ report the limit as 0
 - ☐ multiply by a conjugate
6. A graph shows the curve heading to height 3 from both sides of $x = 2$, with a filled dot at $(2, 7)$. Which statement matches?
- ☐ $\lim_{x \rightarrow 2} f(x) = 3$ and $f(2) = 7$
 - ☐ the limit does not exist
 - ☐ $\lim_{x \rightarrow 2} f(x) = 7$
 - ☐ $f(2) = 3$
7. Select **all** conditions required for f to be continuous at c .
- ☐ f is a polynomial

- ☐ $\lim_{x \rightarrow c} f(x)$ exists
- ☐ $f(c)$ is defined
- ☐ $\lim_{x \rightarrow c} f(x) = f(c)$

Tick every correct option.

8. At the **left** endpoint a of a closed interval $[a, b]$, continuity requires...

- ☐ nothing – endpoints are ignored
- ☐ $f(a) = 0$
- ☐ $\lim_{x \rightarrow a^-} f(x) = f(a)$
- ☐ $\lim_{x \rightarrow a^+} f(x) = f(a)$ (right-continuity)

9. At what x -value does $f(x) = \frac{x+1}{x-4}$ have a vertical asymptote?

10. To apply the IVT on $[a, b]$ with target N , you need **all** of...

- ☐ $f(a)$ and $f(b)$ known
- ☐ f continuous on $[a, b]$
- ☐ N between $f(a)$ and $f(b)$
- ☐ f differentiable on $[a, b]$

Tick every correct option.

AP Calculus BC

1. **average rate of change over that interval**

A secant joins two points, so its slope is the **average rate of change** over that interval.

2. **gets close to c (without equalling it), the output heads toward L**

$x \rightarrow c$ means x **approaches** c without touching it, and $f(x)$ heads toward L .

3. **7**

Both sides close in on 7, so the limit is 7.

4. **8**

Factor: $\frac{(x-4)(x+4)}{x-4} = x+4 \rightarrow 8$.

5. **factor and cancel the common $(x-4)$**

Polynomial over polynomial, both zero at 4 $\rightarrow$ factor: $\frac{x-4}{(x-4)(x+4)} = \frac{1}{x+4} \rightarrow \frac{1}{8}$.

6. **$\lim_{x \rightarrow 2} f(x) = 3$ and $f(2) = 7$**

The curve's approach gives the limit 3; the filled dot gives the separate value $f(2) = 7$.

7. **$\lim_{x \rightarrow c} f(x)$ exists; $f(c)$ is defined; $\lim_{x \rightarrow c} f(x) = f(c)$**

The three conditions are defined, limit-exists, and equal. Being a polynomial is not required.

8. **$\lim_{x \rightarrow a^+} f(x) = f(a)$ (right-continuity)**

You can only approach the left endpoint from the right, so right-continuity is required.

9. **4**

Denominator zero at $x = 4$, numerator $= 5 \neq 0 \rightarrow$ asymptote $x = 4$.

10. **$f(a)$ and $f(b)$ known; f continuous on $[a, b]$; N between $f(a)$ and $f(b)$**

IVT needs continuity and a target in range. Differentiability is **not** required.