

8.13 The Arc Length of a Smooth Curve and Distance Traveled

Name: _____ Class: _____ Date: _____

Total: 14 marks

Objective

Build the skills to answer exam questions on **arc length** —the length of a smooth curve.

You must be able to:

- apply $L = \int_a^b \sqrt{1 + (f'(x))^2} dx$
- set up the integral from a given function
- recognise the same idea as **distance travelled** along a path

1 Worked examples

Study these first. Each one shows the method for a question type used later —follow the steps and you can do the Practice and Exam-style questions yourself.

■ The arc-length formula

The length of the curve $y = f(x)$ from $x = a$ to $x = b$ is

$$L = \int_a^b \sqrt{1 + (f'(x))^2} dx.$$

It adds up tiny hypotenuses $\sqrt{dx^2 + dy^2}$ along the curve.

■ Setting up the integral

For $y = \frac{2}{3}x^{3/2}$: $f'(x) = x^{1/2}$, so $(f')^2 = x$, and

$$L = \int_0^3 \sqrt{1 + x} dx.$$

■ Evaluating

$$\int_0^3 \sqrt{1 + x} dx = \left[\frac{2}{3}(1 + x)^{3/2} \right]_0^3 = \frac{2}{3}(8 - 1) = \frac{14}{3}.$$

■ Distance along a path

If a particle moves along $y = f(x)$, the **distance travelled** is the same arc-length integral. Many arc-length integrals must be evaluated numerically —set them up correctly even when you cannot integrate by hand.

2 Practice

Now apply the methods above.

2.1 State the arc-length formula for $y = f(x)$ on $[a, b]$. [1]

2.2 For $y = \frac{2}{3}x^{3/2}$, find $f'(x)$ and $(f'(x))^2$. [2]

2.3 Set up (do not evaluate) the arc length of $y = x^2$ from $x = 0$ to $x = 2$. [2]

3 Exam-style questions

3.1 The arc length of $y = f(x)$ on $[a, b]$ is [1]

- **A** $\int_a^b f(x) dx$
 - **B** $\int_a^b \sqrt{1 + (f'(x))^2} dx$
 - **C** $\int_a^b f'(x) dx$
 - **D** $\pi \int_a^b (f(x))^2 dx$
-

3.2 A curve is $y = \frac{2}{3}x^{3/2}$ for $0 \leq x \leq 3$.

(a) Set up the arc-length integral. [2]

(b) Evaluate it. [3]

3.3 Set up the integral for the length of $y = \ln(\cos x)$ from $x = 0$ to $x = \frac{\pi}{4}$. (Use $\frac{d}{dx} \ln(\cos x) = -\tan x$ and $1 + \tan^2 x = \sec^2 x$.) [3]

4 Go further

You are now ready for the real exam questions on this subtopic:

- work through the **8.13 The Arc Length of a Smooth Curve and Distance Traveled** lesson on the **Learn** page;
- read the **The Arc Length of a Smooth Curve and Distance Traveled** section of the AP Calculus BC handout on the **Know** page.

Solutions

2.1 $L = \int_a^b \sqrt{1 + (f'(x))^2} dx.$

2.2 $f'(x) = x^{1/2}; (f'(x))^2 = x.$

2.3 $f'(x) = 2x$, so $L = \int_0^2 \sqrt{1 + 4x^2} dx.$

3.1 B —arc length integrates $\sqrt{1 + (f')^2}.$

3.2 (a) $f'(x) = x^{1/2}$, so $L = \int_0^3 \sqrt{1 + x} dx.$ (b) $[\frac{2}{3}(1 + x)^{3/2}]_0^3 = \frac{2}{3}(8 - 1) = \frac{14}{3}.$

3.3 $f'(x) = -\tan x$, so $1 + (f')^2 = 1 + \tan^2 x = \sec^2 x$; $L = \int_0^{\pi/4} \sqrt{\sec^2 x} dx = \int_0^{\pi/4} \sec x dx.$