

5.12 Exploring Behaviors of Implicit Relations

Name: _____ Class: _____ Date: _____ **Total: 10 marks**

KEY WORDS horizontal tangent 水平切线 • vertical tangent 垂直切线 • concavity 凹凸性
 • concave down 下凹 • concave up 上凹

Objective

Build the skills to answer exam questions on the **behavior of implicit relations** — using $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ found by implicit differentiation to locate features of a curve.

You must be able to:

- find where an implicit curve has a **horizontal tangent** 水平切线 ($\frac{dy}{dx} = 0$) or a **vertical tangent** 垂直切线 ($\frac{dy}{dx}$ undefined)
- decide **concavity** 凹凸性 from the sign of $\frac{d^2y}{dx^2}$ on the curve
- evaluate these using a point that lies **on** the curve

1 Worked examples

Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

■ Horizontal tangents

For $x^2 + y^2 = 25$, implicit differentiation gives $2x + 2y \frac{dy}{dx} = 0$, so

$$\frac{dy}{dx} = -\frac{x}{y}.$$

A **horizontal** tangent needs $\frac{dy}{dx} = 0$, i.e. the numerator $x = 0$ (with $y \neq 0$): the points $(0, 5)$ and $(0, -5)$.

■ Vertical tangents

A **vertical** tangent occurs where $\frac{dy}{dx}$ is undefined — the denominator zero, $y = 0$ (with $x \neq 0$): the points $(5, 0)$ and $(-5, 0)$. This matches the circle's shape.

■ Concavity on an implicit curve

Differentiate $\frac{dy}{dx} = -\frac{x}{y}$ again (quotient + chain rule):

$$\frac{d^2y}{dx^2} = -\frac{y - x \frac{dy}{dx}}{y^2} = -\frac{y - x(-x/y)}{y^2} = -\frac{y^2 + x^2}{y^3} = -\frac{25}{y^3}.$$

Where $y > 0$ this is negative (**concave down**, the top of the circle); where $y < 0$ it is positive (**concave up**).

2 Practice

Now apply the methods above.

2.1 For $x^2 + y^2 = 25$, state $\frac{dy}{dx}$. [1]

2.2 Find the point(s) on $x^2 + y^2 = 25$ where the tangent is **horizontal**. [2]

2.3 A curve satisfies $\frac{dy}{dx} = \frac{2x}{y}$. State where its tangent line is **vertical**. [1]

3 Exam-style questions

3.1 A curve has $\frac{dy}{dx} = \frac{x-1}{y}$. A horizontal tangent occurs where [1]

- | | |
|---|---|
| A | B |
| C | D |
- A** $y = 0$
B $x = 0$
C $x = 1$
D $x = y$

3.2 The curve $x^2 + xy + y^2 = 7$ passes through $(1, 2)$. Its derivative is $\frac{dy}{dx} = -\frac{2x + y}{x + 2y}$.

(a) Find the slope of the tangent at $(1, 2)$. [2]

(b) Determine whether the tangent at $(1, 2)$ is horizontal, vertical, or neither. [1]

3.3 For $x^2 + y^2 = 25$ it can be shown that $\frac{d^2y}{dx^2} = -\frac{25}{y^3}$. State, with a reason, whether the curve is concave up or concave down at the point $(3, 4)$. [2]