

6.5 Interpreting the Behavior of Accumulation Functions

Name: _____ Class: _____ Date: _____ **Total: 9 marks****KEY WORDS** local extrema 极值 • accumulation function 累积函数 • accumulation 累积

Objective

Build the skills to answer exam questions on the **behavior of accumulation functions** — reading features of $g(x) = \int_a^x f(t) dt$ from a graph of f .

You must be able to:

- decide where g is **increasing/decreasing** from the sign of f
- locate **local extrema** 极值 of g where f crosses zero (sign change)
- decide **concavity** of g from whether f is increasing or decreasing

1 Worked examples

Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

■ g rises where f is positive

Because $g'(x) = f(x)$, the accumulation function g **increases** on intervals where the graph of f is **above** the axis and **decreases** where f is **below** the axis.

■ Extrema of g at zeros of f

g has a **local maximum** where f changes from $+$ to $-$, and a **local minimum** where f changes from $-$ to $+$. A zero of f where the sign does **not** change is not an extremum of g .

■ Concavity of g from the slope of f

$g''(x) = f'(x)$, so g is **concave up** where f is **increasing** and **concave down** where f is **decreasing**. A point where f has a local max or min is a **point of inflection** of g .

■ Comparing values of g

To compare $g(3)$ and $g(0)$, look at the net signed area of f between them: $g(3) - g(0) = \int_0^3 f dt$. More area above the axis than below makes $g(3) > g(0)$.

2 Practice

Now apply the methods above. Let $g(x) = \int_0^x f(t) dt$, where f is continuous.

2.1 On an interval where $f(t) < 0$, state whether g is increasing or decreasing. [1]

2.2 At a point where f changes from positive to negative, what feature does g have? [1]

2.3 On an interval where f is increasing, state the concavity of g . [1]

3 Exam-style questions

3.1 $g(x) = \int_0^x f(t) dt$. The graph of f crosses from below the axis to above it at $x = 4$. At $x = 4$, g has a [1]

A	B
C	D

A local maximum

B local minimum

C point of inflection

D vertical asymptote

3.2 The graph of f consists of straight segments: $f > 0$ on $(0, 3)$, $f < 0$ on $(3, 7)$, and $f(3) = 0$. Let $g(x) = \int_0^x f(t) dt$.

(a) On what interval is g increasing? [1]

(b) At what x does g attain a local maximum? Justify. [2]

3.3 For $g(x) = \int_0^x f(t) dt$, the graph of f is increasing on $(1, 5)$. State, with a reason, the concavity of g on $(1, 5)$. [2]