Part B (BC): Graphing calculator not allowed
Question 5 — 9 points

General Scoring Notes
The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.



The graphs of the functions f and g are shown in the figure for $0 \le x \le 3$. It is known that $g(x) = \frac{12}{3+x}$ for $x \ge 0$. The twice-differentiable function f, which is not explicitly given, satisfies $f(3) = 2$ and $\int_0^3 f(x)\,dx = 10$.

Model Solution	Scoring	

(a) Find the area of the shaded region enclosed by the graphs of f and g.

$\text{Area} = \int_0^3 (f(x) - g(x))\,dx = \int_0^3 f(x)\,dx - \int_0^3 g(x)\,dx$	Integrand	**1 point**
$= 10 - \int_0^3 \frac{12}{3+x}\,dx = 10 - 12\left[\ln\lvert 3+x\rvert\right]_0^3$	Antiderivative of $g(x)$	**1 point**
$= 10 - 12(\ln 6 - \ln 3) = 10 - 12(\ln 2)$	Answer	**1 point**

Scoring notes:

- The first point is earned for any of the integrands $f(x) - g(x)$, $g(x) - f(x)$, $\lvert f(x) - g(x)\rvert$, or $\lvert g(x) - f(x)\rvert$ in any definite integral. If the limits are incorrect, the response does not earn the third point.

- The first point is earned with an implied integrand for f and explicit integrand for g, such as $10 - \int_0^3 g(x)\,dx$.
- The second point is earned for finding $a\int \frac{dx}{3+x} = a \cdot \ln\lvert 3+x\rvert$ or $a \cdot \ln(3+x)$.
- A response is eligible for the third point only if it has earned the first 2 points. The third point is earned only for the correct answer. The answer does not need to be simplified; however, if simplification is attempted, it must be correct.
- A response is not eligible for the third point with incorrect limits of integration for u-substitution, for example, $\int_0^3 \frac{12}{3+x}\,dx = \int_0^3 \frac{12}{u}\,du = 12\left[\ln(x+3)\right]_0^3$.
- A response with incorrect communication, such as "Area $= \int_0^3 (g(x) - f(x))\,dx = 10 - 12(\ln 2)$," does not earn the third point. However, a response of "$\int_0^3 (g(x) - f(x))\,dx = 12(\ln 2) - 10$, so the area is $10 - 12(\ln 2)$" earns all 3 points.

Total for part (a) 3 points

(b) Evaluate the improper integral $\int_0^\infty (g(x))^2\,dx$, or show that the integral diverges.

$\int_0^\infty (g(x))^2\,dx = \lim_{b\to\infty} \int_0^b \frac{144}{(3+x)^2}\,dx$	Limit notation	**1 point**
$= \lim_{b\to\infty}\left(-\frac{144}{(3+x)}\Big\vert_0^b\right)$	Antiderivative	**1 point**
$= \lim_{b\to\infty}\left(-\frac{144}{3+b} + \frac{144}{3}\right) = 48$	Answer	**1 point**

Scoring notes:

- To earn the first point a response must correctly use limit notation throughout the problem and not include arithmetic with infinity, for example, $\left[-\frac{144}{3+x}\right]_0^\infty$ or $-\frac{144}{3+\infty} + 48$.
- The second point can be earned by finding an antiderivative of the form $-\frac{a}{(3+x)}$ for $a > 0$, from an indefinite or improper integral, with or without correct limit notation. If $a \ne 144$, the response does not earn the third point.
- The third point is earned only for an answer of 48 (or equivalent).
- A response is not eligible for the third point with incorrect limits of integration for u-substitution, for example, $\lim_{b\to\infty} \int_0^b \frac{144}{u^2}\,du = \lim_{b\to\infty}\left[-\frac{144}{3+x}\right]_0^b$.

Total for part (b) 3 points

(c) Let h be the function defined by $h(x) = x \cdot f'(x)$. Find the value of $\int_0^3 h(x)\,dx$.

Using integration by parts, $u = x \qquad dv = f'(x)\,dx$ $du = dx \qquad v = f(x)$	u and dv	**1 point**
$\int h(x)\,dx = \int x \cdot f'(x)\,dx = x \cdot f(x) - \int f(x)\,dx$	$\int h(x)\,dx$ $= x \cdot f(x) - \int f(x)\,dx$	**1 point**
$\int_0^3 h(x)\,dx = \int_0^3 x \cdot f'(x)\,dx = x \cdot f(x)\Big\rvert_0^3 - \int_0^3 f(x)\,dx$ $= (3 \cdot f(3) - 0 \cdot f(0)) - 10 = 3 \cdot 2 - 0 - 10 = -4$	Answer	**1 point**

Scoring notes:

- The first and second points are earned with an implied u and dv in the presence of $x \cdot f(x) - \int f(x)\,dx$ or $x \cdot f(x)\Big\rvert_0^3 - 10$.
- Limits of integration may be present, omitted, or partially present in the work for the first and second points.
- The tabular method may be used to show integration by parts. In this case, the first point is earned by having columns (labeled or unlabeled) that begin with x and $f'(x)$. The second point is earned for $x \cdot f(x) - \int f(x)\,dx$.
- The third point is earned only for the correct answer and can only be earned if the first 2 points were earned.

Total for part (c) 3 points

Total for question 5 9 points