

1. Male birds of a certain species arrive at a nesting area over a thirty-day period. The rate at which the male birds arrive at the nesting area at time t days is modeled by a differentiable function M , where $M(t)$ is measured in number of birds per day. Selected values of $M(t)$ are shown in the table.

t (days)	0	5	10	15	20	25	30
$M(t)$ (birds per day)	2	7	16	6	5	2	0

(Note: Your calculator should be in radian mode.)

Part A

Approximate $M'(7.5)$ using the average rate of change of M over the interval $5 \leq t \leq 10$. Show the work that leads to your answer, and indicate units of measure.

Part B

- i. Use a midpoint Riemann sum with the three subintervals $[0, 10]$, $[10, 20]$, and $[20, 30]$ to approximate $\int_0^{30} M(t) dt$. Show the work that leads to your answer.
- ii. Interpret the meaning of $\int_0^{30} M(t) dt$ in the context of the problem.

Part C

The rate at which female birds of the same species arrive at the same nesting area, in birds per day, is modeled by the function F defined as follows.

$$F(t) = \begin{cases} 0 & \text{for } 0 \leq t < 15 \\ 18 + 16 \sin\left(\frac{\pi}{20}(t + 15)\right) & \text{for } 15 \leq t \leq 45 \end{cases}$$

How many female birds of this species arrive at the nesting area from $t = 15$ to $t = 45$? Show the setup for your calculations, and round your answer to the nearest integer.

Part D

On the interval $15 < t < 30$, the difference in the rates at which male and female birds of this species arrive at the nesting area can be modeled by the differentiable function $D(t) = M(t) - F(t)$, where F is the function defined in part C. Is there a time t in the interval $15 < t < 20$ when $D(t) = 0$? Justify your answer.

3. A student starts reading a book at time $t = 0$ minutes and continues reading for the next 10 minutes. The rate at which the student reads is modeled by the differentiable function R , where $R(t)$ is measured in words per minute. Selected values of $R(t)$ are given in the table shown.

t (minutes)	0	2	8	10
$R(t)$ (words per minute)	90	100	150	162

- A. Approximate $R'(1)$ using the average rate of change of R over the interval $0 \leq t \leq 2$. Show the work that leads to your answer. Indicate units of measure.
- B. Must there be a value c , for $0 < c < 10$, such that $R(c) = 155$? Justify your answer.
- C. Use a trapezoidal sum with the three subintervals indicated by the data in the table to approximate the value of $\int_0^{10} R(t) dt$. Show the work that leads to your answer.
- D. A teacher also starts reading at time $t = 0$ minutes and continues reading for the next 10 minutes. The rate at which the teacher reads is modeled by the function W defined by $W(t) = -\frac{3}{10}t^2 + 8t + 100$, where $W(t)$ is measured in words per minute. Based on the model, how many words has the teacher read by the end of the 10 minutes? Show the work that leads to your answer.

t (days)	0	3	7	10	12
$r'(t)$ (centimeters per day)	-6.1	-5.0	-4.4	-3.8	-3.5

4. An ice sculpture melts in such a way that it can be modeled as a cone that maintains a conical shape as it decreases in size. The radius of the base of the cone is given by a twice-differentiable function r , where $r(t)$ is measured in centimeters and t is measured in days. The table above gives selected values of $r'(t)$, the rate of change of the radius, over the time interval $0 \leq t \leq 12$.

(a) Approximate $r''(8.5)$ using the average rate of change of r' over the interval $7 \leq t \leq 10$. Show the computations that lead to your answer, and indicate units of measure.

(b) Is there a time t , $0 \leq t \leq 3$, for which $r'(t) = -6$? Justify your answer.

(c) Use a right Riemann sum with the four subintervals indicated in the table to approximate the value of

$$\int_0^{12} r'(t) dt.$$

(d) The height of the cone decreases at a rate of 2 centimeters per day. At time $t = 3$ days, the radius is 100 centimeters and the height is 50 centimeters. Find the rate of change of the volume of the cone with respect to time, in cubic centimeters per day, at time $t = 3$ days. (The volume V of a cone with radius r and height h is $V = \frac{1}{3} \pi r^2 h$.)

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

CALCULUS BC
SECTION II, Part A
Time—30 minutes
Number of problems—2

A graphing calculator is required for these problems.

t (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

1. Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.

(a) Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.

(b) Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.

(c) Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.

(d) For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

t (minutes)	0	2	5	8	12
$v_A(t)$ (meters/minute)	0	100	40	-120	-150

4. Train A runs back and forth on an east-west section of railroad track. Train A 's velocity, measured in meters per minute, is given by a differentiable function $v_A(t)$, where time t is measured in minutes. Selected values for $v_A(t)$ are given in the table above.
- (a) Find the average acceleration of train A over the interval $2 \leq t \leq 8$.
- (b) Do the data in the table support the conclusion that train A 's velocity is -100 meters per minute at some time t with $5 < t < 8$? Give a reason for your answer.
- (c) At time $t = 2$, train A 's position is 300 meters east of the Origin Station, and the train is moving to the east. Write an expression involving an integral that gives the position of train A , in meters from the Origin Station, at time $t = 12$. Use a trapezoidal sum with three subintervals indicated by the table to approximate the position of the train at time $t = 12$.
- (d) A second train, train B , travels north from the Origin Station. At time t the velocity of train B is given by $v_B(t) = -5t^2 + 60t + 25$, and at time $t = 2$ the train is 400 meters north of the station. Find the rate, in meters per minute, at which the distance between train A and train B is changing at time $t = 2$.