

6. The Maclaurin series for a function g is given by

$$\sum_{n=0}^{\infty} 2 \left(\frac{-x}{5} \right)^n = 2 - \frac{2}{5}x + \frac{2}{25}x^2 - \frac{2}{125}x^3 + \frac{2}{625}x^4 - \dots + 2 \left(\frac{-x}{5} \right)^n + \dots \text{ and converges to } g(x)$$

for $|x| < 5$.

Part A

For $x > 0$, the Maclaurin series for g is an alternating geometric series. Find $g(3)$. Show the work that leads to your answer.

Part B

The function f is defined by $f(x) = g'(x)$. Write the first four nonzero terms of the Maclaurin series for f .

Part C

The second-degree Taylor polynomial for f about $x = 0$ is used to approximate $f\left(\frac{5}{2}\right)$ as $-\frac{3}{10}$, where f is the function defined in part B. Justify that $\left|f\left(\frac{5}{2}\right) - \left(-\frac{3}{10}\right)\right| \leq \frac{1}{5}$.

Part D

The function h is defined by $h(x) = 25g(x) - 2e^x$.

- i. Write the first three nonzero terms of the Maclaurin series for e^x .
- ii. Write the first three nonzero terms of the Maclaurin series for h .

STOP
END OF EXAM

6. The Maclaurin series for a function f is given by $\sum_{n=1}^{\infty} \frac{(n+1)x^n}{n^2 6^n}$ and converges to $f(x)$ for all x in the interval

of convergence. It can be shown that the Maclaurin series for f has a radius of convergence of 6.

(a) Determine whether the Maclaurin series for f converges or diverges at $x = 6$. Give a reason for your answer.

(b) It can be shown that $f(-3) = \sum_{n=1}^{\infty} \frac{(n+1)(-3)^n}{n^2 6^n} = \sum_{n=1}^{\infty} \frac{n+1}{n^2} \left(-\frac{1}{2}\right)^n$ and that the first three terms of this

series sum to $S_3 = -\frac{125}{144}$. Show that $\left|f(-3) - S_3\right| < \frac{1}{50}$.

(c) Find the general term of the Maclaurin series for f' , the derivative of f . Find the radius of convergence of the Maclaurin series for f' .

(d) Let $g(x) = \sum_{n=1}^{\infty} \frac{(n+1)x^{2n}}{n^2 3^n}$. Use the ratio test to determine the radius of convergence of the Maclaurin series

for g .

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

STOP

END OF EXAM

6. The function f is defined by the power series $f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots + \frac{(-1)^n x^{2n+1}}{2n+1} + \cdots$ for all real numbers x for which the series converges.

(a) Using the ratio test, find the interval of convergence of the power series for f . Justify your answer.

(b) Show that $\left| f\left(\frac{1}{2}\right) - \frac{1}{2} \right| < \frac{1}{10}$. Justify your answer.

(c) Write the first four nonzero terms and the general term for an infinite series that represents $f'(x)$.

(d) Use the result from part (c) to find the value of $f'\left(\frac{1}{6}\right)$.

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

STOP

END OF EXAM

6. The function g has derivatives of all orders for all real numbers. The Maclaurin series for g is given by

$$g(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^n}{2e^n + 3} \text{ on its interval of convergence.}$$

(a) State the conditions necessary to use the integral test to determine convergence of the series $\sum_{n=0}^{\infty} \frac{1}{e^n}$.

Use the integral test to show that $\sum_{n=0}^{\infty} \frac{1}{e^n}$ converges.

(b) Use the limit comparison test with the series $\sum_{n=0}^{\infty} \frac{1}{e^n}$ to show that the series $g(1) = \sum_{n=0}^{\infty} \frac{(-1)^n}{2e^n + 3}$

converges absolutely.

(c) Determine the radius of convergence of the Maclaurin series for g .


(d) The first two terms of the series $g(1) = \sum_{n=0}^{\infty} \frac{(-1)^n}{2e^n + 3}$ are used to approximate $g(1)$. Use the alternating

series error bound to determine an upper bound on the error of the approximation.

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

STOP

END OF EXAM



n	$f^{(n)}(0)$
2	3
3	$-\frac{23}{2}$
4	54

6. A function f has derivatives of all orders for all real numbers x . A portion of the graph of f is shown above, along with the line tangent to the graph of f at $x = 0$. Selected derivatives of f at $x = 0$ are given in the table above.

- Write the third-degree Taylor polynomial for f about $x = 0$.
- Write the first three nonzero terms of the Maclaurin series for e^x . Write the second-degree Taylor polynomial for $e^x f(x)$ about $x = 0$.
- Let h be the function defined by $h(x) = \int_0^x f(t) dt$. Use the Taylor polynomial found in part (a) to find an approximation for $h(1)$.
- It is known that the Maclaurin series for h converges to $h(x)$ for all real numbers x . It is also known that the individual terms of the series for $h(1)$ alternate in sign and decrease in absolute value to 0. Use the alternating series error bound to show that the approximation found in part (c) differs from $h(1)$ by at most 0.45.

STOP
END OF EXAM

6. The Maclaurin series for $\ln(1+x)$ is given by

$$x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots + (-1)^{n+1} \frac{x^n}{n} + \cdots$$

On its interval of convergence, this series converges to $\ln(1+x)$. Let f be the function defined by

$$f(x) = x \ln\left(1 + \frac{x}{3}\right).$$

- Write the first four nonzero terms and the general term of the Maclaurin series for f .
- Determine the interval of convergence of the Maclaurin series for f . Show the work that leads to your answer.
- Let $P_4(x)$ be the fourth-degree Taylor polynomial for f about $x = 0$. Use the alternating series error bound to find an upper bound for $|P_4(2) - f(2)|$.

STOP
END OF EXAM

$$f(0) = 0$$

$$f'(0) = 1$$

$$f^{(n+1)}(0) = -n \cdot f^{(n)}(0) \text{ for all } n \geq 1$$

6. A function f has derivatives of all orders for $-1 < x < 1$. The derivatives of f satisfy the conditions above.

The Maclaurin series for f converges to $f(x)$ for $|x| < 1$.

- (a) Show that the first four nonzero terms of the Maclaurin series for f are $x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4}$, and write the

general term of the Maclaurin series for f .

- (b) Determine whether the Maclaurin series described in part (a) converges absolutely, converges conditionally, or diverges at $x = 1$. Explain your reasoning.

- (c) Write the first four nonzero terms and the general term of the Maclaurin series for $g(x) = \int_0^x f(t) dt$.

- (d) Let $P_n\left(\frac{1}{2}\right)$ represent the n th-degree Taylor polynomial for g about $x = 0$ evaluated at $x = \frac{1}{2}$, where g is

the function defined in part (c). Use the alternating series error bound to show that

$$\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right| < \frac{1}{500}.$$

STOP
END OF EXAM

6. The function f has a Taylor series about $x = 1$ that converges to $f(x)$ for all x in the interval of convergence.

It is known that $f(1) = 1$, $f'(1) = -\frac{1}{2}$, and the n th derivative of f at $x = 1$ is given by $f^{(n)}(1) = (-1)^n \frac{(n-1)!}{2^n}$

for $n \geq 2$.

- (a) Write the first four nonzero terms and the general term of the Taylor series for f about $x = 1$.
- (b) The Taylor series for f about $x = 1$ has a radius of convergence of 2. Find the interval of convergence. Show the work that leads to your answer.
- (c) The Taylor series for f about $x = 1$ can be used to represent $f(1.2)$ as an alternating series. Use the first three nonzero terms of the alternating series to approximate $f(1.2)$.
- (d) Show that the approximation found in part (c) is within 0.001 of the exact value of $f(1.2)$.

STOP
END OF EXAM