

Part B (BC): Graphing calculator not allowed**Question 5****9 points****General Scoring Notes**

- The model solution is presented using standard mathematical notation.
- Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be accurate to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Let $y = f(x)$ be the particular solution to the differential equation $\frac{dy}{dx} = (3 - x)y^2$ with initial condition $f(1) = -1$.

	Model Solution	Scoring
A	Find $f''(1)$, the value of $\frac{d^2y}{dx^2}$ at the point $(1, -1)$. Show the work that leads to your answer.	
	$\frac{d^2y}{dx^2} = -y^2 + (3 - x)2y \frac{dy}{dx}$	Product rule Point 1 (P1)
		Chain rule Point 2 (P2)
	$f'(1) = \frac{dy}{dx} \Big _{(x,y)=(1,-1)} = (3 - 1)(-1)^2 = 2$	$f''(1)$ Point 3 (P3)
	$f''(1) = \frac{d^2y}{dx^2} \Big _{(x,y)=(1,-1)} = -(-1)^2 + (3 - 1)(2)(-1)(2) = -9$	

Scoring Notes for Part A

- The expression $\frac{d^2y}{dx^2} = -y^2 + (3 - x)2y$ or $\frac{d^2y}{dx^2} = 6y - y^2 - 2xy$ earns **P1** but not **P2**. Such a response is not eligible for **P3**.
- The expression $\frac{d^2y}{dx^2} = -2y \frac{dy}{dx}$ or $\frac{d^2y}{dx^2} = 6y \left(\frac{dy}{dx} \right) - 2y \left(\frac{dy}{dx} \right)$ earns **P2** but not **P1**. Such a response is eligible for **P3** for a consistent answer of $f''(1) = 4$ or $f''(1) = -8$, respectively, which is found by correctly substituting correct values for x , y , and $\frac{dy}{dx}$.
- A response of $-(-1)^2 + (3 - 1)(2)(-1)(2)$ earns **P1**, **P2**, and **P3** regardless of any subsequent errors in simplification.
- Alternate approach (using separation of variables):

The particular solution for the differential equation that passes through the point $(1, -1)$ is

$$y = \frac{2}{x^2 - 6x + 3}. \text{ Therefore, } \frac{dy}{dx} = \frac{-2(2x - 6)}{(x^2 - 6x + 3)^2}.$$

This response has not yet earned **P1**, **P2**, or **P3**.

- A response that correctly applies the quotient rule (or product rule) and the chain rule to find that $\frac{d^2y}{dx^2} = \frac{-4(x^2 - 6x + 3)^2 + 4(2x - 6)(x^2 - 6x + 3)(2x - 6)}{(x^2 - 6x + 3)^4}$ earns **P1** and **P2** and is eligible to earn **P3** for the correct answer of $f''(1) = -9$.
- A response that correctly applies the quotient rule (or product rule) but does not correctly apply the chain rule (e.g., $\frac{d^2y}{dx^2} = \frac{-4(x^2 - 6x + 3)^2 + 4(2x - 6)(x^2 - 6x + 3)}{(x^2 - 6x + 3)^4}$) earns **P1**, does not earn **P2**, and is not eligible to earn **P3**.
- A response that does not correctly apply the quotient rule (or product rule) but does correctly apply the chain rule (e.g., $\frac{d^2y}{dx^2} = \frac{-4}{2(x^2 - 6x + 3)(2x - 6)}$) does not earn **P1**, earns **P2**, and is eligible to earn **P3** for a consistent answer.

B Write the second-degree Taylor polynomial for f about $x = 1$.

$$f'(1) = 2 \text{ and } f''(1) = -9$$

Two terms

Point 4 (P4)

$$P_2(x) = -1 + 2(x - 1) - \frac{9}{2}(x - 1)^2$$

Remaining term

Point 5 (P5)

Scoring Notes for Part B

- **P4** and **P5** can be earned with an answer consistent with incorrect values of $f'(1)$ and $f''(1)$ imported from part A.
- Any terms of degree greater than two or “+...” does not earn **P5**.
- A response of $-1 + 2(x - 1) - \frac{9}{2}(x - 1)^2$ earns **P4** and **P5**, regardless of any subsequent algebraic simplification.
- A response that does not present the polynomial as powers of $(x - 1)$ but instead presents a correct expanded/simplified form of the polynomial (e.g., $-\frac{9}{2}x^2 + 11x - \frac{15}{2}$) earns **P4** but not **P5**.

C The second-degree Taylor polynomial for f about $x = 1$ is used to approximate $f(1.1)$. Given that $|f'''(x)| \leq 60$ for all x in the interval $1 \leq x \leq 1.1$, use the Lagrange error bound to show that this approximation differs from $f(1.1)$ by at most 0.01.

$$|f(1.1) - P_2(1.1)| \leq \frac{\max_{1 \leq x \leq 1.1} |f'''(x)|}{3!} |1.1 - 1|^3 \leq \frac{60}{6} (0.1)^3 = 0.01$$

Form of error bound

Point 6 (P6)

Analysis

Point 7 (P7)

Scoring Notes for Part C

- **P6** is earned for presenting either $\frac{\max_{1 \leq x \leq 1.1} |f'''(x)|}{3!} |1.1 - 1|^3$ or $\frac{60}{6} (0.1)^3$. Subsequent errors in simplification will not earn **P7**.
- To earn **P7**, a response must have earned **P6** and must explicitly connect the error bound with 0.01; for example by communicating Error ≤ 0.01 , Error Bound = 0.01, or equivalent.
- A response that declares the error is equal to 0.01 (or any equivalent form of this value) does not earn **P7**.

D Use Euler's method, starting at $x = 1$ with two steps of equal size, to approximate $f(1.4)$. Show the work that leads to your answer.

$$f(1.2) \approx f(1) + (1.2 - 1) \cdot \frac{dy}{dx} \Big|_{(1, -1)}$$

$$= -1 + 0.2(2) = -0.6$$

First step of Euler's method

Point 8 (P8)

$$f(1.4) \approx f(1.2) + (1.4 - 1.2) \cdot \frac{dy}{dx} \Big|_{(1.2, -0.6)}$$

$$\approx -0.6 + 0.2(3 - 1.2)(-0.6)^2 = -0.4704$$

Answer with supporting work

Point 9 (P9)

An approximation of $f(1.4)$ is -0.47 .

Scoring Notes for Part D

- To earn **P8**, a response must demonstrate the first step of Euler's method, with the correct initial condition, correct step size, and correct (or imported) expression for the derivative.
Note: Any subsequent error in simplification or rounding will not affect the scoring for **P8**.
- The two steps of Euler's method may be explicit expressions or may be presented in a table. For example:

x	y	$\frac{dy}{dx} \cdot \Delta x$ (or $\frac{dy}{dx} \cdot 0.2$)
1	-1	0.4
1.2	-0.6	0.1296
1.4	-0.4704	

Note: In the presence of a correct answer, a table does not need to be labeled to earn both **P8** and **P9**. In the presence of no answer or an incorrect answer, such a table must be correctly labeled to earn **P8**.

- A response of $-0.6 + 0.2(3 - 1.2)(-0.6)^2$ earns **P9**, regardless of any subsequent errors in simplification or rounding.
- A response that imports an incorrect value for $f'(1)$ from part A or part B is eligible to earn **P9** with a consistent answer.
- A response may report the final answer as -0.47 , $(1.4, -0.47)$, $-\frac{294}{625}$, or equivalent.

Part B (BC): Graphing calculator not allowed**Question 6****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

The function f has derivatives of all orders for all real numbers. It is known that $f(0) = 2$, $f'(0) = 3$, $f''(x) = -f(x^2)$, and $f'''(x) = -2x \cdot f'(x^2)$.

Model Solution	Scoring
(a) Find $f^{(4)}(x)$, the fourth derivative of f with respect to x . Write the fourth-degree Taylor polynomial for f about $x = 0$. Show the work that leads to your answer.	
$f^{(4)}(x) = -2 \cdot f'(x^2) + (-2x)f''(x^2) \cdot 2x$	Form of product rule 1 point
	$f^{(4)}(x)$ 1 point
$f'''(0) = -f(0) = -2$ $f'''(0) = -2(0) \cdot f'(0) = 0$ $f^{(4)}(0) = -2 \cdot f'(0) + 0 \cdot f''(0) \cdot 0 = -2 \cdot 3 + 0 = -6$	Two terms of polynomial 1 point
	Remaining terms 1 point
The fourth-degree Taylor polynomial for f about $x = 0$ is $T_4(x) = 2 + 3x + \frac{-2}{2!}x^2 + \frac{0}{3!}x^3 + \frac{-6}{4!}x^4$ $= 2 + 3x - x^2 - \frac{1}{4}x^4$	
Scoring notes:	
- The first point is earned for a correct fourth derivative or for $f^{(4)}(x) = -2 \cdot f'(x^2) + (-2x)f''(x^2)$. - The second point is earned only for a completely correct expression for $f^{(4)}(x)$. - A response that earns the first point but not the second may evaluate the presented expression for $f^{(4)}(x)$ at $x = 0$ and use the consistent nonzero value in computing the coefficient of x^4 in the fourth-degree Taylor polynomial. - A polynomial that includes a nonzero third-degree term, any terms of degree greater than four, or $+\dots$ does not earn the fourth point.	
Total for part (a)	4 points

- (b) The fourth-degree Taylor polynomial for f about $x = 0$ is used to approximate $f(0.1)$. Given that $|f^{(5)}(x)| \leq 15$ for $0 \leq x \leq 0.5$, use the Lagrange error bound to show that this approximation is within $\frac{1}{10^5}$ of the exact value of $f(0.1)$.

By the Lagrange error bound,

$$|T_4(0.1) - f(0.1)| \leq \frac{\max_{0 \leq x \leq 0.1} |f^{(5)}(x)|}{5!} \cdot (0.1)^5$$

$$\leq \frac{15}{120} \cdot \frac{1}{10^5} \leq \frac{1}{10^5}$$

Form of error bound **1 point**Shows $|\text{Error}| \leq \frac{1}{10^5}$ **1 point****Scoring notes:**

- The first point is earned for $\frac{\max_{0 \leq x \leq 0.1} |f^{(5)}(x)|}{5!} \cdot (0.1)^5$ or $\frac{15}{5!}(0.1)^5$. Subsequent errors in simplification will not earn the second point.
- To earn the second point a response must communicate the inequality $\text{Error} \leq \frac{15}{5!} \cdot (0.1)^5 \leq \frac{1}{10^5}$.
- A response that states $\text{Error} = \frac{15}{5!} \cdot (0.1)^5$ or $\text{Error} = \frac{1}{10^5}$ does not earn the second point.

Total for part (b) **2 points**

- (c) Let g be the function such that $g(0) = 4$ and $g'(x) = e^x f(x)$. Write the second-degree Taylor polynomial for g about $x = 0$.

$g''(x) = e^x \cdot f(x) + e^x \cdot f'(x)$	$g''(x)$	1 point
$g'(0) = e^0 \cdot f(0) = 2$ $g''(0) = e^0 \cdot f(0) + e^0 \cdot f'(0) = 2 + 3 = 5$	First two terms of polynomial	1 point
	Taylor polynomial	1 point
The second-degree Taylor polynomial for g about $x = 0$ is $T_2(x) = 4 + 2x + \frac{5}{2}x^2$.		

Scoring notes:

- The first point is earned for $g''(x) = e^x \cdot f(x) + e^x \cdot f'(x)$, $g''(0) = e^0 \cdot f(0) + e^0 \cdot f'(0)$, or $g''(0) = f(0) + f'(0)$.
- A presented polynomial of the form $4 + 2x + ax^2$ earns the second point with or without any supporting work for the first two terms.
- A response that earned neither the first nor the second point only earns the third point for a polynomial of the form $a + bx + \frac{c}{2}x^2$, where $c \neq 0$ is declared to be $g''(0)$.

- A polynomial that includes any terms of degree greater than two, or $+\dots$, does not earn the third point.
- Alternate solution:

$$e^x = 1 + x + \frac{x^2}{2} + \dots$$

$$e^x f(x) = \left(1 + x + \frac{x^2}{2} + \dots\right)(2 + 3x - x^2 + \dots) = 2 + 5x + \dots$$

$$g(x) = \int e^x f(x) dx = C + 2x + \frac{5}{2}x^2 + \dots$$

$$g(0) = 4 \Rightarrow C = 4$$

$$g(x) \approx 4 + 2x + \frac{5}{2}x^2$$

- A response that is using this alternate solution method earns the first point for $e^x f(x) = 2 + 5x + \dots$, the second point for any two correct terms in a second-degree polynomial, and the third point for a completely correct second-degree Taylor polynomial with supporting work.
- Note: There is not enough information to conclude that $f(x)$ is equal to its Maclaurin series on its interval of convergence. The second and third lines of the alternate solution are being accepted as identifications of the Maclaurin series for $e^x f(x)$ and $g(x)$, respectively.

Total for part (c) 3 points

Total for question 6 9 points

Part B (BC): Graphing calculator not allowed

Question 5

9 points

General Scoring Notes

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Scoring guidelines and notes contain examples of the most common approaches seen in student responses. These guidelines can be applied to alternate approaches to ensure that these alternate approaches are scored appropriately.

Let $y = f(x)$ be the particular solution to the differential equation $\frac{dy}{dx} = y \cdot (x \ln x)$ with initial condition $f(1) = 4$. It can be shown that $f''(1) = 4$.

Model Solution	Scoring	
(a) Write the second-degree Taylor polynomial for f about $x = 1$. Use the Taylor polynomial to approximate $f(2)$.		
$f'(1) = \frac{dy}{dx} \Big _{(x,y)=(1,4)} = 4 \cdot (1 \ln 1) = 0$	Polynomial	1 point
The second-degree Taylor polynomial for f about $x = 1$ is $f(1) + f'(1)(x-1) + \frac{f''(1)}{2!}(x-1)^2 = 4 + 0(x-1) + \frac{4}{2}(x-1)^2.$		
$f(2) \approx 4 + 2(2-1)^2 = 6$	Approximation	1 point

Scoring notes:

- The first point is earned for $4 + \frac{4 \cdot \ln 1}{1!}(x-1)^1 + \frac{4}{2!}(x-1)^2$ or any correctly simplified equivalent expression. A term involving $(x-1)$ is not necessary. The polynomial must be written about (centered at) $x = 1$.
- If the first point is earned, the second point is earned for just “6” with no additional supporting work.
- If the polynomial is never explicitly written, the first point is not earned. In this case, to earn the second point supporting work of at least “ $4 + 2(1)$ ” is required.

Total for part (a) 2 points

- (b) Use Euler's method, starting at $x = 1$ with two steps of equal size, to approximate $f(2)$. Show the work that leads to your answer.

$f(1.5) \approx f(1) + 0.5 \cdot \frac{dy}{dx} \Big _{(x,y)=(1,4)} = 4 + 0.5 \cdot 0 = 4$	Euler's method with two steps	1 point
$f(2) \approx f(1.5) + 0.5 \cdot \frac{dy}{dx} \Big _{(x,y)=(1.5,4)}$		
$\approx 4 + 0.5 \cdot 4 \cdot (1.5 \ln 1.5) = 4 + 3 \ln 1.5$	Answer	1 point

Scoring notes:

- The first point is earned for two steps (of size 0.5) of Euler's method, with at most one error. If there is any error, the second point is not earned.
- To earn the first point a response must contain two Euler steps, $\Delta x = 0.5$, use of the correct expression for $\frac{dy}{dx}$, and use of the initial condition $f(1) = 4$.
 - The two Euler steps may be explicit expressions or may be presented in a table. Here is a minimal example of a (correctly labeled) table.

x	y	$\Delta y = \frac{dy}{dx} \cdot \Delta x$ or $\Delta y = \frac{dy}{dx} \cdot (0.5)$
1	4	0
1.5	4	$3 \ln 1.5$
2	$4 + 3 \ln 1.5$	

- Note: In the presence of the correct answer, such a table does not need to be labeled in order to earn both points. In the presence of an incorrect answer, the table must be correctly labeled for the response to earn the first point.
- A single error in computing the approximation of $f(1.5)$ is not considered a second error if that incorrect value is imported correctly into an approximation of $f(2)$.
- Both points are earned for " $4 + 0.5 \cdot 0 + 0.5 \cdot 4 \cdot (1.5 \ln 1.5)$ " or " $4 + 0.5 \cdot 4 \cdot (1.5 \ln 1.5)$ ".
- Both points are earned for presenting the ordered pair $(2, 4 + 3 \ln 1.5)$ with sufficient supporting work.

Total for part (b) 2 points

- (c) Find the particular solution $y = f(x)$ to the differential equation $\frac{dy}{dx} = y \cdot (x \ln x)$ with initial condition $f(1) = 4$.

$\frac{1}{y} dy = x \ln x dx$	Separation of variables	1 point
Using integration by parts, $u = \ln x \quad du = \frac{1}{x} dx$ $dv = x dx \quad v = \frac{x^2}{2}$	Antiderivative for $x \ln x$	1 point
$\int x \ln x dx = \frac{x^2}{2} \cdot \ln x - \int \frac{x^2}{2} \cdot \frac{1}{x} dx = \frac{x^2 \ln x}{2} - \frac{x^2}{4} + C$		
$\ln y = \frac{x^2 \ln x}{2} - \frac{x^2}{4} + C$	Antiderivative for $\frac{1}{y}$	1 point
$\ln 4 = 0 - \frac{1}{4} + C \Rightarrow C = \ln 4 + \frac{1}{4}$	Constant of integration and uses initial condition	1 point
$y = e^{\left(\frac{x^2 \ln x}{2} - \frac{x^2}{4} + \ln 4 + \frac{1}{4}\right)}$	Solves for y	1 point

Note: This solution is valid for $x > 0$.

Scoring notes:

- A response with no separation of variables earns 0 out of 5 points. If an error in separation results in one side being correct ($\frac{1}{y} dy$ or $x \ln x dx$), the response is only eligible to earn the corresponding antiderivative point.
- The third point (antiderivative of $\frac{1}{y}$) can be earned for either $\ln y$ or $\ln|y|$.
- A response with no constant of integration can earn at most 3 out of 5 points.
- A response is eligible for the fourth point if it has earned the first point and at least 1 of the 2 antiderivative points.
- A response earns the fourth point by correctly including the constant of integration in an equation and then replacing x with 1 and y with 4.
- A response is eligible for the fifth point only if it has earned the first 4 points.

Total for part (c) 5 points

Total for question 5 9 points

AP[®] Calculus BC

Scoring Guidelines

2019 SCORING GUIDELINES

Question 1

(a) $\int_0^5 E(t) dt = 153.457690$

To the nearest whole number, 153 fish enter the lake from midnight to 5 A.M.

(b) $\frac{1}{5-0} \int_0^5 L(t) dt = 6.059038$

The average number of fish that leave the lake per hour from midnight to 5 A.M. is 6.059 fish per hour.

- (c) The rate of change in the number of fish in the lake at time t is given by $E(t) - L(t)$.

$$E(t) - L(t) = 0 \Rightarrow t = 6.20356$$

$E(t) - L(t) > 0$ for $0 \leq t < 6.20356$, and $E(t) - L(t) < 0$ for $6.20356 < t \leq 8$. Therefore the greatest number of fish in the lake is at time $t = 6.204$ (or 6.203).

— OR —

Let $A(t)$ be the change in the number of fish in the lake from midnight to t hours after midnight.

$$A(t) = \int_0^t (E(s) - L(s)) ds$$

$$A'(t) = E(t) - L(t) = 0 \Rightarrow t = C = 6.20356$$

t	$A(t)$
0	0
C	135.01492
8	80.91998

Therefore the greatest number of fish in the lake is at time $t = 6.204$ (or 6.203).

(d) $E'(5) - L'(5) = -10.7228 < 0$

Because $E'(5) - L'(5) < 0$, the rate of change in the number of fish is decreasing at time $t = 5$.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

3 : $\begin{cases} 1 : \text{sets } E(t) - L(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

2 : $\begin{cases} 1 : \text{considers } E'(5) \text{ and } L'(5) \\ 1 : \text{answer with explanation} \end{cases}$

2019 SCORING GUIDELINES

Question 2

(a) $\frac{1}{2} \int_0^{\sqrt{\pi}} (r(\theta))^2 d\theta = 3.534292$

The area of S is 3.534.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) $\frac{1}{\sqrt{\pi}-0} \int_0^{\sqrt{\pi}} r(\theta) d\theta = 1.579933$

The average distance from the origin to a point on the curve $r = r(\theta)$ for $0 \leq \theta \leq \sqrt{\pi}$ is 1.580 (or 1.579).

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(c) $\tan \theta = \frac{y}{x} = m \Rightarrow \theta = \tan^{-1} m$

$$\frac{1}{2} \int_0^{\tan^{-1} m} (r(\theta))^2 d\theta = \frac{1}{2} \left(\frac{1}{2} \int_0^{\sqrt{\pi}} (r(\theta))^2 d\theta \right)$$

3 : $\begin{cases} 1 : \text{equates polar areas} \\ 1 : \text{inverse trigonometric function} \\ \text{applied to } m \text{ as limit of} \\ \text{integration} \\ 1 : \text{equation} \end{cases}$

(d) As $k \rightarrow \infty$, the circle $r = k \cos \theta$ grows to enclose all points to the right of the y -axis.

$$\begin{aligned} \lim_{k \rightarrow \infty} A(k) &= \frac{1}{2} \int_0^{\pi/2} (r(\theta))^2 d\theta \\ &= \frac{1}{2} \int_0^{\pi/2} (3\sqrt{\theta} \sin(\theta^2))^2 d\theta = 3.324 \end{aligned}$$

2 : $\begin{cases} 1 : \text{limits of integration} \\ 1 : \text{answer with integral} \end{cases}$

2019 SCORING GUIDELINES

Question 3

(a) $\int_{-6}^5 f(x) dx = \int_{-6}^{-2} f(x) dx + \int_{-2}^5 f(x) dx$

$$\Rightarrow 7 = \int_{-6}^{-2} f(x) dx + 2 + \left(9 - \frac{9\pi}{4}\right)$$

$$\Rightarrow \int_{-6}^{-2} f(x) dx = 7 - \left(11 - \frac{9\pi}{4}\right) = \frac{9\pi}{4} - 4$$

3 : $\begin{cases} 1 : \int_{-6}^5 f(x) dx = \int_{-6}^{-2} f(x) dx + \int_{-2}^5 f(x) dx \\ 1 : \int_{-2}^5 f(x) dx \\ 1 : \text{answer} \end{cases}$

(b) $\begin{aligned} \int_3^5 (2f'(x) + 4) dx &= 2 \int_3^5 f'(x) dx + \int_3^5 4 dx \\ &= 2(f(5) - f(3)) + 4(5 - 3) \\ &= 2(0 - (3 - \sqrt{5})) + 8 \\ &= 2(-3 + \sqrt{5}) + 8 = 2 + 2\sqrt{5} \end{aligned}$

2 : $\begin{cases} 1 : \text{Fundamental Theorem of Calculus} \\ 1 : \text{answer} \end{cases}$

— OR —

$$\begin{aligned} \int_3^5 (2f'(x) + 4) dx &= [2f(x) + 4x]_{x=3}^{x=5} \\ &= (2f(5) + 20) - (2f(3) + 12) \\ &= (2 \cdot 0 + 20) - (2(3 - \sqrt{5}) + 12) \\ &= 2 + 2\sqrt{5} \end{aligned}$$

(c) $g'(x) = f(x) = 0 \Rightarrow x = -1, x = \frac{1}{2}, x = 5$

x	$g(x)$
-2	0
-1	$\frac{1}{2}$
$\frac{1}{2}$	$-\frac{1}{4}$
5	$11 - \frac{9\pi}{4}$

3 : $\begin{cases} 1 : g'(x) = f(x) \\ 1 : \text{identifies } x = -1 \text{ as a candidate} \\ 1 : \text{answer with justification} \end{cases}$

On the interval $-2 \leq x \leq 5$, the absolute maximum value of g is $g(5) = 11 - \frac{9\pi}{4}$.

(d) $\begin{aligned} \lim_{x \rightarrow 1} \frac{10^x - 3f'(x)}{f(x) - \arctan x} &= \frac{10^1 - 3f'(1)}{f(1) - \arctan 1} \\ &= \frac{10 - 3 \cdot 2}{1 - \arctan 1} = \frac{4}{1 - \frac{\pi}{4}} \end{aligned}$

1 : answer

2019 SCORING GUIDELINES

Question 4

(a) $V = \pi r^2 h = \pi(1)^2 h = \pi h$

$$\left. \frac{dV}{dt} \right|_{h=4} = \pi \left. \frac{dh}{dt} \right|_{h=4} = \pi \left(-\frac{1}{10} \sqrt{4} \right) = -\frac{\pi}{5} \text{ cubic feet per second}$$

2 : $\begin{cases} 1 : \frac{dV}{dt} = \pi \frac{dh}{dt} \\ 1 : \text{answer with units} \end{cases}$

(b) $\frac{d^2 h}{dt^2} = -\frac{1}{20\sqrt{h}} \cdot \frac{dh}{dt} = -\frac{1}{20\sqrt{h}} \cdot \left(-\frac{1}{10} \sqrt{h} \right) = \frac{1}{200}$

Because $\frac{d^2 h}{dt^2} = \frac{1}{200} > 0$ for $h > 0$, the rate of change of the height is increasing when the height of the water is 3 feet.

3 : $\begin{cases} 1 : \frac{d}{dh} \left(-\frac{1}{10} \sqrt{h} \right) = -\frac{1}{20\sqrt{h}} \\ 1 : \frac{d^2 h}{dt^2} = -\frac{1}{20\sqrt{h}} \cdot \frac{dh}{dt} \\ 1 : \text{answer with explanation} \end{cases}$

(c) $\frac{dh}{\sqrt{h}} = -\frac{1}{10} dt$

$$\int \frac{dh}{\sqrt{h}} = \int -\frac{1}{10} dt$$

$$2\sqrt{h} = -\frac{1}{10}t + C$$

$$2\sqrt{5} = -\frac{1}{10} \cdot 0 + C \Rightarrow C = 2\sqrt{5}$$

$$2\sqrt{h} = -\frac{1}{10}t + 2\sqrt{5}$$

$$h(t) = \left(-\frac{1}{20}t + \sqrt{5} \right)^2$$

4 : $\begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ \text{and uses initial condition} \\ 1 : h(t) \end{cases}$

Note: 0/4 if no separation of variables

Note: max 2/4 [1-1-0-0] if no constant of integration

2019 SCORING GUIDELINES

Question 5

(a) $f'(x) = \frac{-(2x-2)}{(x^2-2x+k)^2}$

$$f'(0) = \frac{2}{k^2} = 6 \Rightarrow k^2 = \frac{1}{3} \Rightarrow k = \frac{1}{\sqrt{3}}$$

3 : $\begin{cases} 1 : \text{denominator of } f'(x) \\ 1 : f'(x) \\ 1 : \text{answer} \end{cases}$

(b) $\frac{1}{x^2-2x-8} = \frac{1}{(x-4)(x+2)} = \frac{A}{x-4} + \frac{B}{x+2}$

$$\Rightarrow 1 = A(x+2) + B(x-4)$$

$$\Rightarrow A = \frac{1}{6}, B = -\frac{1}{6}$$

3 : $\begin{cases} 1 : \text{partial fraction} \\ \text{decomposition} \\ 1 : \text{antiderivatives} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \int_0^1 f(x) dx &= \int_0^1 \left(\frac{1}{6} \frac{1}{x-4} - \frac{1}{6} \frac{1}{x+2} \right) dx \\ &= \left[\frac{1}{6} \ln|x-4| - \frac{1}{6} \ln|x+2| \right]_{x=0}^{x=1} \\ &= \left(\frac{1}{6} \ln 3 - \frac{1}{6} \ln 3 \right) - \left(\frac{1}{6} \ln 4 - \frac{1}{6} \ln 2 \right) = -\frac{1}{6} \ln 2 \end{aligned}$$

(c) $\int_0^2 \frac{1}{x^2-2x+1} dx = \int_0^2 \frac{1}{(x-1)^2} dx = \int_0^1 \frac{1}{(x-1)^2} dx + \int_1^2 \frac{1}{(x-1)^2} dx$

$$= \lim_{b \rightarrow 1^-} \int_0^b \frac{1}{(x-1)^2} dx + \lim_{b \rightarrow 1^+} \int_b^2 \frac{1}{(x-1)^2} dx$$

$$= \lim_{b \rightarrow 1^-} \left(-\frac{1}{x-1} \Big|_{x=0}^{x=b} \right) + \lim_{b \rightarrow 1^+} \left(-\frac{1}{x-1} \Big|_{x=b}^{x=2} \right)$$

$$= \lim_{b \rightarrow 1^-} \left(-\frac{1}{b-1} - 1 \right) + \lim_{b \rightarrow 1^+} \left(-1 + \frac{1}{b-1} \right)$$

Because $\lim_{b \rightarrow 1^-} \left(-\frac{1}{b-1} \right)$ does not exist, the integral diverges.

3 : $\begin{cases} 1 : \text{improper integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer with reason} \end{cases}$

2019 SCORING GUIDELINES

Question 6

- (a)
- $f(0) = 3$
- and
- $f'(0) = -2$

The third-degree Taylor polynomial for f about $x = 0$ is

$$3 - 2x + \frac{3}{2!}x^2 + \frac{-2}{3!}x^3 = 3 - 2x + \frac{3}{2}x^2 - \frac{2}{12}x^3.$$

- (b) The first three nonzero terms of the Maclaurin series for
- e^x
- are

$$1 + x + \frac{1}{2!}x^2.$$

The second-degree Taylor polynomial for $e^x f(x)$ about $x = 0$ is

$$\begin{aligned} 3\left(1 + x + \frac{1}{2!}x^2\right) - 2x(1 + x) + \frac{3}{2}x^2(1) \\ = 3 + (3 - 2)x + \left(\frac{3}{2} - 2 + \frac{3}{2}\right)x^2 \\ = 3 + x + x^2. \end{aligned}$$

- (c)
- $h(1) = \int_0^1 f(t) dt$

$$\begin{aligned} &\approx \int_0^1 \left(3 - 2t + \frac{3}{2}t^2 - \frac{23}{12}t^3\right) dt \\ &= \left[3t - t^2 + \frac{1}{2}t^3 - \frac{23}{48}t^4\right]_{t=0}^{t=1} \\ &= 3 - 1 + \frac{1}{2} - \frac{23}{48} = \frac{97}{48} \end{aligned}$$

- (d) The alternating series error bound is the absolute value of the first omitted term of the series for
- $h(1)$
- .

$$\int_0^1 \left(\frac{54}{4!}t^4\right) dt = \left[\frac{9}{20}t^5\right]_{t=0}^{t=1} = \frac{9}{20}$$

$$\text{Error} \leq \left|\frac{9}{20}\right| = 0.45$$

$$2 : \begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \end{cases}$$

$$2 : \begin{cases} 1 : \text{three terms for } e^x \\ 1 : \text{three terms for } e^x f(x) \end{cases}$$

$$2 : \begin{cases} 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : \text{uses fourth-degree term} \\ \quad \text{of Maclaurin series for } f \\ 1 : \text{uses first omitted term} \\ \quad \text{of series for } h(1) \\ 1 : \text{error bound} \end{cases}$$

AP[®] Calculus BC

2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2016 SCORING GUIDELINES

Question 1

t (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.

- (a) Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.
- (b) Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
- (c) Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
- (d) For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

(a) $R'(2) \approx \frac{R(3) - R(1)}{3 - 1} = \frac{950 - 1190}{3 - 1} = -120$ liters/hr²

2 : $\begin{cases} 1 : \text{estimate} \\ 1 : \text{units} \end{cases}$

(b) The total amount of water removed is given by $\int_0^8 R(t) dt$.

$$\begin{aligned} \int_0^8 R(t) dt &\approx 1 \cdot R(0) + 2 \cdot R(1) + 3 \cdot R(3) + 2 \cdot R(6) \\ &= 1(1340) + 2(1190) + 3(950) + 2(740) \\ &= 8050 \text{ liters} \end{aligned}$$

3 : $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{estimate} \\ 1 : \text{overestimate with reason} \end{cases}$

This is an overestimate since R is a decreasing function.

(c) Total $\approx 50000 + \int_0^8 W(t) dt - 8050$
 $= 50000 + 7836.195325 - 8050 \approx 49786$ liters

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{estimate} \end{cases}$

(d) $W(0) - R(0) > 0$, $W(8) - R(8) < 0$, and $W(t) - R(t)$ is continuous.


2 : $\begin{cases} 1 : \text{considers } W(t) - R(t) \\ 1 : \text{answer with explanation} \end{cases}$

Therefore, the Intermediate Value Theorem guarantees at least one time t , $0 < t < 8$, for which $W(t) - R(t) = 0$, or $W(t) = R(t)$.

For this value of t , the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank.

2016 SCORING GUIDELINES

Question 2



At time t , the position of a particle moving in the xy -plane is given by the parametric functions $(x(t), y(t))$, where $\frac{dx}{dt} = t^2 + \sin(3t^2)$. The graph of y , consisting of three line segments, is shown in the figure above. At $t = 0$, the particle is at position $(5, 1)$.

- (a) Find the position of the particle at $t = 3$.
- (b) Find the slope of the line tangent to the path of the particle at $t = 3$.
- (c) Find the speed of the particle at $t = 3$.
- (d) Find the total distance traveled by the particle from $t = 0$ to $t = 2$.

(a) $x(3) = x(0) + \int_0^3 x'(t) dt = 5 + 9.377035 = 14.377$

$$y(3) = -\frac{1}{2}$$

The position of the particle at $t = 3$ is $(14.377, -0.5)$.

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

(b) Slope $= \frac{y'(3)}{x'(3)} = \frac{0.5}{9.956376} = 0.05$

1 : slope

(c) Speed $= \sqrt{(x'(3))^2 + (y'(3))^2} = 9.969$ (or 9.968)

2 : $\begin{cases} 1 : \text{expression for speed} \\ 1 : \text{answer} \end{cases}$

(d) Distance $= \int_0^2 \sqrt{(x'(t))^2 + (y'(t))^2} dt$
 $= \int_0^1 \sqrt{(x'(t))^2 + (-2)^2} dt + \int_1^2 \sqrt{(x'(t))^2 + 0^2} dt$
 $= 2.237871 + 2.112003 = 4.350$ (or 4.349)

3 : $\begin{cases} 1 : \text{expression for distance} \\ 1 : \text{integrals} \\ 1 : \text{answer} \end{cases}$


2016 SCORING GUIDELINES

Question 3

The figure above shows the graph of the piecewise-linear function f . For $-4 \leq x \leq 12$, the function g is defined by

$$g(x) = \int_2^x f(t) dt.$$

- (a) Does g have a relative minimum, a relative maximum, or neither at $x = 10$? Justify your answer.
- (b) Does the graph of g have a point of inflection at $x = 4$? Justify your answer.
- (c) Find the absolute minimum value and the absolute maximum value of g on the interval $-4 \leq x \leq 12$. Justify your answers.
- (d) For $-4 \leq x \leq 12$, find all intervals for which $g(x) \leq 0$.

Graph of f

- (a) The function g has neither a relative minimum nor a relative maximum at $x = 10$ since $g'(x) = f(x)$ and $f(x) \leq 0$ for $8 \leq x \leq 12$.
- (b) The graph of g has a point of inflection at $x = 4$ since $g'(x) = f(x)$ is increasing for $2 \leq x \leq 4$ and decreasing for $4 \leq x \leq 8$.
- (c) $g'(x) = f(x)$ changes sign only at $x = -2$ and $x = 6$.

x	$g(x)$
-4	-4
-2	-8
6	8
12	-4

On the interval $-4 \leq x \leq 12$, the absolute minimum value is $g(-2) = -8$ and the absolute maximum value is $g(6) = 8$.

- (d) $g(x) \leq 0$ for $-4 \leq x \leq 2$ and $10 \leq x \leq 12$.

1 : $g'(x) = f(x)$ in (a), (b), (c), or (d)

1 : answer with justification

1 : answer with justification

4 : $\begin{cases} 1 : \text{considers } x = -2 \text{ and } x = 6 \\ \quad \text{as candidates} \\ 1 : \text{considers } x = -4 \text{ and } x = 12 \\ 2 : \text{answers with justification} \end{cases}$

2 : intervals

2016 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = x^2 - \frac{1}{2}y$.

- (a) Find $\frac{d^2y}{dx^2}$ in terms of x and y .
- (b) Let $y = f(x)$ be the particular solution to the given differential equation whose graph passes through the point $(-2, 8)$. Does the graph of f have a relative minimum, a relative maximum, or neither at the point $(-2, 8)$? Justify your answer.
- (c) Let $y = g(x)$ be the particular solution to the given differential equation with $g(-1) = 2$. Find $\lim_{x \rightarrow -1} \left(\frac{g(x) - 2}{3(x+1)^2} \right)$. Show the work that leads to your answer.
- (d) Let $y = h(x)$ be the particular solution to the given differential equation with $h(0) = 2$. Use Euler's method, starting at $x = 0$ with two steps of equal size, to approximate $h(1)$.

(a) $\frac{d^2y}{dx^2} = 2x - \frac{1}{2} \frac{dy}{dx} = 2x - \frac{1}{2} \left(x^2 - \frac{1}{2}y \right)$

(b) $\frac{dy}{dx} \Big|_{(x,y)=(-2,8)} = (-2)^2 - \frac{1}{2} \cdot 8 = 0$
 $\frac{d^2y}{dx^2} \Big|_{(x,y)=(-2,8)} = 2(-2) - \frac{1}{2} \left((-2)^2 - \frac{1}{2} \cdot 8 \right) = -4 < 0$

Thus, the graph of f has a relative maximum at the point $(-2, 8)$.

(c) $\lim_{x \rightarrow -1} (g(x) - 2) = 0$ and $\lim_{x \rightarrow -1} 3(x+1)^2 = 0$

Using L'Hospital's Rule,

$$\lim_{x \rightarrow -1} \left(\frac{g(x) - 2}{3(x+1)^2} \right) = \lim_{x \rightarrow -1} \left(\frac{g'(x)}{6(x+1)} \right)$$

$$\lim_{x \rightarrow -1} g'(x) = 0 \text{ and } \lim_{x \rightarrow -1} 6(x+1) = 0$$

Using L'Hospital's Rule,

$$\lim_{x \rightarrow -1} \left(\frac{g'(x)}{6(x+1)} \right) = \lim_{x \rightarrow -1} \left(\frac{g''(x)}{6} \right) = \frac{-2}{6} = -\frac{1}{3}$$

(d) $h\left(\frac{1}{2}\right) \approx h(0) + h'(0) \cdot \frac{1}{2} = 2 + (-1) \cdot \frac{1}{2} = \frac{3}{2}$
 $h(1) \approx h\left(\frac{1}{2}\right) + h'\left(\frac{1}{2}\right) \cdot \frac{1}{2} \approx \frac{3}{2} + \left(-\frac{1}{2}\right) \cdot \frac{1}{2} = \frac{5}{4}$

2 : $\frac{d^2y}{dx^2}$ in terms of x and y


2 : conclusion with justification

3 : $\begin{cases} 2 : \text{L'Hospital's Rule} \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{Euler's method} \\ 1 : \text{approximation} \end{cases}$

2016 SCORING GUIDELINES

Question 5



The inside of a funnel of height 10 inches has circular cross sections, as shown in the figure above. At height h , the radius of the funnel is given by $r = \frac{1}{20}(3 + h^2)$, where $0 \leq h \leq 10$. The units of r and h are inches.

- Find the average value of the radius of the funnel.
- Find the volume of the funnel.
- The funnel contains liquid that is draining from the bottom. At the instant when the height of the liquid is $h = 3$ inches, the radius of the surface of the liquid is decreasing at a rate of $\frac{1}{5}$ inch per second. At this instant, what is the rate of change of the height of the liquid with respect to time?

$$\begin{aligned} \text{(a) Average radius} &= \frac{1}{10} \int_0^{10} \frac{1}{20}(3 + h^2) dh = \frac{1}{200} \left[3h + \frac{h^3}{3} \right]_0^{10} \\ &= \frac{1}{200} \left(\left(30 + \frac{1000}{3} \right) - 0 \right) = \frac{109}{60} \text{ in} \end{aligned}$$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b) Volume} &= \pi \int_0^{10} \left(\frac{1}{20}(3 + h^2) \right)^2 dh = \frac{\pi}{400} \int_0^{10} (9 + 6h^2 + h^4) dh \\ &= \frac{\pi}{400} \left[9h + 2h^3 + \frac{h^5}{5} \right]_0^{10} \\ &= \frac{\pi}{400} \left(\left(90 + 2000 + \frac{100000}{5} \right) - 0 \right) = \frac{2209\pi}{40} \text{ in}^3 \end{aligned}$$

3 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(c) } \frac{dr}{dt} &= \frac{1}{20}(2h) \frac{dh}{dt} \\ -\frac{1}{5} &= \frac{3}{10} \frac{dh}{dt} \\ \frac{dh}{dt} &= -\frac{1}{5} \cdot \frac{10}{3} = -\frac{2}{3} \text{ in/sec} \end{aligned}$$

3 : $\begin{cases} 2 : \text{chain rule} \\ 1 : \text{answer} \end{cases}$

2016 SCORING GUIDELINES

Question 6

The function f has a Taylor series about $x = 1$ that converges to $f(x)$ for all x in the interval of convergence.

It is known that $f(1) = 1$, $f'(1) = -\frac{1}{2}$, and the n th derivative of f at $x = 1$ is given by

$$f^{(n)}(1) = (-1)^n \frac{(n-1)!}{2^n} \text{ for } n \geq 2.$$

- Write the first four nonzero terms and the general term of the Taylor series for f about $x = 1$.
- The Taylor series for f about $x = 1$ has a radius of convergence of 2. Find the interval of convergence. Show the work that leads to your answer.
- The Taylor series for f about $x = 1$ can be used to represent $f(1.2)$ as an alternating series. Use the first three nonzero terms of the alternating series to approximate $f(1.2)$.
- Show that the approximation found in part (c) is within 0.001 of the exact value of $f(1.2)$.

$$\begin{aligned} \text{(a) } f(1) &= 1, \quad f'(1) = -\frac{1}{2}, \quad f''(1) = \frac{1}{2^2}, \quad f'''(1) = -\frac{2}{2^3} \\ f(x) &= 1 - \frac{1}{2}(x-1) + \frac{1}{2^2} \cdot \frac{1}{2}(x-1)^2 - \frac{1}{2^3} \cdot \frac{1}{3}(x-1)^3 + \cdots \\ &\quad + \frac{(-1)^n}{2^n \cdot n}(x-1)^n + \cdots \end{aligned}$$

4 : $\begin{cases} 1 : \text{first two terms} \\ 1 : \text{third term} \\ 1 : \text{fourth term} \\ 1 : \text{general term} \end{cases}$

- $R = 2$. The series converges on the interval $(-1, 3)$.

When $x = -1$, the series is $1 + 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots$.

Since the harmonic series diverges, this series diverges.

When $x = 3$, the series is $1 - 1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \cdots$.

Since the alternating harmonic series converges, this series converges.

Therefore, the interval of convergence is $-1 < x \leq 3$.

$$\text{(c) } f(1.2) \approx 1 - \frac{1}{2}(0.2) + \frac{1}{8}(0.2)^2 = 1 - 0.1 + 0.005 = 0.905$$

1 : approximation

- The series for $f(1.2)$ alternates with terms that decrease in magnitude to 0.

2 : $\begin{cases} 1 : \text{error form} \\ 1 : \text{analysis} \end{cases}$

$$|f(1.2) - T_2(1.2)| \leq \left| \frac{-1}{2^3 \cdot 3}(0.2)^3 \right| = \frac{1}{3000} \leq 0.001$$

AP[®] Calculus BC

2015 Scoring Guidelines

© 2015 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.
Visit the College Board on the Web: www.collegeboard.org.
AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2015 SCORING GUIDELINES

Question 1

The rate at which rainwater flows into a drainpipe is modeled by the function R , where $R(t) = 20\sin\left(\frac{t^2}{35}\right)$ cubic feet per hour, t is measured in hours, and $0 \leq t \leq 8$. The pipe is partially blocked, allowing water to drain out the other end of the pipe at a rate modeled by $D(t) = -0.04t^3 + 0.4t^2 + 0.96t$ cubic feet per hour, for $0 \leq t \leq 8$. There are 30 cubic feet of water in the pipe at time $t = 0$.

- How many cubic feet of rainwater flow into the pipe during the 8-hour time interval $0 \leq t \leq 8$?
- Is the amount of water in the pipe increasing or decreasing at time $t = 3$ hours? Give a reason for your answer.
- At what time t , $0 \leq t \leq 8$, is the amount of water in the pipe at a minimum? Justify your answer.
- The pipe can hold 50 cubic feet of water before overflowing. For $t > 8$, water continues to flow into and out of the pipe at the given rates until the pipe begins to overflow. Write, but do not solve, an equation involving one or more integrals that gives the time w when the pipe will begin to overflow.

(a) $\int_0^8 R(t) dt = 76.570$

2 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b) $R(3) - D(3) = -0.313632 < 0$

Since $R(3) < D(3)$, the amount of water in the pipe is decreasing at time $t = 3$ hours.

2 : $\begin{cases} 1 : \text{considers } R(3) \text{ and } D(3) \\ 1 : \text{answer and reason} \end{cases}$

(c) The amount of water in the pipe at time t , $0 \leq t \leq 8$, is $30 + \int_0^t [R(x) - D(x)] dx$.

$$R(t) - D(t) = 0 \Rightarrow t = 0, 3.271658$$

t	Amount of water in the pipe
0	30
3.271658	27.964561
8	48.543686

The amount of water in the pipe is a minimum at time $t = 3.272$ (or 3.271) hours.

3 : $\begin{cases} 1 : \text{considers } R(t) - D(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

(d) $30 + \int_0^w [R(t) - D(t)] dt = 50$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{equation} \end{cases}$

2015 SCORING GUIDELINES

Question 2

At time $t \geq 0$, a particle moving along a curve in the xy -plane has position $(x(t), y(t))$ with velocity vector $\mathbf{v}(t) = (\cos(t^2), e^{0.5t})$. At $t = 1$, the particle is at the point $(3, 5)$.

- (a) Find the x -coordinate of the position of the particle at time $t = 2$.
 (b) For $0 < t < 1$, there is a point on the curve at which the line tangent to the curve has a slope of 2. At what time is the object at that point?
 (c) Find the time at which the speed of the particle is 3.
 (d) Find the total distance traveled by the particle from time $t = 0$ to time $t = 1$.

(a) $x(2) = 3 + \int_1^2 \cos(t^2) dt = 2.557$ (or 2.556)

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

(b) $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{e^{0.5t}}{\cos(t^2)}$

$$\frac{e^{0.5t}}{\cos(t^2)} = 2$$

$$t = 0.840$$

2 : $\begin{cases} 1 : \text{slope in terms of } t \\ 1 : \text{answer} \end{cases}$

(c) Speed = $\sqrt{\cos^2(t^2) + e^t}$

$$\sqrt{\cos^2(t^2) + e^t} = 3$$

$$t = 2.196$$
 (or 2.195)

2 : $\begin{cases} 1 : \text{speed in terms of } t \\ 1 : \text{answer} \end{cases}$

(d) Distance = $\int_0^1 \sqrt{\cos^2(t^2) + e^t} dt = 1.595$ (or 1.594)

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 3

t (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	-220	150

Johanna jogs along a straight path. For $0 \leq t \leq 40$, Johanna's velocity is given by a differentiable function v . Selected values of $v(t)$, where t is measured in minutes and $v(t)$ is measured in meters per minute, are given in the table above.

- (a) Use the data in the table to estimate the value of $v'(16)$.

- (b) Using correct units, explain the meaning of the definite integral $\int_0^{40} |v(t)| dt$ in the context of the problem.

Approximate the value of $\int_0^{40} |v(t)| dt$ using a right Riemann sum with the four subintervals indicated in the table.

- (c) Bob is riding his bicycle along the same path. For $0 \leq t \leq 10$, Bob's velocity is modeled by $B(t) = t^3 - 6t^2 + 300$, where t is measured in minutes and $B(t)$ is measured in meters per minute.

Find Bob's acceleration at time $t = 5$.

- (d) Based on the model B from part (c), find Bob's average velocity during the interval $0 \leq t \leq 10$.

(a) $v'(16) \approx \frac{240 - 200}{20 - 12} = 5$ meters/min²

1 : approximation

- (b) $\int_0^{40} |v(t)| dt$ is the total distance Johanna jogs, in meters, over the time interval $0 \leq t \leq 40$ minutes.

3 : $\begin{cases} 1 : \text{explanation} \\ 1 : \text{right Riemann sum} \\ 1 : \text{approximation} \end{cases}$

$$\begin{aligned} \int_0^{40} |v(t)| dt &\approx 12 \cdot |v(12)| + 8 \cdot |v(20)| + 4 \cdot |v(24)| + 16 \cdot |v(40)| \\ &= 12 \cdot 200 + 8 \cdot 240 + 4 \cdot 220 + 16 \cdot 150 \\ &= 2400 + 1920 + 880 + 2400 \\ &= 7600 \text{ meters} \end{aligned}$$

- (c) Bob's acceleration is $B'(t) = 3t^2 - 12t$.

$$B'(5) = 3(25) - 12(5) = 15 \text{ meters/min}^2$$

2 : $\begin{cases} 1 : \text{uses } B'(t) \\ 1 : \text{answer} \end{cases}$

- (d) Avg vel = $\frac{1}{10} \int_0^{10} (t^3 - 6t^2 + 300) dt$

$$= \frac{1}{10} \left[\frac{t^4}{4} - 2t^3 + 300t \right]_0^{10}$$

$$= \frac{1}{10} \left[\frac{10000}{4} - 2000 + 3000 \right] = 350 \text{ meters/min}$$


3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = 2x - y$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.
- (b) Find $\frac{d^2y}{dx^2}$ in terms of x and y . Determine the concavity of all solution curves for the given differential equation in Quadrant II. Give a reason for your answer.
- (c) Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(2) = 3$. Does f have a relative minimum, a relative maximum, or neither at $x = 2$? Justify your answer.
- (d) Find the values of the constants m and b for which $y = mx + b$ is a solution to the differential equation.



- 2 : $\begin{cases} 1 : \text{slopes where } x = 0 \\ 1 : \text{slopes where } x = 1 \end{cases}$

(b) $\frac{d^2y}{dx^2} = 2 - \frac{dy}{dx} = 2 - (2x - y) = 2 - 2x + y$

In Quadrant II, $x < 0$ and $y > 0$, so $2 - 2x + y > 0$.
Therefore, all solution curves are concave up in Quadrant II.

- 2 : $\begin{cases} 1 : \frac{d^2y}{dx^2} \\ 1 : \text{concave up with reason} \end{cases}$

(c) $\left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} = 2(2) - 3 = 1 \neq 0$

Therefore, f has neither a relative minimum nor a relative maximum at $x = 2$.

- 2 : $\begin{cases} 1 : \text{considers } \left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} \\ 1 : \text{conclusion with justification} \end{cases}$

(d) $y = mx + b \Rightarrow \frac{dy}{dx} = \frac{d}{dx}(mx + b) = m$

$$2x - y = m$$

$$2x - (mx + b) = m$$

$$(2 - m)x - (m + b) = 0$$

$$2 - m = 0 \Rightarrow m = 2$$

$$b = -m \Rightarrow b = -2$$

- 3 : $\begin{cases} 1 : \frac{d}{dx}(mx + b) = m \\ 1 : 2x - y = m \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 5

Consider the function $f(x) = \frac{1}{x^2 - kx}$, where k is a nonzero constant. The derivative of f is given by

$$f'(x) = \frac{k - 2x}{(x^2 - kx)^2}.$$

- (a) Let $k = 3$, so that $f(x) = \frac{1}{x^2 - 3x}$. Write an equation for the line tangent to the graph of f at the point whose x -coordinate is 4.
- (b) Let $k = 4$, so that $f(x) = \frac{1}{x^2 - 4x}$. Determine whether f has a relative minimum, a relative maximum, or neither at $x = 2$. Justify your answer.
- (c) Find the value of k for which f has a critical point at $x = -5$.
- (d) Let $k = 6$, so that $f(x) = \frac{1}{x^2 - 6x}$. Find the partial fraction decomposition for the function f .

Find $\int f(x) dx$.

(a) $f(4) = \frac{1}{4^2 - 3 \cdot 4} = \frac{1}{4}$ $f'(4) = \frac{3 - 2 \cdot 4}{(4^2 - 3 \cdot 4)^2} = -\frac{5}{16}$

An equation for the line tangent to the graph of f at the point whose x -coordinate is 4 is $y = -\frac{5}{16}(x - 4) + \frac{1}{4}$.

- 2 : $\begin{cases} 1 : \text{slope} \\ 1 : \text{tangent line equation} \end{cases}$

(b) $f'(x) = \frac{4 - 2x}{(x^2 - 4x)^2}$ $f'(2) = \frac{4 - 2 \cdot 2}{(2^2 - 4 \cdot 2)^2} = 0$

$f'(x)$ changes sign from positive to negative at $x = 2$.
Therefore, f has a relative maximum at $x = 2$.

- 2 : $\begin{cases} 1 : \text{considers } f'(2) \\ 1 : \text{answer with justification} \end{cases}$

(c) $f'(-5) = \frac{k - 2 \cdot (-5)}{((-5)^2 - k \cdot (-5))^2} = 0 \Rightarrow k = -10$

- 1 : answer

(d) $\frac{1}{x^2 - 6x} = \frac{1}{x(x - 6)} = \frac{A}{x} + \frac{B}{x - 6} \Rightarrow 1 = A(x - 6) + Bx$

$$x = 0 \Rightarrow 1 = A \cdot (-6) \Rightarrow A = -\frac{1}{6}$$

$$x = 6 \Rightarrow 1 = B \cdot (6) \Rightarrow B = \frac{1}{6}$$

$$\frac{1}{x(x - 6)} = \frac{-1/6}{x} + \frac{1/6}{x - 6}$$

$$\int f(x) dx = \int \left(\frac{-1/6}{x} + \frac{1/6}{x - 6} \right) dx$$

$$= -\frac{1}{6} \ln|x| + \frac{1}{6} \ln|x - 6| + C = \frac{1}{6} \ln \left| \frac{x - 6}{x} \right| + C$$

- 4 : $\begin{cases} 2 : \text{partial fraction decomposition} \\ 2 : \text{general antiderivative} \end{cases}$

2015 SCORING GUIDELINES

Question 6

The Maclaurin series for a function f is given by $\sum_{n=1}^{\infty} \frac{(-3)^{n-1}}{n} x^n = x - \frac{3}{2}x^2 + 3x^3 - \dots + \frac{(-3)^{n-1}}{n} x^n + \dots$ and converges to $f(x)$ for $|x| < R$, where R is the radius of convergence of the Maclaurin series.

- (a) Use the ratio test to find R .
- (b) Write the first four nonzero terms of the Maclaurin series for f' , the derivative of f . Express f' as a rational function for $|x| < R$.
- (c) Write the first four nonzero terms of the Maclaurin series for e^x . Use the Maclaurin series for e^x to write the third-degree Taylor polynomial for $g(x) = e^x f(x)$ about $x = 0$.

- (a) Let a_n be the n th term of the Maclaurin series.

$$\frac{a_{n+1}}{a_n} = \frac{(-3)^n x^{n+1}}{n+1} \cdot \frac{n}{(-3)^{n-1} x^n} = \frac{-3n}{n+1} \cdot x$$

$$\lim_{n \rightarrow \infty} \left| \frac{-3n}{n+1} \cdot x \right| = 3|x|$$

$$3|x| < 1 \Rightarrow |x| < \frac{1}{3}$$

The radius of convergence is $R = \frac{1}{3}$.

- (b) The first four nonzero terms of the Maclaurin series for f' are

$$1 - 3x + 9x^2 - 27x^3.$$

$$f'(x) = \frac{1}{1 - (-3x)} = \frac{1}{1 + 3x}$$

- (c) The first four nonzero terms of the Maclaurin series for e^x are

$$1 + x + \frac{x^2}{2!} + \frac{x^3}{3!}.$$

The product of the Maclaurin series for e^x and the Maclaurin series for f is

$$\left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots\right) \left(x - \frac{3}{2}x^2 + 3x^3 - \dots\right) \\ = x - \frac{1}{2}x^2 + 2x^3 + \dots$$

The third-degree Taylor polynomial for $g(x) = e^x f(x)$

about $x = 0$ is $T_3(x) = x - \frac{1}{2}x^2 + 2x^3$.

1 : sets up ratio
1 : computes limit of ratio
1 : determines radius of convergence

2 : first four nonzero terms
1 : rational function

1 : first four nonzero terms
of the Maclaurin series for e^x
2 : Taylor polynomial