

6. The Taylor series for a function f about $x = 4$ is given by

$\sum_{n=1}^{\infty} \frac{(x-4)^{n+1}}{(n+1)3^n} = \frac{(x-4)^2}{2 \cdot 3} + \frac{(x-4)^3}{3 \cdot 3^2} + \frac{(x-4)^4}{4 \cdot 3^3} + \dots + \frac{(x-4)^{n+1}}{(n+1)3^n} + \dots$ and converges to $f(x)$ on its interval of convergence.

- A. Using the ratio test, find the interval of convergence of the Taylor series for f about $x = 4$. Justify your answer.
- B. Find the first three nonzero terms and the general term of the Taylor series for f' , the derivative of f , about $x = 4$.
- C. The Taylor series for f' described in part B is a geometric series. For all x in the interval of convergence of the Taylor series for f' , show that $f'(x) = \frac{x-4}{7-x}$.
- D. It is known that the radius of convergence of the Taylor series for f about $x = 4$ is the same as the radius of convergence of the Taylor series for f' about $x = 4$. Does the Taylor series for f' described in part B converge to $f'(x) = \frac{x-4}{7-x}$ at $x = 8$? Give a reason for your answer.

STOP
END OF EXAM

6. The Maclaurin series for a function f is given by $\sum_{n=1}^{\infty} \frac{(n+1)x^n}{n^2 6^n}$ and converges to $f(x)$ for all x in the interval of convergence. It can be shown that the Maclaurin series for f has a radius of convergence of 6.

- (a) Determine whether the Maclaurin series for f converges or diverges at $x = 6$. Give a reason for your answer.
- (b) It can be shown that $f(-3) = \sum_{n=1}^{\infty} \frac{(n+1)(-3)^n}{n^2 6^n} = \sum_{n=1}^{\infty} \frac{n+1}{n^2} \left(-\frac{1}{2}\right)^n$ and that the first three terms of this series sum to $S_3 = -\frac{125}{144}$. Show that $\left|f(-3) - S_3\right| < \frac{1}{50}$.
- (c) Find the general term of the Maclaurin series for f' , the derivative of f . Find the radius of convergence of the Maclaurin series for f' .
- (d) Let $g(x) = \sum_{n=1}^{\infty} \frac{(n+1)x^{2n}}{n^2 3^n}$. Use the ratio test to determine the radius of convergence of the Maclaurin series for g .

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

STOP
END OF EXAM

6. The function f is defined by the power series $f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots + \frac{(-1)^n x^{2n+1}}{2n+1} + \cdots$ for all real numbers x for which the series converges.

(a) Using the ratio test, find the interval of convergence of the power series for f . Justify your answer.

(b) Show that $\left| f\left(\frac{1}{2}\right) - \frac{1}{2} \right| < \frac{1}{10}$. Justify your answer.

(c) Write the first four nonzero terms and the general term for an infinite series that represents $f'(x)$.

(d) Use the result from part (c) to find the value of $f'\left(\frac{1}{6}\right)$.

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

STOP

END OF EXAM

6. The function g has derivatives of all orders for all real numbers. The Maclaurin series for g is given by

$$g(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^n}{2e^n + 3} \text{ on its interval of convergence.}$$

(a) State the conditions necessary to use the integral test to determine convergence of the series $\sum_{n=0}^{\infty} \frac{1}{e^n}$.

Use the integral test to show that $\sum_{n=0}^{\infty} \frac{1}{e^n}$ converges.

(b) Use the limit comparison test with the series $\sum_{n=0}^{\infty} \frac{1}{e^n}$ to show that the series $g(1) = \sum_{n=0}^{\infty} \frac{(-1)^n}{2e^n + 3}$

converges absolutely.

(c) Determine the radius of convergence of the Maclaurin series for g .

(d) The first two terms of the series $g(1) = \sum_{n=0}^{\infty} \frac{(-1)^n}{2e^n + 3}$ are used to approximate $g(1)$. Use the alternating

series error bound to determine an upper bound on the error of the approximation.

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

STOP

END OF EXAM

6. The Maclaurin series for $\ln(1+x)$ is given by

$$x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots + (-1)^{n+1} \frac{x^n}{n} + \cdots$$

On its interval of convergence, this series converges to $\ln(1+x)$. Let f be the function defined by

$$f(x) = x \ln\left(1 + \frac{x}{3}\right).$$

- (a) Write the first four nonzero terms and the general term of the Maclaurin series for f .
- (b) Determine the interval of convergence of the Maclaurin series for f . Show the work that leads to your answer.
- (c) Let $P_4(x)$ be the fourth-degree Taylor polynomial for f about $x = 0$. Use the alternating series error bound to find an upper bound for $|P_4(2) - f(2)|$.

STOP
END OF EXAM

6. The function f has a Taylor series about $x = 1$ that converges to $f(x)$ for all x in the interval of convergence.

It is known that $f(1) = 1$, $f'(1) = -\frac{1}{2}$, and the n th derivative of f at $x = 1$ is given by $f^{(n)}(1) = (-1)^n \frac{(n-1)!}{2^n}$ for $n \geq 2$.

- (a) Write the first four nonzero terms and the general term of the Taylor series for f about $x = 1$.
- (b) The Taylor series for f about $x = 1$ has a radius of convergence of 2. Find the interval of convergence. Show the work that leads to your answer.
- (c) The Taylor series for f about $x = 1$ can be used to represent $f(1.2)$ as an alternating series. Use the first three nonzero terms of the alternating series to approximate $f(1.2)$.
- (d) Show that the approximation found in part (c) is within 0.001 of the exact value of $f(1.2)$.

STOP
END OF EXAM

6. The Maclaurin series for a function f is given by $\sum_{n=1}^{\infty} \frac{(-3)^{n-1}}{n} x^n = x - \frac{3}{2}x^2 + 3x^3 - \cdots + \frac{(-3)^{n-1}}{n} x^n + \cdots$ and converges to $f(x)$ for $|x| < R$, where R is the radius of convergence of the Maclaurin series.
- (a) Use the ratio test to find R .
- (b) Write the first four nonzero terms of the Maclaurin series for f' , the derivative of f . Express f' as a rational function for $|x| < R$.
- (c) Write the first four nonzero terms of the Maclaurin series for e^x . Use the Maclaurin series for e^x to write the third-degree Taylor polynomial for $g(x) = e^x f(x)$ about $x = 0$.

STOP**END OF EXAM**

6. The Taylor series for a function f about $x = 1$ is given by $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{f^{(n)}(1)}{n!} (x-1)^n$ and converges to $f(x)$ for $|x-1| < R$, where R is the radius of convergence of the Taylor series.
- (a) Find the value of R .
- (b) Find the first three nonzero terms and the general term of the Taylor series for f' , the derivative of f , about $x = 1$.
- (c) The Taylor series for f' about $x = 1$, found in part (b), is a geometric series. Find the function f' to which the series converges for $|x-1| < R$. Use this function to determine f for $|x-1| < R$.

STOP**END OF EXAM**