

Part B (BC): Graphing calculator not allowed**Question 6****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

The function f is defined by the power series $f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots + \frac{(-1)^n x^{2n+1}}{2n+1} + \cdots$ for all real numbers x for which the series converges.

Model Solution**Scoring**

- (a) Using the ratio test, find the interval of convergence of the power series for f . Justify your answer.

$\lim_{n \rightarrow \infty} \left \frac{(-1)^{n+1} x^{2n+3}}{2n+3} \cdot \frac{2n+1}{(-1)^n x^{2n+1}} \right = \lim_{n \rightarrow \infty} \left \frac{x^{2n+3}}{2n+3} \cdot \frac{2n+1}{x^{2n+1}} \right = \lim_{n \rightarrow \infty} \left x^2 \left(\frac{2n+1}{2n+3} \right) \right = x^2 $	Sets up ratio	1 point
$ x^2 < 1$ for $ x < 1$. The series converges when $-1 < x < 1$.	Identifies interior of interval of convergence	1 point
When $x = -1$, the series is $-1 + \frac{1}{3} - \frac{1}{5} + \cdots + \frac{(-1)^{n+1}}{2n+1} + \cdots$.	Considers both endpoints	1 point
The series is an alternating series whose terms decrease in absolute value to 0. The series converges by the Alternating Series Test.	Analysis and interval of convergence	1 point
When $x = 1$, the series is $1 - \frac{1}{3} + \frac{1}{5} - \cdots + \frac{(-1)^n}{2n+1} + \cdots$.		
The series is an alternating series whose terms decrease in absolute value to 0. The series converges by the Alternating Series Test.		
Therefore, the interval of convergence is $-1 \leq x \leq 1$.		

Scoring notes:

- A response that includes the substitution error of the form $x^{2(n+1)+1} = x^{2n+3}$ appearing as $x^{2n+1+1} = x^{2n+2}$ in setting up a ratio is eligible for the first 3 points but does not earn the fourth point.
- The first point is earned by presenting a correct ratio with or without absolute values.
- To earn the second point a response must:
 - use the absolute value of the ratio, or resolve the lack of absolute values by concluding $x^2 < 1$ (without any errors), and correctly evaluate the limit of the ratio, including correct limit notation, and
 - identify the interior of the interval of convergence. The response can use either interval notation or the compound inequality $-1 < x < 1$ ($|x| < 1$ is insufficient).
- The only incorrect interval eligible for the third point is $0 < x < 1$. In this case, to earn the third point, the response needs to evaluate the general term at $x = 1$.

Total for part (a) 4 points

- (b) Show that $\left| f\left(\frac{1}{2}\right) - \frac{1}{2} \right| < \frac{1}{10}$. Justify your answer.

The series for $f\left(\frac{1}{2}\right)$ is an alternating series whose terms decrease in absolute value to 0. The first term of the series for $f\left(\frac{1}{2}\right)$ is $\frac{1}{2}$.	Uses second term	1 point
Using the alternating series error bound, $f\left(\frac{1}{2}\right)$ differs from $\frac{1}{2}$ by at most the absolute value of the second term of the series.	Justification	1 point
$\left f\left(\frac{1}{2}\right) - \frac{1}{2} \right < \left \frac{(-1)^1 \left(\frac{1}{2}\right)^3}{3} \right = \frac{1}{24} < \frac{1}{10}$		

Scoring notes:

- The first point is earned by correctly using $x = \frac{1}{2}$ in the second term (listing the second term as part of a polynomial is insufficient). Using $x = \frac{1}{2}$ in any term of degree five or higher does not earn this point.
- To earn the second point a response must:
 - have earned the first point,
 - state that the series is alternating and that its terms decrease to zero, and
 - present the inequality $\text{Error} < \frac{1}{24} < \frac{1}{10}$ (or the equivalent).
- A response that states $\text{Error} = \frac{1}{24}$ does not earn the second point.

Total for part (b) 2 points

- (c) Write the first four nonzero terms and the general term for an infinite series that represents $f'(x)$.

$f'(x) = 1 - x^2 + x^4 - x^6 + \cdots + (-1)^n x^{2n} + \cdots$	First four terms	1 point
	General term	1 point

Scoring notes:

- The first point is earned by presenting the first four nonzero terms in a list or as part of a polynomial or series.
- The second point is earned by identifying the general term (either individually or as part of a polynomial or series).
- Read “=” as “ $\approx$ ” as necessary.

Total for part (c) 2 points

- (d) Use the result from part (c) to find the value of $f'\left(\frac{1}{6}\right)$.

$f'\left(\frac{1}{6}\right) = 1 - \left(\frac{1}{6}\right)^2 + \left(\frac{1}{6}\right)^4 - \left(\frac{1}{6}\right)^6 + \cdots$ $f'\left(\frac{1}{6}\right) \text{ is a geometric series with } a = 1 \text{ and } r = -\frac{1}{36}.$ $f'\left(\frac{1}{6}\right) = \frac{a}{1-r} = \frac{1}{1-\left(-\frac{1}{36}\right)} = \frac{1}{\frac{37}{36}} = \frac{36}{37}$	Answer	1 point
---	--------	---------

Scoring notes:

- The result from part (c) must be geometric in order to be eligible for this point.
- If a response imports an incorrect geometric series from part (c), this point is earned only for a consistent answer.

Total for part (d) 1 point

Total for question 6 9 points

2017

AP[®] CollegeBoard

AP Calculus BC

Scoring Guidelines

2017 SCORING GUIDELINES

Question 1

	1 : units in parts (a), (c), and (d)
(a) Volume = $\int_0^{10} A(h) \, dh$ $\approx (2 - 0) \cdot A(0) + (5 - 2) \cdot A(2) + (10 - 5) \cdot A(5)$ $= 2 \cdot 50.3 + 3 \cdot 14.4 + 5 \cdot 6.5$ $= 176.3 \text{ cubic feet}$	2 : $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{approximation} \end{cases}$
(b) The approximation in part (a) is an overestimate because a left Riemann sum is used and A is decreasing.	1 : overestimate with reason
(c) $\int_0^{10} f(h) \, dh = 101.325338$ The volume is 101.325 cubic feet.	2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$
(d) Using the model, $V(h) = \int_0^h f(x) \, dx$. $\left. \frac{dV}{dt} \right _{h=5} = \left[\frac{dV}{dh} \cdot \frac{dh}{dt} \right]_{h=5}$ $= \left[f(h) \cdot \frac{dh}{dt} \right]_{h=5}$ $= f(5) \cdot 0.26 = 1.694419$ When $h = 5$, the volume of water is changing at a rate of 1.694 cubic feet per minute.	3 : $\begin{cases} 2 : \frac{dV}{dt} \\ 1 : \text{answer} \end{cases}$

2017 SCORING GUIDELINES

Question 2

$$(a) \frac{1}{2} \int_0^{\pi/2} (f(\theta))^2 d\theta = 0.648414$$

The area of R is 0.648.

$$(b) \int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \frac{1}{2} \int_0^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

— OR —

$$\int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \int_k^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

$$(c) w(\theta) = g(\theta) - f(\theta)$$

$$w_A = \frac{\int_0^{\pi/2} w(\theta) d\theta}{\frac{\pi}{2} - 0} = 0.485446$$

The average value of $w(\theta)$ on the interval $\left[0, \frac{\pi}{2}\right]$ is 0.485.

$$(d) w(\theta) = w_A \text{ for } 0 \leq \theta \leq \frac{\pi}{2} \Rightarrow \theta = 0.517688$$

$$w(\theta) = w_A \text{ at } \theta = 0.518 \text{ (or } 0.517\text{)}.$$

$$w'(0.518) < 0 \Rightarrow w(\theta) \text{ is decreasing at } \theta = 0.518.$$

$$2: \begin{cases} 1: \text{integral} \\ 1: \text{answer} \end{cases}$$

$$2: \begin{cases} 1: \text{integral expression} \\ \quad \text{for one region} \\ 1: \text{equation} \end{cases}$$

$$3: \begin{cases} 1: w(\theta) \\ 1: \text{integral} \\ 1: \text{average value} \end{cases}$$

$$2: \begin{cases} 1: \text{solves } w(\theta) = w_A \\ 1: \text{answer with reason} \end{cases}$$

2017 SCORING GUIDELINES

Question 3

$$(a) f(-6) = f(-2) + \int_{-2}^{-6} f'(x) dx = 7 - \int_{-6}^{-2} f'(x) dx = 7 - 4 = 3$$

$$f(5) = f(-2) + \int_{-2}^5 f'(x) dx = 7 - 2\pi + 3 = 10 - 2\pi$$

$$(b) f'(x) > 0 \text{ on the intervals } [-6, -2) \text{ and } (2, 5).$$

Therefore, f is increasing on the intervals $[-6, -2]$ and $[2, 5]$.

$$(c) \text{The absolute minimum will occur at a critical point where } f'(x) = 0 \text{ or at an endpoint.}$$

$$f'(x) = 0 \Rightarrow x = -2, x = 2$$

x	$f(x)$
-6	3
-2	7
2	$7 - 2\pi$
5	$10 - 2\pi$

The absolute minimum value is $f(2) = 7 - 2\pi$.

$$(d) f''(-5) = \frac{2 - 0}{-6 - (-2)} = -\frac{1}{2}$$

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} = 2 \text{ and } \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3} = -1$$

$f''(3)$ does not exist because

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} \neq \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3}.$$

$$3: \begin{cases} 1: \text{uses initial condition} \\ 1: f(-6) \\ 1: f(5) \end{cases}$$

$$2: \text{answer with justification}$$

$$2: \begin{cases} 1: \text{considers } x = 2 \\ 1: \text{answer with justification} \end{cases}$$

$$2: \begin{cases} 1: f''(-5) \\ 1: f''(3) \text{ does not exist,} \\ \quad \text{with explanation} \end{cases}$$

2017 SCORING GUIDELINES

Question 4

(a) $H'(0) = -\frac{1}{4}(91 - 27) = -16$
 $H(0) = 91$

An equation for the tangent line is $y = 91 - 16t$.

The internal temperature of the potato at time $t = 3$ minutes is approximately $91 - 16 \cdot 3 = 43$ degrees Celsius.

(b) $\frac{d^2H}{dt^2} = -\frac{1}{4} \frac{dH}{dt} = \left(-\frac{1}{4}\right)\left(-\frac{1}{4}\right)(H - 27) = \frac{1}{16}(H - 27)$

$$H > 27 \text{ for } t > 0 \Rightarrow \frac{d^2H}{dt^2} = \frac{1}{16}(H - 27) > 0 \text{ for } t > 0$$

Therefore, the graph of H is concave up for $t > 0$. Thus, the answer in part (a) is an underestimate.

(c) $\frac{dG}{(G - 27)^{2/3}} = -dt$
 $\int \frac{dG}{(G - 27)^{2/3}} = \int (-1) dt$
 $3(G - 27)^{1/3} = -t + C$
 $3(91 - 27)^{1/3} = 0 + C \Rightarrow C = 12$
 $3(G - 27)^{1/3} = 12 - t$
 $G(t) = 27 + \left(\frac{12 - t}{3}\right)^3 \text{ for } 0 \leq t < 10$

The internal temperature of the potato at time $t = 3$ minutes is $27 + \left(\frac{12 - 3}{3}\right)^3 = 54$ degrees Celsius.

3 : $\begin{cases} 1 : \text{slope} \\ 1 : \text{tangent line} \\ 1 : \text{approximation} \end{cases}$

1 : underestimate with reason

5 : $\begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration and uses initial condition} \\ 1 : \text{equation involving } G \text{ and } t \\ 1 : G(t) \text{ and } G(3) \end{cases}$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

2017 SCORING GUIDELINES

Question 5

(a) $f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2}$

$$f'(3) = \frac{(-3)(5)}{(18 - 21 + 5)^2} = -\frac{15}{4}$$

(b) $f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2} = 0 \Rightarrow x = \frac{7}{4}$

The only critical point in the interval $1 < x < 2.5$ has x -coordinate $\frac{7}{4}$.

f' changes sign from positive to negative at $x = \frac{7}{4}$.

Therefore, f has a relative maximum at $x = \frac{7}{4}$.

(c) $\int_5^\infty f(x) dx = \lim_{b \rightarrow \infty} \int_5^b \frac{3}{2x^2 - 7x + 5} dx = \lim_{b \rightarrow \infty} \int_5^b \left(\frac{2}{2x - 5} - \frac{1}{x - 1} \right) dx$
 $= \lim_{b \rightarrow \infty} \left[\ln(2x - 5) - \ln(x - 1) \right]_5^b = \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2x - 5}{x - 1} \right) \right]_5^b$
 $= \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2b - 5}{b - 1} \right) - \ln \left(\frac{5}{4} \right) \right] = \ln 2 - \ln \left(\frac{5}{4} \right) = \ln \left(\frac{8}{5} \right)$

(d) f is continuous, positive, and decreasing on $[5, \infty)$.

The series converges by the integral test since $\int_5^\infty \frac{3}{2x^2 - 7x + 5} dx$ converges.

— OR —

$$\frac{3}{2n^2 - 7n + 5} > 0 \text{ and } \frac{1}{n^2} > 0 \text{ for } n \geq 5.$$

Since $\lim_{n \rightarrow \infty} \frac{\frac{3}{2n^2 - 7n + 5}}{\frac{1}{n^2}} = \frac{3}{2}$ and the series $\sum_{n=5}^\infty \frac{1}{n^2}$ converges,

the series $\sum_{n=5}^\infty \frac{3}{2n^2 - 7n + 5}$ converges by the limit comparison test.

2 : $f'(3)$

2 : $\begin{cases} 1 : x\text{-coordinate} \\ 1 : \text{relative maximum with justification} \end{cases}$

3 : $\begin{cases} 1 : \text{antiderivative} \\ 1 : \text{limit expression} \\ 1 : \text{answer} \end{cases}$

2 : answer with conditions

2017 SCORING GUIDELINES

Question 6

(a) $f(0) = 0$

$f'(0) = 1$

$f''(0) = -1(1) = -1$

$f'''(0) = -2(-1) = 2$

$f^{(4)}(0) = -3(2) = -6$

The first four nonzero terms are

$$0 + 1x + \frac{-1}{2!}x^2 + \frac{2}{3!}x^3 + \frac{-6}{4!}x^4 = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4}.$$

The general term is $\frac{(-1)^{n+1}x^n}{n}$.

(b) For $x = 1$, the Maclaurin series becomes $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n}$.

The series does not converge absolutely because the harmonic series diverges.

The series alternates with terms that decrease in magnitude to 0, and therefore the series converges conditionally.

(c)
$$\begin{aligned} \int_0^x f(t) dt &= \int_0^x \left(t - \frac{t^2}{2} + \frac{t^3}{3} - \frac{t^4}{4} + \cdots + \frac{(-1)^{n+1}t^n}{n} + \cdots \right) dt \\ &= \left[\frac{t^2}{2} - \frac{t^3}{3 \cdot 2} + \frac{t^4}{4 \cdot 3} - \frac{t^5}{5 \cdot 4} + \cdots + \frac{(-1)^{n+1}t^{n+1}}{(n+1)n} + \cdots \right]_{t=0}^{t=x} \\ &= \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{12} - \frac{x^5}{20} + \cdots + \frac{(-1)^{n+1}x^{n+1}}{(n+1)n} + \cdots \end{aligned}$$

(d) The terms alternate in sign and decrease in magnitude to 0. By the alternating series error bound, the error $\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right|$ is boundedby the magnitude of the first unused term, $\left| -\frac{(1/2)^5}{20} \right|$.

Thus, $\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right| \leq \left| -\frac{(1/2)^5}{20} \right| = \frac{1}{32 \cdot 20} < \frac{1}{500}$.

3 : $\begin{cases} 1 : f''(0), f'''(0), \text{ and } f^{(4)}(0) \\ 1 : \text{verify terms} \\ 1 : \text{general term} \end{cases}$

2 : converges conditionally
with reason

3 : $\begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \\ 1 : \text{general term} \end{cases}$

1 : error bound

AP[®] Calculus BC
2015 Scoring Guidelines

© 2015 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2015 SCORING GUIDELINES

Question 1

The rate at which rainwater flows into a drainpipe is modeled by the function R , where $R(t) = 20\sin\left(\frac{t^2}{35}\right)$ cubic feet per hour, t is measured in hours, and $0 \leq t \leq 8$. The pipe is partially blocked, allowing water to drain out the other end of the pipe at a rate modeled by $D(t) = -0.04t^3 + 0.4t^2 + 0.96t$ cubic feet per hour, for $0 \leq t \leq 8$. There are 30 cubic feet of water in the pipe at time $t = 0$.

- (a) How many cubic feet of rainwater flow into the pipe during the 8-hour time interval $0 \leq t \leq 8$?
- (b) Is the amount of water in the pipe increasing or decreasing at time $t = 3$ hours? Give a reason for your answer.
- (c) At what time t , $0 \leq t \leq 8$, is the amount of water in the pipe at a minimum? Justify your answer.
- (d) The pipe can hold 50 cubic feet of water before overflowing. For $t > 8$, water continues to flow into and out of the pipe at the given rates until the pipe begins to overflow. Write, but do not solve, an equation involving one or more integrals that gives the time w when the pipe will begin to overflow.

(a) $\int_0^8 R(t) dt = 76.570$

2 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b) $R(3) - D(3) = -0.313632 < 0$
 Since $R(3) < D(3)$, the amount of water in the pipe is decreasing at time $t = 3$ hours.

2 : $\begin{cases} 1 : \text{considers } R(3) \text{ and } D(3) \\ 1 : \text{answer and reason} \end{cases}$

(c) The amount of water in the pipe at time t , $0 \leq t \leq 8$, is $30 + \int_0^t [R(x) - D(x)] dx$.

3 : $\begin{cases} 1 : \text{considers } R(t) - D(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

$$R(t) - D(t) = 0 \Rightarrow t = 0, 3.271658$$

t	Amount of water in the pipe
0	30
3.271658	27.964561
8	48.543686

The amount of water in the pipe is a minimum at time $t = 3.272$ (or 3.271) hours.

(d) $30 + \int_0^w [R(t) - D(t)] dt = 50$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{equation} \end{cases}$

2015 SCORING GUIDELINES

Question 2

At time $t \geq 0$, a particle moving along a curve in the xy -plane has position $(x(t), y(t))$ with velocity vector $v(t) = (\cos(t^2), e^{0.5t})$. At $t = 1$, the particle is at the point $(3, 5)$.

- (a) Find the x -coordinate of the position of the particle at time $t = 2$.
- (b) For $0 < t < 1$, there is a point on the curve at which the line tangent to the curve has a slope of 2. At what time is the object at that point?
- (c) Find the time at which the speed of the particle is 3.
- (d) Find the total distance traveled by the particle from time $t = 0$ to time $t = 1$.

(a) $x(2) = 3 + \int_1^2 \cos(t^2) dt = 2.557$ (or 2.556)

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

(b) $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{e^{0.5t}}{\cos(t^2)}$

$$\frac{e^{0.5t}}{\cos(t^2)} = 2$$

$$t = 0.840$$

2 : $\begin{cases} 1 : \text{slope in terms of } t \\ 1 : \text{answer} \end{cases}$

(c) Speed = $\sqrt{\cos^2(t^2) + e^t}$

$$\sqrt{\cos^2(t^2) + e^t} = 3$$

$$t = 2.196$$
 (or 2.195)

2 : $\begin{cases} 1 : \text{speed in terms of } t \\ 1 : \text{answer} \end{cases}$

(d) Distance = $\int_0^1 \sqrt{\cos^2(t^2) + e^t} dt = 1.595$ (or 1.594)

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 3

t (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	-220	150

Johanna jogs along a straight path. For $0 \leq t \leq 40$, Johanna's velocity is given by a differentiable function v . Selected values of $v(t)$, where t is measured in minutes and $v(t)$ is measured in meters per minute, are given in the table above.

- (a) Use the data in the table to estimate the value of $v'(16)$.
- (b) Using correct units, explain the meaning of the definite integral $\int_0^{40} |v(t)| dt$ in the context of the problem.

Approximate the value of $\int_0^{40} |v(t)| dt$ using a right Riemann sum with the four subintervals indicated in the table.

- (c) Bob is riding his bicycle along the same path. For $0 \leq t \leq 10$, Bob's velocity is modeled by $B(t) = t^3 - 6t^2 + 300$, where t is measured in minutes and $B(t)$ is measured in meters per minute.

Find Bob's acceleration at time $t = 5$.

- (d) Based on the model B from part (c), find Bob's average velocity during the interval $0 \leq t \leq 10$.

(a) $v'(16) \approx \frac{240 - 200}{20 - 12} = 5$ meters/min²

1 : approximation

- (b) $\int_0^{40} |v(t)| dt$ is the total distance Johanna jogs, in meters, over the time interval $0 \leq t \leq 40$ minutes.

3 : $\begin{cases} 1 : \text{explanation} \\ 1 : \text{right Riemann sum} \\ 1 : \text{approximation} \end{cases}$

$$\begin{aligned} \int_0^{40} |v(t)| dt &\approx 12 \cdot |v(12)| + 8 \cdot |v(20)| + 4 \cdot |v(24)| + 16 \cdot |v(40)| \\ &= 12 \cdot 200 + 8 \cdot 240 + 4 \cdot 220 + 16 \cdot 150 \\ &= 2400 + 1920 + 880 + 2400 \\ &= 7600 \text{ meters} \end{aligned}$$

- (c) Bob's acceleration is $B'(t) = 3t^2 - 12t$.

$$B'(5) = 3(25) - 12(5) = 15 \text{ meters/min}^2$$

2 : $\begin{cases} 1 : \text{uses } B'(t) \\ 1 : \text{answer} \end{cases}$

- (d) Avg vel = $\frac{1}{10} \int_0^{10} (t^3 - 6t^2 + 300) dt$

$$\begin{aligned} &= \frac{1}{10} \left[\frac{t^4}{4} - 2t^3 + 300t \right]_0^{10} \\ &= \frac{1}{10} \left[\frac{10000}{4} - 2000 + 3000 \right] = 350 \text{ meters/min} \end{aligned}$$


3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = 2x - y$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.
- (b) Find $\frac{d^2y}{dx^2}$ in terms of x and y . Determine the concavity of all solution curves for the given differential equation in Quadrant II. Give a reason for your answer.
- (c) Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(2) = 3$. Does f have a relative minimum, a relative maximum, or neither at $x = 2$? Justify your answer.
- (d) Find the values of the constants m and b for which $y = mx + b$ is a solution to the differential equation.



2 : $\begin{cases} 1 : \text{slopes where } x = 0 \\ 1 : \text{slopes where } x = 1 \end{cases}$

(b) $\frac{d^2y}{dx^2} = 2 - \frac{dy}{dx} = 2 - (2x - y) = 2 - 2x + y$

In Quadrant II, $x < 0$ and $y > 0$, so $2 - 2x + y > 0$.

Therefore, all solution curves are concave up in Quadrant II.

2 : $\begin{cases} 1 : \frac{d^2y}{dx^2} \\ 1 : \text{concave up with reason} \end{cases}$

(c) $\left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} = 2(2) - 3 = 1 \neq 0$

Therefore, f has neither a relative minimum nor a relative maximum at $x = 2$.

2 : $\begin{cases} 1 : \text{considers } \left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} \\ 1 : \text{conclusion with justification} \end{cases}$

(d) $y = mx + b \Rightarrow \frac{dy}{dx} = \frac{d}{dx}(mx + b) = m$

$$2x - y = m$$

$$2x - (mx + b) = m$$

$$(2 - m)x - (m + b) = 0$$

$$2 - m = 0 \Rightarrow m = 2$$

$$b = -m \Rightarrow b = -2$$

3 : $\begin{cases} 1 : \frac{d}{dx}(mx + b) = m \\ 1 : 2x - y = m \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 5

Consider the function $f(x) = \frac{1}{x^2 - kx}$, where k is a nonzero constant. The derivative of f is given by

$$f'(x) = \frac{k - 2x}{(x^2 - kx)^2}.$$

- (a) Let $k = 3$, so that $f(x) = \frac{1}{x^2 - 3x}$. Write an equation for the line tangent to the graph of f at the point whose x -coordinate is 4.
- (b) Let $k = 4$, so that $f(x) = \frac{1}{x^2 - 4x}$. Determine whether f has a relative minimum, a relative maximum, or neither at $x = 2$. Justify your answer.
- (c) Find the value of k for which f has a critical point at $x = -5$.
- (d) Let $k = 6$, so that $f(x) = \frac{1}{x^2 - 6x}$. Find the partial fraction decomposition for the function f .

Find $\int f(x) dx$.

(a) $f(4) = \frac{1}{4^2 - 3 \cdot 4} = \frac{1}{4}$ $f'(4) = \frac{3 - 2 \cdot 4}{(4^2 - 3 \cdot 4)^2} = -\frac{5}{16}$

An equation for the line tangent to the graph of f at the point whose x -coordinate is 4 is $y = -\frac{5}{16}(x - 4) + \frac{1}{4}$.

(b) $f'(x) = \frac{4 - 2x}{(x^2 - 4x)^2}$ $f'(2) = \frac{4 - 2 \cdot 2}{(2^2 - 4 \cdot 2)^2} = 0$

$f'(x)$ changes sign from positive to negative at $x = 2$.
Therefore, f has a relative maximum at $x = 2$.

(c) $f'(-5) = \frac{k - 2 \cdot (-5)}{((-5)^2 - k \cdot (-5))^2} = 0 \Rightarrow k = -10$

(d) $\frac{1}{x^2 - 6x} = \frac{1}{x(x - 6)} = \frac{A}{x} + \frac{B}{x - 6} \Rightarrow 1 = A(x - 6) + Bx$

$$x = 0 \Rightarrow 1 = A \cdot (-6) \Rightarrow A = -\frac{1}{6}$$

$$x = 6 \Rightarrow 1 = B \cdot (6) \Rightarrow B = \frac{1}{6}$$

$$\frac{1}{x(x - 6)} = \frac{-1/6}{x} + \frac{1/6}{x - 6}$$

$$\int f(x) dx = \int \left(\frac{-1/6}{x} + \frac{1/6}{x - 6} \right) dx$$

$$= -\frac{1}{6} \ln|x| + \frac{1}{6} \ln|x - 6| + C = \frac{1}{6} \ln \left| \frac{x - 6}{x} \right| + C$$

2 : $\begin{cases} 1 : \text{slope} \\ 1 : \text{tangent line equation} \end{cases}$

2 : $\begin{cases} 1 : \text{considers } f'(2) \\ 1 : \text{answer with justification} \end{cases}$

1 : answer

4 : $\begin{cases} 2 : \text{partial fraction decomposition} \\ 2 : \text{general antiderivative} \end{cases}$

2015 SCORING GUIDELINES

Question 6

The Maclaurin series for a function f is given by $\sum_{n=1}^{\infty} \frac{(-3)^{n-1}}{n} x^n = x - \frac{3}{2}x^2 + 3x^3 - \dots + \frac{(-3)^{n-1}}{n} x^n + \dots$ and converges to $f(x)$ for $|x| < R$, where R is the radius of convergence of the Maclaurin series.

- (a) Use the ratio test to find R .
- (b) Write the first four nonzero terms of the Maclaurin series for f' , the derivative of f . Express f' as a rational function for $|x| < R$.
- (c) Write the first four nonzero terms of the Maclaurin series for e^x . Use the Maclaurin series for e^x to write the third-degree Taylor polynomial for $g(x) = e^x f(x)$ about $x = 0$.

- (a) Let a_n be the n th term of the Maclaurin series.

$$\frac{a_{n+1}}{a_n} = \frac{(-3)^n x^{n+1}}{n+1} \cdot \frac{n}{(-3)^{n-1} x^n} = \frac{-3n}{n+1} \cdot x$$

$$\lim_{n \rightarrow \infty} \left| \frac{-3n}{n+1} \cdot x \right| = 3|x|$$

$$3|x| < 1 \Rightarrow |x| < \frac{1}{3}$$

The radius of convergence is $R = \frac{1}{3}$.

- (b) The first four nonzero terms of the Maclaurin series for f' are $1 - 3x + 9x^2 - 27x^3$.

$$f'(x) = \frac{1}{1 - (-3x)} = \frac{1}{1 + 3x}$$

- (c) The first four nonzero terms of the Maclaurin series for e^x are $1 + x + \frac{x^2}{2!} + \frac{x^3}{3!}$.

The product of the Maclaurin series for e^x and the Maclaurin series for f' is

$$\left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \right) \left(x - \frac{3}{2}x^2 + 3x^3 - \dots \right)$$

$$= x - \frac{1}{2}x^2 + 2x^3 + \dots$$

The third-degree Taylor polynomial for $g(x) = e^x f(x)$ about $x = 0$ is $T_3(x) = x - \frac{1}{2}x^2 + 2x^3$.

3 : $\begin{cases} 1 : \text{sets up ratio} \\ 1 : \text{computes limit of ratio} \\ 1 : \text{determines radius of convergence} \end{cases}$

3 : $\begin{cases} 2 : \text{first four nonzero terms} \\ 1 : \text{rational function} \end{cases}$

3 : $\begin{cases} 1 : \text{first four nonzero terms} \\ \quad \text{of the Maclaurin series for } e^x \\ 2 : \text{Taylor polynomial} \end{cases}$