

2017

AP Calculus BC

Scoring Guidelines

2017 SCORING GUIDELINES

Question 1

	1 : units in parts (a), (c), and (d)
(a) $\text{Volume} = \int_0^{10} A(h) \, dh$ $\approx (2 - 0) \cdot A(0) + (5 - 2) \cdot A(2) + (10 - 5) \cdot A(5)$ $= 2 \cdot 50.3 + 3 \cdot 14.4 + 5 \cdot 6.5$ $= 176.3 \text{ cubic feet}$	2 : $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{approximation} \end{cases}$
(b) The approximation in part (a) is an overestimate because a left Riemann sum is used and A is decreasing.	1 : overestimate with reason
(c) $\int_0^{10} f(h) \, dh = 101.325338$ The volume is 101.325 cubic feet.	2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$
(d) Using the model, $V(h) = \int_0^h f(x) \, dx$. $\left. \frac{dV}{dt} \right _{h=5} = \left[\frac{dV}{dh} \cdot \frac{dh}{dt} \right]_{h=5}$ $= \left[f(h) \cdot \frac{dh}{dt} \right]_{h=5}$ $= f(5) \cdot 0.26 = 1.694419$ When $h = 5$, the volume of water is changing at a rate of 1.694 cubic feet per minute.	3 : $\begin{cases} 2 : \frac{dV}{dt} \\ 1 : \text{answer} \end{cases}$

2017 SCORING GUIDELINES

Question 2

$$(a) \frac{1}{2} \int_0^{\pi/2} (f(\theta))^2 d\theta = 0.648414$$

The area of R is 0.648.

$$(b) \int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \frac{1}{2} \int_0^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

— OR —

$$\int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \int_k^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

$$(c) w(\theta) = g(\theta) - f(\theta)$$

$$w_A = \frac{\int_0^{\pi/2} w(\theta) d\theta}{\frac{\pi}{2} - 0} = 0.485446$$

The average value of $w(\theta)$ on the interval $\left[0, \frac{\pi}{2}\right]$ is 0.485.

$$(d) w(\theta) = w_A \text{ for } 0 \leq \theta \leq \frac{\pi}{2} \Rightarrow \theta = 0.517688$$

$$w(\theta) = w_A \text{ at } \theta = 0.518 \text{ (or } 0.517).$$

$$w'(0.518) < 0 \Rightarrow w(\theta) \text{ is decreasing at } \theta = 0.518.$$

$$2: \begin{cases} 1: \text{integral} \\ 1: \text{answer} \end{cases}$$

$$2: \begin{cases} 1: \text{integral expression} \\ \quad \text{for one region} \\ 1: \text{equation} \end{cases}$$

$$3: \begin{cases} 1: w(\theta) \\ 1: \text{integral} \\ 1: \text{average value} \end{cases}$$

$$2: \begin{cases} 1: \text{solves } w(\theta) = w_A \\ 1: \text{answer with reason} \end{cases}$$

2017 SCORING GUIDELINES

Question 3

$$(a) f(-6) = f(-2) + \int_{-2}^{-6} f'(x) dx = 7 - \int_{-6}^{-2} f'(x) dx = 7 - 4 = 3$$

$$f(5) = f(-2) + \int_{-2}^5 f'(x) dx = 7 - 2\pi + 3 = 10 - 2\pi$$

$$(b) f'(x) > 0 \text{ on the intervals } [-6, -2) \text{ and } (2, 5).$$

Therefore, f is increasing on the intervals $[-6, -2]$ and $[2, 5]$.

$$(c) \text{The absolute minimum will occur at a critical point where } f'(x) = 0 \text{ or at an endpoint.}$$

$$f'(x) = 0 \Rightarrow x = -2, x = 2$$

x	$f(x)$
-6	3
-2	7
2	$7 - 2\pi$
5	$10 - 2\pi$

The absolute minimum value is $f(2) = 7 - 2\pi$.

$$(d) f''(-5) = \frac{2 - 0}{-6 - (-2)} = -\frac{1}{2}$$

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} = 2 \text{ and } \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3} = -1$$

$f''(3)$ does not exist because

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} \neq \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3}.$$

$$3: \begin{cases} 1: \text{uses initial condition} \\ 1: f(-6) \\ 1: f(5) \end{cases}$$

$$2: \text{answer with justification}$$

$$2: \begin{cases} 1: \text{considers } x = 2 \\ 1: \text{answer with justification} \end{cases}$$

$$2: \begin{cases} 1: f''(-5) \\ 1: f''(3) \text{ does not exist,} \\ \quad \text{with explanation} \end{cases}$$

2017 SCORING GUIDELINES

Question 4

(a) $H'(0) = -\frac{1}{4}(91 - 27) = -16$
 $H(0) = 91$

An equation for the tangent line is $y = 91 - 16t$.

The internal temperature of the potato at time $t = 3$ minutes is approximately $91 - 16 \cdot 3 = 43$ degrees Celsius.

(b) $\frac{d^2H}{dt^2} = -\frac{1}{4} \frac{dH}{dt} = \left(-\frac{1}{4}\right)\left(-\frac{1}{4}\right)(H - 27) = \frac{1}{16}(H - 27)$

$$H > 27 \text{ for } t > 0 \Rightarrow \frac{d^2H}{dt^2} = \frac{1}{16}(H - 27) > 0 \text{ for } t > 0$$

Therefore, the graph of H is concave up for $t > 0$. Thus, the answer in part (a) is an underestimate.

(c) $\frac{dG}{(G - 27)^{2/3}} = -dt$
 $\int \frac{dG}{(G - 27)^{2/3}} = \int (-1) dt$
 $3(G - 27)^{1/3} = -t + C$
 $3(91 - 27)^{1/3} = 0 + C \Rightarrow C = 12$
 $3(G - 27)^{1/3} = 12 - t$
 $G(t) = 27 + \left(\frac{12 - t}{3}\right)^3 \text{ for } 0 \leq t < 10$

The internal temperature of the potato at time $t = 3$ minutes is $27 + \left(\frac{12 - 3}{3}\right)^3 = 54$ degrees Celsius.

3 : $\begin{cases} 1 : \text{slope} \\ 1 : \text{tangent line} \\ 1 : \text{approximation} \end{cases}$

1 : underestimate with reason

5 : $\begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration and uses initial condition} \\ 1 : \text{equation involving } G \text{ and } t \\ 1 : G(t) \text{ and } G(3) \end{cases}$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

2017 SCORING GUIDELINES

Question 5

(a) $f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2}$

$$f'(3) = \frac{(-3)(5)}{(18 - 21 + 5)^2} = -\frac{15}{4}$$

(b) $f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2} = 0 \Rightarrow x = \frac{7}{4}$

The only critical point in the interval $1 < x < 2.5$ has x -coordinate $\frac{7}{4}$.

f' changes sign from positive to negative at $x = \frac{7}{4}$.

Therefore, f has a relative maximum at $x = \frac{7}{4}$.

(c) $\int_5^\infty f(x) dx = \lim_{b \rightarrow \infty} \int_5^b \frac{3}{2x^2 - 7x + 5} dx = \lim_{b \rightarrow \infty} \int_5^b \left(\frac{2}{2x - 5} - \frac{1}{x - 1} \right) dx$
 $= \lim_{b \rightarrow \infty} \left[\ln(2x - 5) - \ln(x - 1) \right]_5^b = \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2x - 5}{x - 1} \right) \right]_5^b$
 $= \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2b - 5}{b - 1} \right) - \ln \left(\frac{5}{4} \right) \right] = \ln 2 - \ln \left(\frac{5}{4} \right) = \ln \left(\frac{8}{5} \right)$

(d) f is continuous, positive, and decreasing on $[5, \infty)$.

The series converges by the integral test since $\int_5^\infty \frac{3}{2x^2 - 7x + 5} dx$ converges.

— OR —

$$\frac{3}{2n^2 - 7n + 5} > 0 \text{ and } \frac{1}{n^2} > 0 \text{ for } n \geq 5.$$

Since $\lim_{n \rightarrow \infty} \frac{\frac{3}{2n^2 - 7n + 5}}{\frac{1}{n^2}} = \frac{3}{2}$ and the series $\sum_{n=5}^\infty \frac{1}{n^2}$ converges,

the series $\sum_{n=5}^\infty \frac{3}{2n^2 - 7n + 5}$ converges by the limit comparison test.

2 : $f'(3)$

2 : $\begin{cases} 1 : x\text{-coordinate} \\ 1 : \text{relative maximum with justification} \end{cases}$

3 : $\begin{cases} 1 : \text{antiderivative} \\ 1 : \text{limit expression} \\ 1 : \text{answer} \end{cases}$

2 : answer with conditions

2017 SCORING GUIDELINES

Question 6

(a) $f(0) = 0$

$f'(0) = 1$

$f''(0) = -1(1) = -1$

$f'''(0) = -2(-1) = 2$

$f^{(4)}(0) = -3(2) = -6$

The first four nonzero terms are

$$0 + 1x + \frac{-1}{2!}x^2 + \frac{2}{3!}x^3 + \frac{-6}{4!}x^4 = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4}.$$

The general term is $\frac{(-1)^{n+1}x^n}{n}$.

(b) For $x = 1$, the Maclaurin series becomes $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n}$.

The series does not converge absolutely because the harmonic series diverges.

The series alternates with terms that decrease in magnitude to 0, and therefore the series converges conditionally.

(c)
$$\begin{aligned} \int_0^x f(t) dt &= \int_0^x \left(t - \frac{t^2}{2} + \frac{t^3}{3} - \frac{t^4}{4} + \cdots + \frac{(-1)^{n+1}t^n}{n} + \cdots \right) dt \\ &= \left[\frac{t^2}{2} - \frac{t^3}{3 \cdot 2} + \frac{t^4}{4 \cdot 3} - \frac{t^5}{5 \cdot 4} + \cdots + \frac{(-1)^{n+1}t^{n+1}}{(n+1)n} + \cdots \right]_{t=0}^{t=x} \\ &= \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{12} - \frac{x^5}{20} + \cdots + \frac{(-1)^{n+1}x^{n+1}}{(n+1)n} + \cdots \end{aligned}$$

(d) The terms alternate in sign and decrease in magnitude to 0. By the alternating series error bound, the error $\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right|$ is boundedby the magnitude of the first unused term, $\left| -\frac{(1/2)^5}{20} \right|$.

Thus, $\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right| \leq \left| -\frac{(1/2)^5}{20} \right| = \frac{1}{32 \cdot 20} < \frac{1}{500}.$

$$3 : \begin{cases} 1 : f''(0), f'''(0), \text{ and } f^{(4)}(0) \\ 1 : \text{verify terms} \\ 1 : \text{general term} \end{cases}$$

2 : converges conditionally
with reason

$$3 : \begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \\ 1 : \text{general term} \end{cases}$$

1 : error bound