


Part B (AB or BC): Graphing calculator not allowed**Question 4****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Graph of f'

The function f is defined on the closed interval $[-2, 8]$ and satisfies $f(2) = 1$. The graph of f' , the derivative of f , consists of two line segments and a semicircle, as shown in the figure.

Model Solution**Scoring**

- (a) Does f have a relative minimum, a relative maximum, or neither at $x = 6$? Give a reason for your answer.

$$f'(x) > 0 \text{ on } (2, 6) \text{ and } f'(x) > 0 \text{ on } (6, 8).$$

$f'(x)$ does not change sign at $x = 6$, so there is neither a relative maximum nor a relative minimum at this location.

Answer with reason

1 point**Scoring notes:**

- A response that declares $f'(x)$ does not change sign at $x = 6$, so neither, is sufficient to earn the point.
- A response does not have to present intervals on which $f'(x)$ is positive or negative, but if any are given, they must be correct. Any presented intervals may include none, one, or both endpoints.
- A response that declares $f'(x) > 0$ before and after $x = 6$ does not earn the point.

Total for part (a)**1 point**

- (b) On what open intervals, if any, is the graph of f concave down? Give a reason for your answer.

The graph of f is concave down on $(-2, 0)$ and $(4, 6)$ because f' is decreasing on these intervals.

Intervals

1 point

Reason

1 point**Scoring notes:**

- The first point is earned only by an answer of $(-2, 0)$ and $(4, 6)$, or these intervals including one or both endpoints.
- A response must earn the first point to be eligible for second point.
- To earn the second point a response must correctly discuss the behavior of f' or the slopes of f' .
- Special case: A response that presents exactly one of the two correct intervals with a correct reason earns 1 out of 2 points.

Total for part (b)**2 points**

- (c) Find the value of $\lim_{x \rightarrow 2} \frac{6f(x) - 3x}{x^2 - 5x + 6}$, or show that it does not exist. Justify your answer.

Because f is differentiable at $x = 2$, f is continuous at $x = 2$, so $\lim_{x \rightarrow 2} f(x) = f(2) = 1$.

$$\lim_{x \rightarrow 2} (6f(x) - 3x) = 6 \cdot 1 - 3 \cdot 2 = 0$$

$$\lim_{x \rightarrow 2} (x^2 - 5x + 6) = 0$$

Limits of numerator and denominator

1 point

Because $\lim_{x \rightarrow 2} \frac{6f(x) - 3x}{x^2 - 5x + 6}$ is of indeterminate form $\frac{0}{0}$,

L'Hospital's Rule can be applied.

Uses L'Hospital's Rule

1 point

Answer

1 point

Using L'Hospital's Rule,

$$\lim_{x \rightarrow 2} \frac{6f(x) - 3x}{x^2 - 5x + 6} = \lim_{x \rightarrow 2} \frac{6f'(x) - 3}{2x - 5} = \frac{6 \cdot 0 - 3}{2 \cdot 2 - 5} = 3.$$

Scoring notes:

- The first point is earned by the presentation of two separate limits for the numerator and denominator.
- A response that presents a limit explicitly equal to $\frac{0}{0}$ does not earn the first point.
- The second point is earned by applying L'Hospital's Rule, that is, by presenting at least one correct derivative in the limit of a ratio of derivatives.
- The third point is earned for the correct answer with supporting work.

Total for part (c)**3 points**

- (d) Find the absolute minimum value of f on the closed interval $[-2, 8]$. Justify your answer.

$$f'(x) = 0 \Rightarrow x = -1, x = 2, x = 6$$

Considers $f'(x) = 0$ **1 point**

The function f is continuous on $[-2, 8]$, so the candidates for the location of an absolute minimum for f are $x = -2$, $x = -1$, $x = 2$, $x = 6$, and $x = 8$.

Justification **1 point**

Answer **1 point**

x	$f(x)$
-2	3
-1	4
2	1
6	$7 - \pi$
8	$11 - 2\pi$

The absolute minimum value of f is $f(2) = 1$.

Scoring notes:

- To earn the first point a response must state $f' = 0$ or equivalent. Listing the zeros of f' is not sufficient.
- A response that presents any error in evaluating f at any critical point or endpoint will not earn the justification point.
- A response need not present the value of $f(-1)$ provided $x = -1$ is eliminated because it is the location of a local maximum. A response need not present the value of $f(6)$ provided $x = 6$ is eliminated by reference to part (a) or eliminated through analysis.
- A response need not present the value of $f(8)$ provided there is a presentation that argues $f'(x) \geq 0$ for $x > 2$ and, therefore, $f(8) > f(2)$.
- A response that does not consider both endpoints does not earn the justification point.
- The answer point is earned only for indicating that the minimum value is 1. It is not earned for noting that the minimum occurs at $x = 2$.

Total for part (d) 3 points

Total for question 4 9 points

AP[®] Calculus BC 2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2016 SCORING GUIDELINES

Question 1

t (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.

- (a) Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.
- (b) Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
- (c) Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
- (d) For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

(a) $R'(2) \approx \frac{R(3) - R(1)}{3 - 1} = \frac{950 - 1190}{3 - 1} = -120$ liters/hr²

2 : $\begin{cases} 1 : \text{estimate} \\ 1 : \text{units} \end{cases}$

(b) The total amount of water removed is given by $\int_0^8 R(t) dt$.

$$\begin{aligned} \int_0^8 R(t) dt &\approx 1 \cdot R(0) + 2 \cdot R(1) + 3 \cdot R(3) + 2 \cdot R(6) \\ &= 1(1340) + 2(1190) + 3(950) + 2(740) \\ &= 8050 \text{ liters} \end{aligned}$$

3 : $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{estimate} \\ 1 : \text{overestimate with reason} \end{cases}$

This is an overestimate since R is a decreasing function.

(c) Total $\approx 50000 + \int_0^8 W(t) dt - 8050$
 $= 50000 + 7836.195325 - 8050 \approx 49786$ liters

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{estimate} \end{cases}$

(d) $W(0) - R(0) > 0$, $W(8) - R(8) < 0$, and $W(t) - R(t)$ is continuous.


2 : $\begin{cases} 1 : \text{considers } W(t) - R(t) \\ 1 : \text{answer with explanation} \end{cases}$

Therefore, the Intermediate Value Theorem guarantees at least one time t , $0 < t < 8$, for which $W(t) - R(t) = 0$, or $W(t) = R(t)$.

For this value of t , the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank.

2016 SCORING GUIDELINES

Question 2



At time t , the position of a particle moving in the xy -plane is given by the parametric functions $(x(t), y(t))$, where $\frac{dx}{dt} = t^2 + \sin(3t^2)$. The graph of y , consisting of three line segments, is shown in the figure above. At $t = 0$, the particle is at position $(5, 1)$.

- (a) Find the position of the particle at $t = 3$.
- (b) Find the slope of the line tangent to the path of the particle at $t = 3$.
- (c) Find the speed of the particle at $t = 3$.
- (d) Find the total distance traveled by the particle from $t = 0$ to $t = 2$.

(a) $x(3) = x(0) + \int_0^3 x'(t) dt = 5 + 9.377035 = 14.377$

$$y(3) = -\frac{1}{2}$$

The position of the particle at $t = 3$ is $(14.377, -0.5)$.

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

(b) Slope $= \frac{y'(3)}{x'(3)} = \frac{0.5}{9.956376} = 0.05$

1 : slope

(c) Speed $= \sqrt{(x'(3))^2 + (y'(3))^2} = 9.969$ (or 9.968)

2 : $\begin{cases} 1 : \text{expression for speed} \\ 1 : \text{answer} \end{cases}$

(d) Distance $= \int_0^2 \sqrt{(x'(t))^2 + (y'(t))^2} dt$
 $= \int_0^1 \sqrt{(x'(t))^2 + (-2)^2} dt + \int_1^2 \sqrt{(x'(t))^2 + 0^2} dt$
 $= 2.237871 + 2.112003 = 4.350$ (or 4.349)

3 : $\begin{cases} 1 : \text{expression for distance} \\ 1 : \text{integrals} \\ 1 : \text{answer} \end{cases}$


2016 SCORING GUIDELINES

Question 3

The figure above shows the graph of the piecewise-linear function f . For $-4 \leq x \leq 12$, the function g is defined by

$$g(x) = \int_2^x f(t) dt.$$

- (a) Does g have a relative minimum, a relative maximum, or neither at $x = 10$? Justify your answer.
- (b) Does the graph of g have a point of inflection at $x = 4$? Justify your answer.
- (c) Find the absolute minimum value and the absolute maximum value of g on the interval $-4 \leq x \leq 12$. Justify your answers.
- (d) For $-4 \leq x \leq 12$, find all intervals for which $g(x) \leq 0$.

Graph of f

- (a) The function g has neither a relative minimum nor a relative maximum at $x = 10$ since $g'(x) = f(x)$ and $f(x) \leq 0$ for $8 \leq x \leq 12$.
- (b) The graph of g has a point of inflection at $x = 4$ since $g'(x) = f(x)$ is increasing for $2 \leq x \leq 4$ and decreasing for $4 \leq x \leq 8$.
- (c) $g'(x) = f(x)$ changes sign only at $x = -2$ and $x = 6$.

x	$g(x)$
-4	-4
-2	-8
6	8
12	-4

On the interval $-4 \leq x \leq 12$, the absolute minimum value is $g(-2) = -8$ and the absolute maximum value is $g(6) = 8$.

- (d) $g(x) \leq 0$ for $-4 \leq x \leq 2$ and $10 \leq x \leq 12$.

1 : $g'(x) = f(x)$ in (a), (b), (c), or (d)

1 : answer with justification

1 : answer with justification

4 : $\begin{cases} 1 : \text{considers } x = -2 \text{ and } x = 6 \\ \quad \text{as candidates} \\ 1 : \text{considers } x = -4 \text{ and } x = 12 \\ 2 : \text{answers with justification} \end{cases}$

2 : intervals

2016 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = x^2 - \frac{1}{2}y$.

- (a) Find $\frac{d^2y}{dx^2}$ in terms of x and y .
- (b) Let $y = f(x)$ be the particular solution to the given differential equation whose graph passes through the point $(-2, 8)$. Does the graph of f have a relative minimum, a relative maximum, or neither at the point $(-2, 8)$? Justify your answer.
- (c) Let $y = g(x)$ be the particular solution to the given differential equation with $g(-1) = 2$. Find $\lim_{x \rightarrow -1} \left(\frac{g(x) - 2}{3(x+1)^2} \right)$. Show the work that leads to your answer.
- (d) Let $y = h(x)$ be the particular solution to the given differential equation with $h(0) = 2$. Use Euler's method, starting at $x = 0$ with two steps of equal size, to approximate $h(1)$.

(a) $\frac{d^2y}{dx^2} = 2x - \frac{1}{2} \frac{dy}{dx} = 2x - \frac{1}{2} \left(x^2 - \frac{1}{2}y \right)$

2 : $\frac{d^2y}{dx^2}$ in terms of x and y

(b) $\left. \frac{dy}{dx} \right|_{(x,y)=(-2,8)} = (-2)^2 - \frac{1}{2} \cdot 8 = 0$
 $\left. \frac{d^2y}{dx^2} \right|_{(x,y)=(-2,8)} = 2(-2) - \frac{1}{2} \left((-2)^2 - \frac{1}{2} \cdot 8 \right) = -4 < 0$

Thus, the graph of f has a relative maximum at the point $(-2, 8)$.

2 : conclusion with justification

(c) $\lim_{x \rightarrow -1} (g(x) - 2) = 0$ and $\lim_{x \rightarrow -1} 3(x+1)^2 = 0$

Using L'Hospital's Rule,

$$\lim_{x \rightarrow -1} \left(\frac{g(x) - 2}{3(x+1)^2} \right) = \lim_{x \rightarrow -1} \left(\frac{g'(x)}{6(x+1)} \right)$$

$$\lim_{x \rightarrow -1} g'(x) = 0 \text{ and } \lim_{x \rightarrow -1} 6(x+1) = 0$$

Using L'Hospital's Rule,

$$\lim_{x \rightarrow -1} \left(\frac{g'(x)}{6(x+1)} \right) = \lim_{x \rightarrow -1} \left(\frac{g''(x)}{6} \right) = \frac{-2}{6} = -\frac{1}{3}$$


3 : $\begin{cases} 2 : \text{L'Hospital's Rule} \\ 1 : \text{answer} \end{cases}$

(d) $h\left(\frac{1}{2}\right) \approx h(0) + h'(0) \cdot \frac{1}{2} = 2 + (-1) \cdot \frac{1}{2} = \frac{3}{2}$
 $h(1) \approx h\left(\frac{1}{2}\right) + h'\left(\frac{1}{2}\right) \cdot \frac{1}{2} \approx \frac{3}{2} + \left(-\frac{1}{2}\right) \cdot \frac{1}{2} = \frac{5}{4}$

2 : $\begin{cases} 1 : \text{Euler's method} \\ 1 : \text{approximation} \end{cases}$

2016 SCORING GUIDELINES

Question 5



The inside of a funnel of height 10 inches has circular cross sections, as shown in the figure above. At height h , the radius of the funnel is given by $r = \frac{1}{20}(3 + h^2)$, where $0 \leq h \leq 10$. The units of r and h are inches.

- Find the average value of the radius of the funnel.
- Find the volume of the funnel.
- The funnel contains liquid that is draining from the bottom. At the instant when the height of the liquid is $h = 3$ inches, the radius of the surface of the liquid is decreasing at a rate of $\frac{1}{5}$ inch per second. At this instant, what is the rate of change of the height of the liquid with respect to time?

$$\begin{aligned} \text{(a) Average radius} &= \frac{1}{10} \int_0^{10} \frac{1}{20}(3 + h^2) dh = \frac{1}{200} \left[3h + \frac{h^3}{3} \right]_0^{10} \\ &= \frac{1}{200} \left(\left(30 + \frac{1000}{3} \right) - 0 \right) = \frac{109}{60} \text{ in} \end{aligned}$$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b) Volume} &= \pi \int_0^{10} \left(\frac{1}{20}(3 + h^2) \right)^2 dh = \frac{\pi}{400} \int_0^{10} (9 + 6h^2 + h^4) dh \\ &= \frac{\pi}{400} \left[9h + 2h^3 + \frac{h^5}{5} \right]_0^{10} \\ &= \frac{\pi}{400} \left(\left(90 + 2000 + \frac{100000}{5} \right) - 0 \right) = \frac{2209\pi}{40} \text{ in}^3 \end{aligned}$$

3 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(c) } \frac{dr}{dt} &= \frac{1}{20}(2h) \frac{dh}{dt} \\ -\frac{1}{5} &= \frac{3}{10} \frac{dh}{dt} \\ \frac{dh}{dt} &= -\frac{1}{5} \cdot \frac{10}{3} = -\frac{2}{3} \text{ in/sec} \end{aligned}$$

3 : $\begin{cases} 2 : \text{chain rule} \\ 1 : \text{answer} \end{cases}$

2016 SCORING GUIDELINES

Question 6

The function f has a Taylor series about $x = 1$ that converges to $f(x)$ for all x in the interval of convergence.

It is known that $f(1) = 1$, $f'(1) = -\frac{1}{2}$, and the n th derivative of f at $x = 1$ is given by

$$f^{(n)}(1) = (-1)^n \frac{(n-1)!}{2^n} \text{ for } n \geq 2.$$

- Write the first four nonzero terms and the general term of the Taylor series for f about $x = 1$.
- The Taylor series for f about $x = 1$ has a radius of convergence of 2. Find the interval of convergence. Show the work that leads to your answer.
- The Taylor series for f about $x = 1$ can be used to represent $f(1.2)$ as an alternating series. Use the first three nonzero terms of the alternating series to approximate $f(1.2)$.
- Show that the approximation found in part (c) is within 0.001 of the exact value of $f(1.2)$.

$$\begin{aligned} \text{(a) } f(1) &= 1, \quad f'(1) = -\frac{1}{2}, \quad f''(1) = \frac{1}{2^2}, \quad f'''(1) = -\frac{2}{2^3} \\ f(x) &= 1 - \frac{1}{2}(x-1) + \frac{1}{2^2} \cdot \frac{1}{2}(x-1)^2 - \frac{1}{2^3} \cdot \frac{1}{3}(x-1)^3 + \cdots \\ &\quad + \frac{(-1)^n}{2^n \cdot n}(x-1)^n + \cdots \end{aligned}$$

4 : $\begin{cases} 1 : \text{first two terms} \\ 1 : \text{third term} \\ 1 : \text{fourth term} \\ 1 : \text{general term} \end{cases}$

- $R = 2$. The series converges on the interval $(-1, 3)$.

When $x = -1$, the series is $1 + 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots$.

Since the harmonic series diverges, this series diverges.

When $x = 3$, the series is $1 - 1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \cdots$.

Since the alternating harmonic series converges, this series converges.

Therefore, the interval of convergence is $-1 < x \leq 3$.

$$\text{(c) } f(1.2) \approx 1 - \frac{1}{2}(0.2) + \frac{1}{8}(0.2)^2 = 1 - 0.1 + 0.005 = 0.905$$

1 : approximation

- The series for $f(1.2)$ alternates with terms that decrease in magnitude to 0.

2 : $\begin{cases} 1 : \text{error form} \\ 1 : \text{analysis} \end{cases}$

$$|f(1.2) - T_2(1.2)| \leq \left| \frac{-1}{2^3 \cdot 3}(0.2)^3 \right| = \frac{1}{3000} \leq 0.001$$