

**Part B (BC): Graphing calculator not allowed****Question 5****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

|         |   |       |        |
|---------|---|-------|--------|
| $x$     | 0 | $\pi$ | $2\pi$ |
| $f'(x)$ | 5 | 6     | 0      |

The function  $f$  is twice differentiable for all  $x$  with  $f(0) = 0$ . Values of  $f'$ , the derivative of  $f$ , are given in the table for selected values of  $x$ .

**Model Solution****Scoring**

- (a) For  $x \geq 0$ , the function  $h$  is defined by  $h(x) = \int_0^x \sqrt{1 + (f'(t))^2} dt$ . Find the value of  $h'(\pi)$ . Show the work that leads to your answer.

|                                                                 |                                 |                |
|-----------------------------------------------------------------|---------------------------------|----------------|
| $h'(x) = \sqrt{1 + (f'(x))^2}$                                  | Fundamental Theorem of Calculus | <b>1 point</b> |
| $h'(\pi) = \sqrt{1 + (f'(\pi))^2} = \sqrt{1 + 6^2} = \sqrt{37}$ | Answer                          | <b>1 point</b> |

**Scoring notes:**

- A response of  $\sqrt{1 + (f'(\pi))^2}$  earns the first point.
- A response of  $\sqrt{1 + 6^2}$  alone earns both points.
- A response such as  $h'(x) = \sqrt{1 + (f'(x))^2} = \sqrt{37}$ , that equates a variable expression to a numeric value, earns at most 1 of the 2 points.
- A response that equates  $h'(x)$  or  $h'(\pi)$  to a derivative of a constant, such as

$$h'(x) = \frac{d}{dx} \int_0^x \sqrt{1 + (f'(t))^2} dt, \text{ earns at most 1 of the 2 points.}$$

**Total for part (a) 2 points**

- (b) What information does  $\int_0^\pi \sqrt{1 + (f'(x))^2} dx$  provide about the graph of  $f$ ?

|                                                                                            |                     |                |
|--------------------------------------------------------------------------------------------|---------------------|----------------|
| $\int_0^\pi \sqrt{1 + (f'(x))^2} dx$ is the arc length of the graph of $f$ on $[0, \pi]$ . | Arc length of $f$   | <b>1 point</b> |
|                                                                                            | Interval $[0, \pi]$ | <b>1 point</b> |

**Scoring notes:**

- A response of “arc length” or “length” earns the first point. Such a response does not need to reference  $f$ . However, if the response references a different function, the response does not earn the first point and is eligible to earn the second point.
- A response referring to distance explicitly connected to the graph or  $f$  (or equivalent) earns the first point. For example, a response of “distance along the curve” or “distance traveled by a particle moving along  $f$ ” earns the first point and is eligible to earn the second point.
- A response referring to distance that is not explicitly connected to the graph of  $f$  does not earn the first point but is eligible to earn the second point. For example, a response of “distance” or “distance traveled” does not earn the first point but is eligible to earn the second point.
- To earn the second point a response must connect the interval  $[0, \pi]$  to arc length, length, or distance.

**Total for part (b) 2 points**

- (c) Use Euler's method, starting at  $x = 0$  with two steps of equal size, to approximate  $f(2\pi)$ . Show the computations that lead to your answer.

|                                                     |                |         |
|-----------------------------------------------------|----------------|---------|
| $f(\pi) \approx f(0) + \pi f'(0) = 0 + 5\pi = 5\pi$ | Euler's method | 1 point |
| $f(2\pi) \approx f(\pi) + \pi f'(\pi)$              |                |         |
| $\approx 5\pi + 6\pi = 11\pi$                       | Answer         | 1 point |

**Scoring notes:**

- To earn the first point a response must demonstrate two Euler's steps, with use of the correct expression for  $\frac{dy}{dx}$ , and at most one error. If there is an error, the second point is not earned.
- In order to earn the first point, a response that presents a single error in computing the approximation of  $f(\pi)$  must import the incorrect value in computing the approximation of  $f(2\pi)$ .
- The two Euler's steps may be explicit expressions or may be presented in a table. For example:

| $x$    | $y$     | $\frac{dy}{dx} \cdot \Delta x$ (or $\frac{dy}{dx} \cdot \pi$ ) |
|--------|---------|----------------------------------------------------------------|
| 0      | 0       | $5\pi$                                                         |
| $\pi$  | $5\pi$  | $6\pi$                                                         |
| $2\pi$ | $11\pi$ |                                                                |

- In the presence of a correct answer, a table does not need to be labeled in order to earn both points. In the presence of no answer or an incorrect answer, such a table must be correctly labeled in order to earn the first point.
- Both points are earned for  $5\pi + 6\pi$ .
- The response may report the final answer as  $(2\pi, 11\pi)$ .

**Total for part (c) 2 points**

- (d) Find  $\int (t+5)\cos\left(\frac{t}{4}\right) dt$ . Show the work that leads to your answer.

|                                                                                                                          |                  |         |
|--------------------------------------------------------------------------------------------------------------------------|------------------|---------|
| $u = t + 5 \quad dv = \cos\left(\frac{t}{4}\right) dt$<br>$du = dt \quad v = 4\sin\left(\frac{t}{4}\right)$              | $u$ and $dv$     | 1 point |
| $\int (t+5)\cos\left(\frac{t}{4}\right) dt = 4(t+5)\sin\left(\frac{t}{4}\right) - \int 4\sin\left(\frac{t}{4}\right) dt$ | $uv - \int v du$ | 1 point |
| $= 4(t+5)\sin\left(\frac{t}{4}\right) + 16\cos\left(\frac{t}{4}\right) + C$                                              | Answer           | 1 point |

**Scoring notes:**

- The first and second points are earned with an implied  $u$  and  $dv$  in the presence of  $4(t+5)\sin\left(\frac{t}{4}\right) - \int 4\sin\left(\frac{t}{4}\right) dt$  or a mathematically equivalent expression.
- The tabular method may be used to show integration by parts. In this case, the first point is earned by columns (labeled or unlabeled) that begin with  $t+5$  and  $\cos\left(\frac{t}{4}\right)$ . The second point is earned for  $4(t+5)\sin\left(\frac{t}{4}\right) - \int 4\sin\left(\frac{t}{4}\right) dt$  or a mathematically equivalent expression.
- The third point is earned only for an expression mathematically equivalent to  $4(t+5)\sin\left(\frac{t}{4}\right) + 16\cos\left(\frac{t}{4}\right) + C$  (such as  $4t\sin\left(\frac{t}{4}\right) + 20\sin\left(\frac{t}{4}\right) + 16\cos\left(\frac{t}{4}\right) + C$ ) in the presence of correct supporting work.
- To earn the third point a response must have a final answer that includes a constant of integration.
- Alternate solution:

$$\int (t+5)\cos\left(\frac{t}{4}\right) dt = \int t\cos\left(\frac{t}{4}\right) dt + \int 5\cos\left(\frac{t}{4}\right) dt$$

$$u = t \quad dv = \cos\left(\frac{t}{4}\right) dt$$

$$du = dt \quad v = 4\sin\left(\frac{t}{4}\right)$$

$$\begin{aligned} \int t\cos\left(\frac{t}{4}\right) dt + \int 5\cos\left(\frac{t}{4}\right) dt &= 4t\sin\left(\frac{t}{4}\right) - \int 4\sin\left(\frac{t}{4}\right) dt + \int 5\cos\left(\frac{t}{4}\right) dt \\ &= 4t\sin\left(\frac{t}{4}\right) + 16\cos\left(\frac{t}{4}\right) + 20\sin\left(\frac{t}{4}\right) + C \end{aligned}$$


- A response can earn the first and second points for correctly applying integration by parts to  $\int t\cos\left(\frac{t}{4}\right) dt$ . The tabular method may be used to show integration by parts. The third point is earned for the correct answer.

**Total for part (d) 3 points****Total for question 5 9 points**

**Part B (BC): Graphing calculator not allowed****Question 5****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.



The graphs of the functions  $f$  and  $g$  are shown in the figure for  $0 \leq x \leq 3$ . It is known that  $g(x) = \frac{12}{3+x}$  for  $x \geq 0$ . The twice-differentiable function  $f$ , which is not explicitly given, satisfies  $f(3) = 2$  and  $\int_0^3 f(x) dx = 10$ .

**Model Solution****Scoring**

(a) Find the area of the shaded region enclosed by the graphs of  $f$  and  $g$ .

|                                                                                 |                          |                |
|---------------------------------------------------------------------------------|--------------------------|----------------|
| $\text{Area} = \int_0^3 (f(x) - g(x)) dx = \int_0^3 f(x) dx - \int_0^3 g(x) dx$ | Integrand                | <b>1 point</b> |
| $= 10 - \int_0^3 \frac{12}{3+x} dx = 10 - 12[\ln 3+x ]_0^3$                     | Antiderivative of $g(x)$ | <b>1 point</b> |
| $= 10 - 12(\ln 6 - \ln 3) = 10 - 12(\ln 2)$                                     | Answer                   | <b>1 point</b> |

**Scoring notes:**

- The first point is earned for any of the integrands  $f(x) - g(x)$ ,  $g(x) - f(x)$ ,  $|f(x) - g(x)|$ , or  $|g(x) - f(x)|$  in any definite integral. If the limits are incorrect, the response does not earn the third point.

- The first point is earned with an implied integrand for  $f$  and explicit integrand for  $g$ , such as

$$10 - \int_0^3 g(x) dx.$$

- The second point is earned for finding  $a \int \frac{dx}{3+x} = a \cdot \ln|3+x|$  or  $a \cdot \ln(3+x)$ .
- A response is eligible for the third point only if it has earned the first 2 points. The third point is earned only for the correct answer. The answer does not need to be simplified; however, if simplification is attempted, it must be correct.
- A response is not eligible for the third point with incorrect limits of integration for  $u$ -substitution, for example,  $\int_0^3 \frac{12}{3+x} dx = \int_0^3 \frac{12}{u} du = 12[\ln(x+3)]_0^3$ .
- A response with incorrect communication, such as “Area =  $\int_0^3 (g(x) - f(x)) dx = 10 - 12(\ln 2)$ ,” does not earn the third point. However, a response of “ $\int_0^3 (g(x) - f(x)) dx = 12(\ln 2) - 10$ , so the area is  $10 - 12(\ln 2)$ ” earns all 3 points.

**Total for part (a) 3 points**

(b) Evaluate the improper integral  $\int_0^\infty (g(x))^2 dx$ , or show that the integral diverges.

|                                                                                           |                |                |
|-------------------------------------------------------------------------------------------|----------------|----------------|
| $\int_0^\infty (g(x))^2 dx = \lim_{b \rightarrow \infty} \int_0^b \frac{144}{(3+x)^2} dx$ | Limit notation | <b>1 point</b> |
| $= \lim_{b \rightarrow \infty} \left( -\frac{144}{(3+x)} \Big _0^b \right)$               | Antiderivative | <b>1 point</b> |
| $= \lim_{b \rightarrow \infty} \left( -\frac{144}{3+b} + \frac{144}{3} \right) = 48$      | Answer         | <b>1 point</b> |

**Scoring notes:**

- To earn the first point a response must correctly use limit notation throughout the problem and not include arithmetic with infinity, for example,  $\left[ -\frac{144}{3+x} \right]_0^\infty$  or  $-\frac{144}{3+\infty} + 48$ .
- The second point can be earned by finding an antiderivative of the form  $-\frac{a}{(3+x)}$  for  $a > 0$ , from an indefinite or improper integral, with or without correct limit notation. If  $a \neq 144$ , the response does not earn the third point.
- The third point is earned only for an answer of 48 (or equivalent).
- A response is not eligible for the third point with incorrect limits of integration for  $u$ -substitution, for example,  $\lim_{b \rightarrow \infty} \int_0^b \frac{144}{u^2} du = \lim_{b \rightarrow \infty} \left[ -\frac{144}{u} \right]_0^b$ .

**Total for part (b) 3 points**

- (c) Let  $h$  be the function defined by  $h(x) = x \cdot f'(x)$ . Find the value of  $\int_0^3 h(x) \, dx$ .

|                                                                                                                                                                         |                                                         |                |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------|----------------|
| Using integration by parts,<br>$u = x \quad dv = f'(x) \, dx$<br>$du = dx \quad v = f(x)$                                                                               | $u$ and $dv$                                            | <b>1 point</b> |
| $\int h(x) \, dx = \int x \cdot f'(x) \, dx = x \cdot f(x) - \int f(x) \, dx$                                                                                           | $\int h(x) \, dx$<br>$= x \cdot f(x) - \int f(x) \, dx$ | <b>1 point</b> |
| $\int_0^3 h(x) \, dx = \int_0^3 x \cdot f'(x) \, dx = x \cdot f(x) \Big _0^3 - \int_0^3 f(x) \, dx$<br>$= (3 \cdot f(3) - 0 \cdot f(0)) - 10 = 3 \cdot 2 - 0 - 10 = -4$ | Answer                                                  | <b>1 point</b> |

**Scoring notes:**

- The first and second points are earned with an implied  $u$  and  $dv$  in the presence of  $x \cdot f(x) - \int f(x) \, dx$  or  $x \cdot f(x) \Big|_0^3 - 10$ .
- Limits of integration may be present, omitted, or partially present in the work for the first and second points.
- The tabular method may be used to show integration by parts. In this case, the first point is earned by having columns (labeled or unlabeled) that begin with  $x$  and  $f'(x)$ . The second point is earned for  $x \cdot f(x) - \int f(x) \, dx$ .
- The third point is earned only for the correct answer and can only be earned if the first 2 points were earned.

**Total for part (c)    3 points**

**Total for question 5    9 points**