

Part B (AB or BC): Graphing calculator not allowed**Question 3****9 points****General Scoring Notes**

- The model solution is presented using standard mathematical notation.
- Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be accurate to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

A student starts reading a book at time $t = 0$ minutes and continues reading for the next 10 minutes. The rate at which the student reads is modeled by the differentiable function R , where $R(t)$ is measured in words per minute. Selected values of $R(t)$ are given in the table shown.

t (minutes)	0	2	8	10
$R(t)$ (words per minute)	90	100	150	162

	Model Solution	Scoring	
A	Approximate $R'(1)$ using the average rate of change of R over the interval $0 \leq t \leq 2$. Show the work that leads to your answer. Indicate units of measure.		
	$R'(1) \approx \frac{R(2) - R(0)}{2 - 0}$ $= \frac{100 - 90}{2} = \frac{10}{2} = 5 \text{ words per minute per minute}$	Answer with setup	Point 1 (P1)
		Units	Point 2 (P2)

Scoring Notes for Part A

- To earn **P1**, a response must present the answer along with the supporting work of a difference and a quotient using values from the table.
 - $\frac{100 - 90}{2 - 0}$, $\frac{10}{2 - 0}$, $\frac{100 - 90}{2}$, or $\frac{R(2) - R(0)}{2 - 0} = 5$ is sufficient to earn **P1**.
 - $\frac{R(2) - R(0)}{2 - 0}$ by itself is not sufficient to earn **P1**.
- P2** is earned for correct units, whether or not they are attached to a numerical value for the average rate of change.
- P2** is also earned for the units “words / minute² .”

B Must there be a value c , for $0 < c < 10$, such that $R(c) = 155$? Justify your answer.

R is differentiable implies R is continuous.

Differentiable
implies continuous

Point 3 (P3)

$$R(0) = 90 < 155 < R(10) = 162$$

Answer with
justification

Point 4 (P4)

Therefore, by the Intermediate Value Theorem, there must be a value c , with $0 < c < 10$, such that $R(c) = 155$.

Scoring Notes for Part B

- To earn **P3**, a response must state that R is continuous because R is differentiable (or equivalent). A response that simply states “ R is continuous” without justification does not earn **P3**.
- A response does not need to earn **P3** to be eligible for **P4**.
- To earn **P4**, a response must indicate that $R(0) < 155$ (or $R(2) < 155$ or $R(8) < 155$) and $R(10) > 155$, state that “ R is continuous,” and answer “yes” in some way.
- To earn **P4**, a response need not explicitly name the Intermediate Value Theorem, but if a theorem is named, it must be correct.

- C Use a trapezoidal sum with the three subintervals indicated by the data in the table to approximate the value of $\int_0^{10} R(t) dt$. Show the work that leads to your answer.

$\int_0^{10} R(t) dt$ $\approx \frac{R(0) + R(2)}{2}(2 - 0) + \frac{R(2) + R(8)}{2}(8 - 2)$ $+ \frac{R(8) + R(10)}{2}(10 - 8)$	Form of trapezoidal sum	Point 5 (P5)
$= \frac{90 + 100}{2}(2 - 0) + \frac{100 + 150}{2}(8 - 2) + \frac{150 + 162}{2}(10 - 8)$ $= \frac{190}{2}(2) + \frac{250}{2}(6) + \frac{312}{2}(2) = 190 + 750 + 312 = 1252$	Answer with supporting work	Point 6 (P6)

Scoring Notes for Part C

- Read “=” as “ $\approx$ ” for **P5**.
- The form of a trapezoidal sum includes three terms, each of which includes a product of two factors, where one of the factors incorporates the $\frac{1}{2}$ as part of the product. To earn **P5**, at least five of the six factors must be correct. If any of the six factors is incorrect, the response does not earn **P6**. Consider the following examples:
 - $\frac{90 + 100}{2}(2 - 0) + \frac{100 + 150}{2}(8 - 2) + \frac{150 + 162}{2}(10 - 8)$ earns **P5** and is sufficient to earn **P6**.
 - $\frac{190}{2}(2) + \frac{250}{2}(6) + \frac{312}{2}(2)$ earns **P5** and is sufficient to earn **P6**.
 - $\frac{1}{2}((R(0) + R(2))(2) + (R(2) + R(8))(6) + (R(8) + R(10))(2))$ earns **P5** and is eligible for **P6**.
 - $\frac{90 + 100}{2}(2) + \frac{100 + 150}{2}(2) + \frac{150 + 162}{2}(2)$ earns **P5** but is not eligible for **P6**.
(Note that the factor of 2 in the second term of this expression is incorrect.)
- Special case:** A response of $(90 + 100) + (100 + 150)3 + (150 + 162)$ earns both **P5** and **P6**.
- To be eligible for **P6**, a response must have earned **P5**.
Special case: A response of $95 \cdot 2 + 125 \cdot 6 + 156 \cdot 2$ earns **P6** but does not earn **P5**.
- A response of $\frac{90 + 100}{2}(2 - 0) + \frac{100 + 150}{2}(8 - 2) + \frac{150 + 162}{2}(10 - 8)$ or equivalent earns **P6** (i.e., subsequent errors in simplification will not be considered in scoring for **P6**).
- A response of $\frac{(90 \cdot 2 + 100 \cdot 6 + 150 \cdot 2) + (100 \cdot 2 + 150 \cdot 6 + 162 \cdot 2)}{2}$ or equivalent earns both **P5** and **P6**. (Note that the average of the left Riemann sum and right Riemann sum is equivalent to the trapezoidal sum.)
- A completely correct left Riemann sum (e.g., $90 \cdot 2 + 100 \cdot 6 + 150 \cdot 2 = 1080$) or a completely correct right Riemann sum (e.g., $100 \cdot 2 + 150 \cdot 6 + 162 \cdot 2 = 1424$) earns **P5** but does not earn **P6**.

- D** A teacher also starts reading at time $t = 0$ minutes and continues reading for the next 10 minutes. The rate at which the teacher reads is modeled by the function W defined by $W(t) = -\frac{3}{10}t^2 + 8t + 100$, where $W(t)$ is measured in words per minute. Based on the model, how many words has the teacher read by the end of the 10 minutes? Show the work that leads to your answer.

$$\int_0^{10} W(t) \, dt = \int_0^{10} \left(-\frac{3}{10}t^2 + 8t + 100 \right) dt$$

Integrand **Point 7 (P7)**

$$= \left(-\frac{1}{10}t^3 + 4t^2 + 100t \right) \Big|_0^{10}$$

Antiderivative **Point 8 (P8)**

$$= \left(-\frac{1}{10} \cdot 1000 + 4 \cdot 100 + 100 \cdot 10 \right) - \left(-\frac{1}{10} \cdot 0 + 4 \cdot 0 + 100 \cdot 0 \right) \\ = 1300$$

Answer **Point 9 (P9)**

Based on the model, the teacher has read 1300 words by the end of the 10 minutes.

Scoring Notes for Part D

- **P7** is earned for an indefinite or definite integral with integrand $W(t)$, with or without the differential dt .
- **P8** is earned for the correct antiderivative, with or without the constant of integration.
- To be eligible for **P9**, a response must have earned **P8**.
- A response of $\left(-\frac{1}{10} \cdot 1000 + 4 \cdot 100 + 100 \cdot 10 \right) - \left(-\frac{1}{10} \cdot 0 + 4 \cdot 0 + 100 \cdot 0 \right)$ or equivalent banks **P9** (i.e., subsequent errors in simplification will not be considered in scoring for **P9**).

Part A (AB or BC): Graphing calculator required**Question 1****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

t (minutes)	0	3	7	12
$C(t)$ (degrees Celsius)	100	85	69	55

The temperature of coffee in a cup at time t minutes is modeled by a decreasing differentiable function C , where $C(t)$ is measured in degrees Celsius. For $0 \leq t \leq 12$, selected values of $C(t)$ are given in the table shown.

Model Solution**Scoring**

- (a) Approximate $C'(5)$ using the average rate of change of C over the interval $3 \leq t \leq 7$. Show the work that leads to your answer and include units of measure.

$$C'(5) \approx \frac{C(7) - C(3)}{7 - 3} = \frac{69 - 85}{4} = -4 \text{ degrees Celsius per minute}$$

Estimate with supporting work **1 point**

Units **1 point**

Scoring notes:

- To earn the first point a response must include a difference and a quotient as the supporting work.
- $\frac{-16}{7-3}$, $\frac{69-85}{7-3}$, or $\frac{69-85}{4}$ is sufficient to earn the first point.
- A response that presents only units without a numerical approximation for $C'(5)$ does not earn the second point.
- The second point is also earned for “degrees per minute” attached to a numerical value.

Total for part (a) 2 points

- (b) Use a left Riemann sum with the three subintervals indicated by the data in the table to approximate the value of $\int_0^{12} C(t) dt$. Interpret the meaning of $\frac{1}{12} \int_0^{12} C(t) dt$ in the context of the problem.

$\int_0^{12} C(t) dt \approx (3 - 0) \cdot C(0) + (7 - 3) \cdot C(3) + (12 - 7) \cdot C(7)$ $= 3 \cdot 100 + 4 \cdot 85 + 5 \cdot 69 = 985$	Form of left Riemann sum	1 point
	Estimate	1 point
$\frac{1}{12} \int_0^{12} C(t) dt$ is the average temperature of the coffee (in degrees Celsius) over the interval from $t = 0$ to $t = 12$.	Interpretation	1 point

Scoring notes:

- Read “=” as “ $\approx$ ” for the first point.
- To earn the first point at least five of the six factors in the Riemann sum must be correct. If any of the six factors is incorrect, the response does not earn the second point.
- A response of $(3 - 0) \cdot C(0) + (7 - 3) \cdot C(3) + (12 - 7) \cdot C(7)$ earns the first point. Values must be pulled from the table to earn the second point.
- A response of $3 \cdot 100 + 4 \cdot 85 + 5 \cdot 69$ earns both the first and second points, unless there is a subsequent error in simplification, in which case the response would earn only the first point.
- A completely correct right Riemann sum (e.g., $3 \cdot 85 + 4 \cdot 69 + 5 \cdot 55$) earns 1 of the first 2 points. An unsupported answer of 806 does not earn either of the first 2 points.
- Units will not affect scoring for the second point.
- To earn the third point the interpretation must include both “average temperature” and the time interval. The response need not include a reference to units. However, if incorrect units are given in the interpretation, the response does not earn the third point.

Total for part (b) 3 points

- (c) For $12 \leq t \leq 20$, the rate of change of the temperature of the coffee is modeled by $C'(t) = \frac{-24.55e^{0.01t}}{t}$, where $C'(t)$ is measured in degrees Celsius per minute. Find the temperature of the coffee at time $t = 20$. Show the setup for your calculations.

$C(20) = C(12) + \int_{12}^{20} C'(t) dt$	Integral	1 point
	Uses initial condition	1 point
$= 55 - 14.670812 = 40.329188$	Answer	1 point
The temperature of the coffee at time $t = 20$ is 40.329 degrees Celsius.		

Scoring notes:

- The first point is earned for a definite integral with integrand $C'(t)$. If the limits of integration are incorrect, the response does not earn the third point.
- A linkage error such as $C(20) = \int_{12}^{20} C'(t) dt = 55 - 14.670812$ or $\int_{12}^{20} C'(t) dt = -14.670812 = 40.329188$ earns the first 2 points but does not earn the third point.
- Missing differential (dt):
 - Unambiguous responses of $C(20) = C(12) + \int_{12}^{20} C'(t)$ or $C(20) = 55 + \int_{12}^{20} C'(t)$ earn the first 2 points and are eligible for the third point.
 - Ambiguous responses of $C(20) = \int_{12}^{20} C'(t) + C(12)$ or $C(20) = \int_{12}^{20} C'(t) + 55$ do not earn the first point, earn the second point, and earn the third point if the given numeric answer is correct. If there is no numeric answer given, these responses do not earn the third point.
- The second point is earned for adding $C(12)$ or 55 to a definite integral with a lower limit of 12, either symbolically or numerically.
- The third point is earned for an answer of $55 - 14.671$ or $-14.671 + 55$ with no additional simplification, provided there is some supporting work for these values.
- An answer of just 40.329 with no supporting work does not earn any points.

Total for part (c) 3 points

(d) For the model defined in part (c), it can be shown that $C''(t) = \frac{0.2455e^{0.01t}(100-t)}{t^2}$. For $12 < t < 20$, determine whether the temperature of the coffee is changing at a decreasing rate or at an increasing rate. Give a reason for your answer.		
Because $C''(t) > 0$ on the interval $12 < t < 20$, the rate of change in the temperature of the coffee, $C'(t)$, is increasing on this interval. That is, on the interval $12 < t < 20$, the temperature of the coffee is changing at an increasing rate.	Answer with reason	1 point

Scoring notes:

- This point is earned only for a correct answer with a correct reason that references the sign of the second derivative of C .
- A response that provides a reason based on the evaluation of $C''(t)$ at a single point does not earn this point.
- A response that uses ambiguous pronouns (such as “It is positive, so increasing”) does not earn this point.
- A response does not need to reference the interval $12 < t < 20$ to earn the point.

Total for part (d) 1 point**Total for question 1 9 points**

Part A (AB or BC): Graphing calculator required**Question 1****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

t (seconds)	0	60	90	120	135	150
$f(t)$ (gallons per second)	0	0.1	0.15	0.1	0.05	0

A customer at a gas station is pumping gasoline into a gas tank. The rate of flow of gasoline is modeled by a differentiable function f , where $f(t)$ is measured in gallons per second and t is measured in seconds since pumping began. Selected values of $f(t)$ are given in the table.

Model Solution**Scoring**

- (a) Using correct units, interpret the meaning of $\int_{60}^{135} f(t) dt$ in the context of the problem. Use a right Riemann sum with the three subintervals $[60, 90]$, $[90, 120]$, and $[120, 135]$ to approximate the value of $\int_{60}^{135} f(t) dt$.

$\int_{60}^{135} f(t) dt$ represents the total number of gallons of gasoline pumped into the gas tank from time $t = 60$ seconds to time $t = 135$ seconds.

Interpretation with units **1 point**

$\int_{60}^{135} f(t) dt$
 $\approx f(90)(90 - 60) + f(120)(120 - 90) + f(135)(135 - 120)$
 $= (0.15)(30) + (0.1)(30) + (0.05)(15) = 8.25$

Form of Riemann sum **1 point**

Answer **1 point**

Scoring notes:

- To earn the first point the response must reference gallons of gasoline added/pumped and the time interval $t = 60$ to $t = 135$.
- To earn the second point at least five of the six factors in the Riemann sum must be correct.
- If there is any error in the Riemann sum, the response does not earn the third point.
- A response of $(0.15)(30) + (0.1)(30) + (0.05)(15)$ earns both the second and third points, unless there is a subsequent error in simplification, in which case the response would earn only the second point.

- A response that presents a correct value with accompanying work that shows the three products in the Riemann sum but does not show all six of the factors and/or the sum process, does not earn the second point but does earn the third point. For example, responses of either $4.5 + 3.0 + 0.75$ or $(0.15)(30)$, $0.1(30)$, $0.05(15) \rightarrow 8.25$ earn the third point but not the second.
- A response of $f(90)(90 - 60) + f(120)(120 - 90) + f(135)(135 - 120) = 8.25$ earns both the second and the third points.
- A response that presents an answer of only 8.25 does not earn either the second or third point.
- A response that provides a completely correct left Riemann sum with accompanying work, $f(60)(30) + f(90)(30) + f(120)(15) = 9$, or $(0.1)(30) + (0.15)(30) + (0.1)(15)$ earns 1 of the last 2 points. A response with any errors or missing factors in a left Riemann sum earns neither of the last 2 points.

Total for part (a) 3 points

(b) Must there exist a value of c , for $60 < c < 120$, such that $f'(c) = 0$? Justify your answer.

f is differentiable. $\Rightarrow f$ is continuous on $[60, 120]$.	$f(120) - f(60) = 0$	1 point
$\frac{f(120) - f(60)}{120 - 60} = \frac{0.1 - 0.1}{60} = 0$	Answer with justification	1 point
By the Mean Value Theorem, there must exist a c , for $60 < c < 120$, such that $f'(c) = 0$.		

Scoring notes:

- To earn the first point a response must present either $f(120) - f(60) = 0$, $0.1 - 0.1 = 0$ (perhaps as the numerator of a quotient), or $f(60) = f(120)$.
- To earn the second point a response must:
 - have earned the first point,
 - state that f is continuous because f is differentiable (or equivalent), and
 - answer “yes” in some way.
- A response may reference either the Mean Value Theorem or Rolle’s Theorem.
- A response that references the Intermediate Value Theorem cannot earn the second point.

Total for part (b) 2 points

(c) The rate of flow of gasoline, in gallons per second, can also be modeled by

$$g(t) = \left(\frac{t}{500}\right) \cos\left(\left(\frac{t}{120}\right)^2\right) \text{ for } 0 \leq t \leq 150. \text{ Using this model, find the average rate of flow of}$$

gasoline over the time interval $0 \leq t \leq 150$. Show the setup for your calculations.

$\frac{1}{150 - 0} \int_0^{150} g(t) dt$	Average value formula	1 point
$= 0.0959967$	Answer	1 point

The average rate of flow of gasoline, in gallons per second, is 0.096 (or 0.095).

Scoring notes:

- The exact value of $\frac{1}{150} \int_0^{150} g(t) dt$ is $\frac{12}{125} \sin\left(\frac{25}{16}\right)$.
- A response may present the average value formula in single or multiple steps. For example, the following response earns both points: $\int_0^{150} g(t) dt = 14.399504$ so the average rate is 0.0959967.
- A response that presents the average value formula in multiple steps but provides incorrect or incomplete communication (e.g., $\int_0^{150} g(t) dt = \frac{14.399504}{150} = 0.0959967$) earns 1 out of 2 points.
- A response of $\int_0^{150} g(t) dt = 0.0959967$ does not earn either point.
- Degree mode: A response that presents answers obtained by using a calculator in degree mode does not earn the first point it would have otherwise earned. The response is generally eligible for all subsequent points (unless no answer is possible in degree mode or the question is made simpler by using degree mode). In degree mode, $\frac{1}{150} \int_0^{150} g(t) dt = 0.149981$ or 0.002618.

Total for part (c) 2 points

- (d) Using the model g defined in part (c), find the value of $g'(140)$. Interpret the meaning of your answer in the context of the problem.

$g'(140) \approx -0.004908$ $g'(140) = -0.005$ (or -0.004)	$g'(140)$ 1 point
The rate at which gasoline is flowing into the tank is decreasing at a rate of 0.005 (or 0.004) gallon per second per second at time $t = 140$ seconds.	Interpretation 1 point

Scoring notes:

- The exact value of $g'(140)$ is $\frac{1}{500} \cos\left(\frac{49}{36}\right) - \frac{49}{9000} \sin\left(\frac{49}{36}\right)$.
- The value of $g'(140)$ may appear only in the interpretation.
- To be eligible for the second point a response must present some numerical value for $g'(140)$.
- To earn the second point the interpretation must include “the rate of flow of gasoline is changing at a rate of [the declared value of $g'(140)$]” and “at $t = 140$ ” (or equivalent).
- An interpretation of “decreasing at a rate of -0.005 ” or “increasing at a rate of 0.005” does not earn the second point.
- Degree mode: In degree mode, $g'(140) = 0.001997$ or 0.00187.

Total for part (d) 2 points**Total for question 1 9 points**

Part B (AB or BC): Graphing calculator not allowed**Question 4****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

t (days)	0	3	7	10	12
$r'(t)$ (centimeters per day)	-6.1	-5.0	-4.4	-3.8	-3.5

An ice sculpture melts in such a way that it can be modeled as a cone that maintains a conical shape as it decreases in size. The radius of the base of the cone is given by a twice-differentiable function r , where $r(t)$ is measured in centimeters and t is measured in days. The table above gives selected values of $r'(t)$, the rate of change of the radius, over the time interval $0 \leq t \leq 12$.

Model Solution**Scoring**

- (a) Approximate $r''(8.5)$ using the average rate of change of r' over the interval $7 \leq t \leq 10$. Show the computations that lead to your answer, and indicate units of measure.

$$r''(8.5) \approx \frac{r'(10) - r'(7)}{10 - 7} = \frac{-3.8 - (-4.4)}{10 - 7}$$

$r''(8.5)$ with
supporting work **1 point**

$$= \frac{0.6}{3} = 0.2 \text{ centimeter per day per day}$$

Units **1 point**

Scoring notes:

- To earn the first point the supporting work must include at least a difference and a quotient.
- Simplification is not required to earn the first point. If the numerical value is simplified, it must be correct.
- The second point can be earned with an incorrect approximation for $r''(8.5)$ but cannot be earned without some value for $r''(8.5)$ presented.
- Units may be written in any equivalent form (such as cm/day^2).

Total for part (a) 2 points

(b) Is there a time t , $0 \leq t \leq 3$, for which $r'(t) = -6$? Justify your answer.

$r(t)$ is twice-differentiable. $\Rightarrow r'(t)$ is differentiable. $\Rightarrow r'(t)$ is continuous.	$r'(0) < -6 < r'(3)$	1 point
$r'(0) = -6.1 < -6 < -5.0 = r'(3)$ Therefore, by the Intermediate Value Theorem, there is a time t , $0 < t < 3$, such that $r'(t) = -6$.	Conclusion using Intermediate Value Theorem	1 point

Scoring notes:

- To earn the first point, the response must establish that -6 is between $r'(0)$ and $r'(3)$ (or -6.1 and -5). This statement may be represented symbolically (with or without including one or both endpoints in an inequality) or verbally. A response of “ $r'(t) = -6$ because $r'(0) = -6.1$ and $r'(3) = -5$ ” does not state that -6 is between -6.1 and -5 . Thus this response does not earn the first point.
- To earn the second point:
 - The response must state that $r'(t)$ is continuous because $r'(t)$ is differentiable (or because $r(t)$ is twice differentiable).
 - The response must have earned the first point.
 - Exception: A response of “ $r'(t) = -6$ because $r'(0) = -6.1$ and $r'(3) = -5$ ” does not earn the first point because of imprecise communication but may nonetheless earn the second point if all other criteria for the second point are met.
 - The response must conclude that there is a time t such that $r'(t) = -6$. (A statement of “yes” would be sufficient.)
- To earn the second point, the response need not explicitly name the Intermediate Value Theorem, but if a theorem is named, it must be correct.

Total for part (b) 2 points

- (c) Use a right Riemann sum with the four subintervals indicated in the table to approximate the value of $\int_0^{12} r'(t) dt$.

$\int_0^{12} r'(t) dt \approx 3r'(3) + 4r'(7) + 3r'(10) + 2r'(12)$	Form of right Riemann sum	1 point
$= 3(-5.0) + 4(-4.4) + 3(-3.8) + 2(-3.5)$ $= -51$	Answer	1 point

Scoring notes:

- To earn the first point, at least seven of the eight factors in the Riemann sum must be correct. If there is any error in the Riemann sum, the response does not earn the second point.
- A response of $3(-5.0) + 4(-4.4) + 3(-3.8) + 2(-3.5)$ earns both the first and second points, unless there is a subsequent error in simplification, in which case the response would earn only the first point.
- A response that presents the correct answer, with accompanying work that shows the four products in the Riemann sum (without explicitly showing all of the factors and/or the sum process) does not earn the first point but earns the second point. For example, $-15 + 4(-4.4) + 3(-3.8) + -7$ does not earn the first point but earns the second point. Similarly, $-15, -17.6, -11.4, -7 \rightarrow -51$ does not earn the first point but earns the second point.
- A response that presents the correct answer (-51) with no supporting work earns no points.
- A response that provides a completely correct left Riemann sum and approximation $\int_0^{12} r'(t) dt$ (i.e., $3r'(0) + 4r'(3) + 3r'(7) + 2r'(10) = 3(-6.1) + 4(-5.0) + 3(-4.4) + 2(-3.8) = -59.1$) earns 1 of the 2 points. A response that has any error in a left Riemann sum or evaluation for $\int_0^{12} r'(t) dt$ earns no points.
- Units are not required or read in this part.

Total for part (c) 2 points

- (d) The height of the cone decreases at a rate of 2 centimeters per day. At time $t = 3$ days, the radius is 100 centimeters and the height is 50 centimeters. Find the rate of change of the volume of the cone with respect to time, in cubic centimeters per day, at time $t = 3$ days. (The volume V of a cone with radius r and height h is $V = \frac{1}{3}\pi r^2 h$.)

$\frac{dV}{dt} = \frac{2}{3}\pi r h \frac{dr}{dt} + \frac{1}{3}\pi r^2 \frac{dh}{dt}$	Product rule	1 point
	Chain rule	1 point
$\left. \frac{dV}{dt} \right _{t=3} = \frac{2}{3}\pi(100)(50)(-5) + \frac{1}{3}\pi(100)^2(-2) = -\frac{70,000\pi}{3}$	Answer	1 point
The rate of change of the volume of the sculpture at $t = 3$ is approximately $-\frac{70,000\pi}{3}$ cubic centimeters per day.		

Scoring notes:

- The first 2 points could be earned in either order.
- A response with a completely correct product rule, missing one or both of the correct differentials, earns the product rule point, but not the chain rule point. For example, $\frac{dV}{dt} = \frac{2}{3}\pi r h + \frac{1}{3}\pi r^2$ earns the first point, but not the second.
- A response that treats r or h (but not both) as a constant is eligible for the chain rule point but not the product rule point. For example, $\frac{dV}{dt} = \frac{2}{3}\pi r h \frac{dr}{dt}$ is correct if h is constant, and thus earns the chain rule point.
- Note: Neither $\frac{dV}{dt} = \frac{2}{3}\pi r \frac{dh}{dt}$ nor $\frac{dV}{dt} = \frac{2}{3}\pi r h \frac{dr}{dt} \frac{dh}{dt}$ earns any points.
- A response that assumes a functional relationship between r and h (such as $r = 2h$), and uses this relationship to create a function for volume in terms of one variable, is eligible for at most the chain rule point. For example, $r = 2h \rightarrow V = \frac{1}{3}\pi(2h)^2 h = \frac{4}{3}\pi h^3 \rightarrow \frac{dV}{dt} = 4\pi h^2 \frac{dh}{dt}$ earns only the chain rule point.
- A response that mishandles the constant $\frac{1}{3}\pi$ cannot earn the third point but is eligible for the first 2 points.
- The third point cannot be earned without both of the first 2 points.
- $\frac{dV}{dt} = \frac{2}{3}\pi(100)(50)(-5) + \frac{1}{3}\pi(100)^2(-2)$ earns all 3 points.
- Units are not required or read in this part.

Total for part (d) 3 points

Total for question 4 9 points

Part A (AB or BC): Graphing calculator required**Question 1****9 points****General Scoring Notes**

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Scoring guidelines and notes contain examples of the most common approaches seen in student responses. These guidelines can be applied to alternate approaches to ensure that these alternate approaches are scored appropriately.

r (centimeters)	0	1	2	2.5	4
$f(r)$ (milligrams per square centimeter)	1	2	6	10	18

The density of a bacteria population in a circular petri dish at a distance r centimeters from the center of the dish is given by an increasing, differentiable function f , where $f(r)$ is measured in milligrams per square centimeter. Values of $f(r)$ for selected values of r are given in the table above.

Model Solution**Scoring**

- (a) Use the data in the table to estimate $f'(2.25)$. Using correct units, interpret the meaning of your answer in the context of this problem.

$$f'(2.25) \approx \frac{f(2.5) - f(2)}{2.5 - 2} = \frac{10 - 6}{0.5} = 8$$

Estimate **1 point**

At a distance of $r = 2.25$ centimeters from the center of the petri dish, the density of the bacteria population is increasing at a rate of 8 milligrams per square centimeter per centimeter.

Interpretation with units **1 point**

Scoring notes:

- To earn the first point the response must provide both a difference and a quotient and must explicitly use values of f from the table.
- Simplification of the numerical value is not required to earn the first point. If the numerical value is simplified, it must be correct.
- The interpretation requires all of the following: distance $r = 2.25$, density of bacteria (population) is increasing or changing, at a rate of 8, and units of milligrams per square centimeter per centimeter.
- The second point (interpretation) cannot be earned without a nonzero presented value for $f'(2.25)$.
- To earn the second point the interpretation must be consistent with the presented nonzero value for $f'(2.25)$. In particular, if the presented value for $f'(2.25)$ is negative, the interpretation must include “decreasing at a rate of $|f'(2.25)|$ ” or “changing at a rate of $f'(2.25)$.” The second point cannot be earned for an incorrect statement such as “the bacteria density is decreasing at a rate of $-8 \dots$ ” even for a presented $f'(2.25) = -8$.
- The units ($\text{mg}/\text{cm}^2/\text{cm}$) may be attached to the estimate of $f'(2.25)$ and, if so, do not need to be repeated in the interpretation.
- If units attached to the estimate do not agree with units in the interpretation, read the units in the interpretation.

Total for part (a) 2 points

- (b) The total mass, in milligrams, of bacteria in the petri dish is given by the integral expression

$2\pi \int_0^4 r f(r) dr$. Approximate the value of $2\pi \int_0^4 r f(r) dr$ using a right Riemann sum with the four subintervals indicated by the data in the table.

$2\pi \int_0^4 r f(r) dr \approx 2\pi(1 \cdot f(1) \cdot (1 - 0) + 2 \cdot f(2) \cdot (2 - 1) + 2.5 \cdot f(2.5) \cdot (2.5 - 2) + 4 \cdot f(4) \cdot (4 - 2.5))$	Right Riemann sum setup	1 point
$= 2\pi(1 \cdot 2 \cdot 1 + 2 \cdot 6 \cdot 1 + 2.5 \cdot 10 \cdot 0.5 + 4 \cdot 18 \cdot 1.5)$ $= 269\pi = 845.088$	Approximation	1 point

Scoring notes:

- The presence or absence of 2π has no bearing on earning the first point.
- The first point is earned for a sum of four products with at most one error in any single value among the four products. Multiplication by 1 in any term does not need to be shown, but all other products must be explicitly shown.
- A response of $1 \cdot f(1) \cdot (1 - 0) + 2 \cdot f(2) \cdot (2 - 1) + 2.5 \cdot f(2.5) \cdot (2.5 - 2) + 4 \cdot f(4) \cdot (4 - 2.5)$ earns the first point but not the second.
- A response with any error in the Riemann sum is not eligible for the second point.
- A response that provides a completely correct left Riemann sum for $2\pi \int_0^4 r f(r) dr$ and approximation (91π) earns one of the two points. A response that has any error in a left Riemann sum or evaluation for $2\pi \int_0^4 r f(r) dr$ earns no points.
- A response that provides a completely correct right Riemann sum for $2\pi \int_0^4 r f(r) dr$ and approximation (80π) earns one of the two points. A response that has any error in a right Riemann sum or evaluation for $2\pi \int_0^4 r f(r) dr$ earns no points.
- Simplification of the numerical value is not required to earn the second point. If a numerical value is given, it must be correct to three decimal places.

Total for part (b) 2 points

- (c) Is the approximation found in part (b) an overestimate or underestimate of the total mass of bacteria in the petri dish? Explain your reasoning.

$\frac{d}{dr}(rf(r)) = f(r) + rf'(r)$	Product rule expression for $\frac{d}{dr}(rf(r))$	1 point
Because f is nonnegative and increasing, $\frac{d}{dr}(rf(r)) > 0$ on the interval $0 \leq r \leq 4$. Thus, the integrand $rf(r)$ is strictly increasing. Therefore, the right Riemann sum approximation of $2\pi \int_0^4 rf(r) dr$ is an overestimate.	Answer with explanation	1 point

Scoring notes:

- To earn the second point a response must explain that $rf(r)$ is increasing and, therefore, the right Riemann sum is an overestimate. The second point can be earned without having earned the first point.
- A response that attempts to explain based on a left Riemann sum for $2\pi \int_0^4 rf(r) dr$ from part (b) earns no points.
- A response that attempts to explain based on a right Riemann sum for $2\pi \int_0^4 f(r) dr$ from part (b) earns no points.

Total for part (c) 2 points

- (d) The density of bacteria in the petri dish, for $1 \leq r \leq 4$, is modeled by the function g defined by $g(r) = 2 - 16(\cos(1.57\sqrt{r}))^3$. For what value of k , $1 < k < 4$, is $g(k)$ equal to the average value of $g(r)$ on the interval $1 \leq r \leq 4$?

Average value = $g_{\text{avg}} = \frac{1}{4-1} \int_1^4 g(r) dr$	Definite integral	1 point
$\frac{1}{4-1} \int_1^4 g(r) dr = 9.875795$	Average value	1 point
$g(k) = g_{\text{avg}} \Rightarrow k = 2.497$	Answer	1 point

Scoring notes:

- The first point is earned for a definite integral, with or without $\frac{1}{4-1}$ or $\frac{1}{3}$.
- A response that presents a definite integral with incorrect limits but a correct integrand earns the first point.
- Presentation of the numerical value 9.875795 is not required to earn the second point. This point can be earned by the average value setup: $\frac{1}{3} \int_1^4 g(r) dr$.
- Once earned for the average value setup, the second point cannot be lost. Subsequent errors will result in not earning the third point.
- The third point is earned only for the value $k = 2.497$.
- The third point cannot be earned without the second.
- Special case: A response that does not provide the average value setup but presents an average value of -13.955 is using degree mode on their calculator. This response would not earn the second point but could earn the third point for an answer of $k = 2.5$ (or 2.499).

Total for part (d) 3 points

Total for question 1 9 points

2018

AP[®]

CollegeBoard

AP Calculus BC

Scoring Guidelines

2018 SCORING GUIDELINES

Question 1

(a) $\int_0^{300} r(t) dt = 270$

According to the model, 270 people enter the line for the escalator during the time interval $0 \leq t \leq 300$.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) $20 + \int_0^{300} (r(t) - 0.7) dt = 20 + \int_0^{300} r(t) dt - 0.7 \cdot 300 = 80$

According to the model, 80 people are in line at time $t = 300$.

2 : $\begin{cases} 1 : \text{considers rate out} \\ 1 : \text{answer} \end{cases}$

(c) Based on part (b), the number of people in line at time $t = 300$ is 80.

The first time t that there are no people in line is

$$300 + \frac{80}{0.7} = 414.286 \text{ (or 414.285) seconds.}$$

1 : answer

(d) The total number of people in line at time t , $0 \leq t \leq 300$, is modeled by

$$20 + \int_0^t r(x) dx - 0.7t.$$

$$r(t) - 0.7 = 0 \Rightarrow t_1 = 33.013298, t_2 = 166.574719$$

4 : $\begin{cases} 1 : \text{considers } r(t) - 0.7 = 0 \\ 1 : \text{identifies } t = 33.013 \\ 1 : \text{answers} \\ 1 : \text{justification} \end{cases}$

t	People in line for escalator
0	20
t_1	3.803
t_2	158.070
300	80

The number of people in line is a minimum at time $t = 33.013$ seconds, when there are 4 people in line.

2018 SCORING GUIDELINES

Question 2

(a) $p'(25) = -1.179$

At a depth of 25 meters, the density of plankton cells is changing at a rate of -1.179 million cells per cubic meter per meter.

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{meaning with units} \end{cases}$$

(b) $\int_0^{30} 3p(h) \, dh = 1675.414936$

There are 1675 million plankton cells in the column of water between $h = 0$ and $h = 30$ meters.

$$2 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$$

(c) $\int_{30}^K 3f(h) \, dh$ represents the number of plankton cells, in millions, in the column of water from a depth of 30 meters to a depth of K meters.

The number of plankton cells, in millions, in the entire column of water is given by $\int_0^{30} 3p(h) \, dh + \int_{30}^K 3f(h) \, dh$.

Because $0 \leq f(h) \leq u(h)$ for all $h \geq 30$,

$$3 \int_{30}^K f(h) \, dh \leq 3 \int_{30}^K u(h) \, dh \leq 3 \int_{30}^{\infty} u(h) \, dh = 3 \cdot 105 = 315.$$

The total number of plankton cells in the column of water is bounded by $1675.415 + 315 = 1990.415 \leq 2000$ million.

$$3 : \begin{cases} 1 : \text{integral expression} \\ 1 : \text{compares improper integral} \\ 1 : \text{explanation} \end{cases}$$

(d) $\int_0^1 \sqrt{(x'(t))^2 + (y'(t))^2} \, dt = 757.455862$

The total distance traveled by the boat over the time interval $0 \leq t \leq 1$ is 757.456 (or 757.455) meters.

$$2 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{total distance} \end{cases}$$

2018 SCORING GUIDELINES

Question 3

$$\begin{aligned} \text{(a)} \quad f(-5) &= f(1) + \int_1^{-5} g(x) \, dx = f(1) - \int_{-5}^1 g(x) \, dx \\ &= 3 - \left(-9 - \frac{3}{2} + 1 \right) = 3 - \left(-\frac{19}{2} \right) = \frac{25}{2} \end{aligned}$$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b)} \quad \int_1^6 g(x) \, dx &= \int_1^3 g(x) \, dx + \int_3^6 g(x) \, dx \\ &= \int_1^3 2 \, dx + \int_3^6 2(x-4)^2 \, dx \\ &= 4 + \left[\frac{2}{3}(x-4)^3 \right]_{x=3}^{x=6} = 4 + \frac{16}{3} - \left(-\frac{2}{3} \right) = 10 \end{aligned}$$

3 : $\begin{cases} 1 : \text{split at } x = 3 \\ 1 : \text{antiderivative of } 2(x-4)^2 \\ 1 : \text{answer} \end{cases}$

(c) The graph of f is increasing and concave up on $0 < x < 1$ and $4 < x < 6$ because $f'(x) = g(x) > 0$ and $f''(x) = g'(x)$ is increasing on those intervals.

2 : $\begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$

(d) The graph of f has a point of inflection at $x = 4$ because $f'(x) = g(x)$ changes from decreasing to increasing at $x = 4$.

2 : $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

2018 SCORING GUIDELINES

Question 4

$$(a) \quad H'(6) \approx \frac{H(7) - H(5)}{7 - 5} = \frac{11 - 6}{2} = \frac{5}{2}$$

$H'(6)$ is the rate at which the height of the tree is changing, in meters per year, at time $t = 6$ years.

$$(b) \quad \frac{H(5) - H(3)}{5 - 3} = \frac{6 - 2}{2} = 2$$

Because H is differentiable on $3 \leq t \leq 5$, H is continuous on $3 \leq t \leq 5$.

By the Mean Value Theorem, there exists a value c , $3 < c < 5$, such that $H'(c) = 2$.

$$(c) \quad \text{The average height of the tree over the time interval } 2 \leq t \leq 10 \text{ is given by } \frac{1}{10 - 2} \int_2^{10} H(t) \, dt.$$

$$\begin{aligned} \frac{1}{8} \int_2^{10} H(t) \, dt &\approx \frac{1}{8} \left(\frac{1.5 + 2}{2} \cdot 1 + \frac{2 + 6}{2} \cdot 2 + \frac{6 + 11}{2} \cdot 2 + \frac{11 + 15}{2} \cdot 3 \right) \\ &= \frac{1}{8} (65.75) = \frac{263}{32} \end{aligned}$$

The average height of the tree over the time interval $2 \leq t \leq 10$ is $\frac{263}{32}$ meters.

$$(d) \quad G(x) = 50 \Rightarrow x = 1$$

$$\frac{d}{dt}(G(x)) = \frac{d}{dx}(G(x)) \cdot \frac{dx}{dt} = \frac{(1+x)100 - 100x \cdot 1}{(1+x)^2} \cdot \frac{dx}{dt} = \frac{100}{(1+x)^2} \cdot \frac{dx}{dt}$$

$$\left. \frac{d}{dt}(G(x)) \right|_{x=1} = \frac{100}{(1+1)^2} \cdot 0.03 = \frac{3}{4}$$

According to the model, the rate of change of the height of the tree with respect to time when the tree is 50 meters tall is $\frac{3}{4}$ meter per year.

$$2 : \begin{cases} 1 : \text{estimate} \\ 1 : \text{interpretation with units} \end{cases}$$

$$2 : \begin{cases} 1 : \frac{H(5) - H(3)}{5 - 3} \\ 1 : \text{conclusion using Mean Value Theorem} \end{cases}$$

$$2 : \begin{cases} 1 : \text{trapezoidal sum} \\ 1 : \text{approximation} \end{cases}$$

$$3 : \begin{cases} 2 : \frac{d}{dt}(G(x)) \\ 1 : \text{answer} \end{cases}$$

Note: max 1/3 [1-0] if no chain rule

2018 SCORING GUIDELINES

Question 5

$$(a) \text{ Area} = \frac{1}{2} \int_{\pi/3}^{5\pi/3} (4^2 - (3 + 2\cos \theta)^2) d\theta$$

$$(b) \frac{dr}{d\theta} = -2\sin \theta \Rightarrow \left. \frac{dr}{d\theta} \right|_{\theta=\pi/2} = -2$$

$$r\left(\frac{\pi}{2}\right) = 3 + 2\cos\left(\frac{\pi}{2}\right) = 3$$

$$y = r \sin \theta \Rightarrow \frac{dy}{d\theta} = \frac{dr}{d\theta} \sin \theta + r \cos \theta$$

$$x = r \cos \theta \Rightarrow \frac{dx}{d\theta} = \frac{dr}{d\theta} \cos \theta - r \sin \theta$$

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/2} = \left. \frac{dy/d\theta}{dx/d\theta} \right|_{\theta=\pi/2} = \frac{-2\sin\left(\frac{\pi}{2}\right) + 3\cos\left(\frac{\pi}{2}\right)}{-2\cos\left(\frac{\pi}{2}\right) - 3\sin\left(\frac{\pi}{2}\right)} = \frac{2}{3}$$

The slope of the line tangent to the graph of $r = 3 + 2\cos \theta$

at $\theta = \frac{\pi}{2}$ is $\frac{2}{3}$.

— OR —

$$y = r \sin \theta = (3 + 2\cos \theta) \sin \theta \Rightarrow \frac{dy}{d\theta} = 3\cos \theta + 2\cos^2 \theta - 2\sin^2 \theta$$

$$x = r \cos \theta = (3 + 2\cos \theta) \cos \theta \Rightarrow \frac{dx}{d\theta} = -3\sin \theta - 4\sin \theta \cos \theta$$

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/2} = \left. \frac{dy/d\theta}{dx/d\theta} \right|_{\theta=\pi/2} = \frac{3\cos\left(\frac{\pi}{2}\right) + 2\cos^2\left(\frac{\pi}{2}\right) - 2\sin^2\left(\frac{\pi}{2}\right)}{-3\sin\left(\frac{\pi}{2}\right) - 4\sin\left(\frac{\pi}{2}\right)\cos\left(\frac{\pi}{2}\right)} = \frac{2}{3}$$

The slope of the line tangent to the graph of $r = 3 + 2\cos \theta$

at $\theta = \frac{\pi}{2}$ is $\frac{2}{3}$.

$$(c) \frac{dr}{dt} = \frac{dr}{d\theta} \cdot \frac{d\theta}{dt} = -2\sin \theta \cdot \frac{d\theta}{dt} \Rightarrow \frac{d\theta}{dt} = \frac{dr}{dt} \cdot \frac{1}{-2\sin \theta}$$

$$\left. \frac{d\theta}{dt} \right|_{\theta=\pi/3} = 3 \cdot \frac{1}{-2\sin\left(\frac{\pi}{3}\right)} = \frac{3}{-\sqrt{3}} = -\sqrt{3} \text{ radians per second}$$

$$3 : \begin{cases} 1 : \text{constant and limits} \\ 2 : \text{integrand} \end{cases}$$

$$3 : \begin{cases} 1 : \frac{dy}{d\theta} = \frac{dr}{d\theta} \sin \theta + r \cos \theta \\ \text{or } \frac{dx}{d\theta} = \frac{dr}{d\theta} \cos \theta - r \sin \theta \\ 1 : \frac{dy}{dx} = \frac{\frac{dr}{d\theta} \sin \theta + r \cos \theta}{\frac{dr}{d\theta} \cos \theta - r \sin \theta} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : \frac{dr}{dt} = \frac{dr}{d\theta} \cdot \frac{d\theta}{dt} \\ 1 : \frac{d\theta}{dt} = \frac{dr}{dt} \cdot \frac{1}{-2\sin \theta} \\ 1 : \text{answer with units} \end{cases}$$

2018 SCORING GUIDELINES

Question 6

(a) The first four nonzero terms are $\frac{x^2}{3} - \frac{x^3}{2 \cdot 3^2} + \frac{x^4}{3 \cdot 3^3} - \frac{x^5}{4 \cdot 3^4}$.

The general term is $(-1)^{n+1} \frac{x^{n+1}}{n \cdot 3^n}$.

(b)
$$\lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+2} x^{n+2}}{(n+1)(3^{n+1})}}{\frac{(-1)^{n+1} x^{n+1}}{n \cdot 3^n}} \right| = \lim_{n \rightarrow \infty} \left| \frac{-x}{3} \cdot \frac{n}{n+1} \right| = \left| \frac{x}{3} \right|$$

$\left| \frac{x}{3} \right| < 1$ for $|x| < 3$

Therefore, the radius of convergence of the Maclaurin series for f is 3.

— OR —

The radius of convergence of the Maclaurin series for $\ln(1+x)$ is 1, so the series for $f(x) = x \ln\left(1 + \frac{x}{3}\right)$ converges absolutely for $\left|\frac{x}{3}\right| < 1$.

$\left|\frac{x}{3}\right| < 1 \Rightarrow |x| < 3$

Therefore, the radius of convergence of the Maclaurin series for f is 3.

When $x = -3$, the series is $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{(-3)^{n+1}}{n \cdot 3^n} = \sum_{n=1}^{\infty} \frac{3}{n}$, which diverges by comparison to the harmonic series.

When $x = 3$, the series is $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{3^{n+1}}{n \cdot 3^n} = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{3}{n}$, which converges by the alternating series test.

The interval of convergence of the Maclaurin series for f is $-3 < x \leq 3$.

(c) By the alternating series error bound, an upper bound for $|P_4(2) - f(2)|$ is the magnitude of the next term of the alternating series.

$|P_4(2) - f(2)| < \left| -\frac{2^5}{4 \cdot 3^4} \right| = \frac{8}{81}$

2 : $\begin{cases} 1 : \text{first four terms} \\ 1 : \text{general term} \end{cases}$

5 : $\begin{cases} 1 : \text{sets up ratio} \\ 1 : \text{computes limit of ratio} \\ 1 : \text{radius of convergence} \\ 1 : \text{considers both endpoints} \\ 1 : \text{analysis and interval of convergence} \end{cases}$

— OR —

5 : $\begin{cases} 1 : \text{radius for } \ln(1+x) \text{ series} \\ 1 : \text{substitutes } \frac{x}{3} \\ 1 : \text{radius of convergence} \\ 1 : \text{considers both endpoints} \\ 1 : \text{analysis and interval of convergence} \end{cases}$

2 : $\begin{cases} 1 : \text{uses fifth-degree term as error bound} \\ 1 : \text{answer} \end{cases}$

2017**AP[®]** **CollegeBoard**

AP Calculus BC

Scoring Guidelines

2017 SCORING GUIDELINES

Question 1

(a) Volume = $\int_0^{10} A(h) \, dh$ $\approx (2 - 0) \cdot A(0) + (5 - 2) \cdot A(2) + (10 - 5) \cdot A(5)$ $= 2 \cdot 50.3 + 3 \cdot 14.4 + 5 \cdot 6.5$ $= 176.3$ cubic feet (b) The approximation in part (a) is an overestimate because a left Riemann sum is used and A is decreasing. (c) $\int_0^{10} f(h) \, dh = 101.325338$ The volume is 101.325 cubic feet. (d) Using the model, $V(h) = \int_0^h f(x) \, dx$. $\begin{aligned} \left. \frac{dV}{dt} \right _{h=5} &= \left[\frac{dV}{dh} \cdot \frac{dh}{dt} \right]_{h=5} \\ &= \left[f(h) \cdot \frac{dh}{dt} \right]_{h=5} \\ &= f(5) \cdot 0.26 = 1.694419 \end{aligned}$ When $h = 5$, the volume of water is changing at a rate of 1.694 cubic feet per minute.	1 : units in parts (a), (c), and (d) 2 : $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{approximation} \end{cases}$ 1 : overestimate with reason 2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$ 3 : $\begin{cases} 2 : \frac{dV}{dt} \\ 1 : \text{answer} \end{cases}$
--	---

2017 SCORING GUIDELINES

Question 2

(a) $\frac{1}{2} \int_0^{\pi/2} (f(\theta))^2 d\theta = 0.648414$

The area of R is 0.648.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) $\int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \frac{1}{2} \int_0^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$

— OR —

$$\int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \int_k^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

2 : $\begin{cases} 1 : \text{integral expression} \\ \quad \text{for one region} \\ 1 : \text{equation} \end{cases}$

(c) $w(\theta) = g(\theta) - f(\theta)$

$$w_A = \frac{\int_0^{\pi/2} w(\theta) d\theta}{\frac{\pi}{2} - 0} = 0.485446$$

The average value of $w(\theta)$ on the interval $\left[0, \frac{\pi}{2}\right]$ is 0.485.

3 : $\begin{cases} 1 : w(\theta) \\ 1 : \text{integral} \\ 1 : \text{average value} \end{cases}$

(d) $w(\theta) = w_A$ for $0 \leq \theta \leq \frac{\pi}{2} \Rightarrow \theta = 0.517688$

$w(\theta) = w_A$ at $\theta = 0.518$ (or 0.517).

$w'(0.518) < 0 \Rightarrow w(\theta)$ is decreasing at $\theta = 0.518$.

2 : $\begin{cases} 1 : \text{solves } w(\theta) = w_A \\ 1 : \text{answer with reason} \end{cases}$

2017 SCORING GUIDELINES

Question 3

$$(a) \quad f(-6) = f(-2) + \int_{-2}^{-6} f'(x) \, dx = 7 - \int_{-6}^{-2} f'(x) \, dx = 7 - 4 = 3$$

$$f(5) = f(-2) + \int_{-2}^5 f'(x) \, dx = 7 - 2\pi + 3 = 10 - 2\pi$$

$$3 : \begin{cases} 1 : \text{uses initial condition} \\ 1 : f(-6) \\ 1 : f(5) \end{cases}$$

$$(b) \quad f'(x) > 0 \text{ on the intervals } [-6, -2) \text{ and } (2, 5).$$

Therefore, f is increasing on the intervals $[-6, -2]$ and $[2, 5]$.

2 : answer with justification

$$(c) \quad \text{The absolute minimum will occur at a critical point where } f'(x) = 0 \text{ or at an endpoint.}$$

$$f'(x) = 0 \Rightarrow x = -2, x = 2$$

$$2 : \begin{cases} 1 : \text{considers } x = 2 \\ 1 : \text{answer with justification} \end{cases}$$

x	$f(x)$
-6	3
-2	7
2	$7 - 2\pi$
5	$10 - 2\pi$

The absolute minimum value is $f(2) = 7 - 2\pi$.

$$(d) \quad f''(-5) = \frac{2 - 0}{-6 - (-2)} = -\frac{1}{2}$$

$$2 : \begin{cases} 1 : f''(-5) \\ 1 : f''(3) \text{ does not exist,} \\ \quad \text{with explanation} \end{cases}$$

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} = 2 \text{ and } \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3} = -1$$

$f''(3)$ does not exist because

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} \neq \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3}.$$

2017 SCORING GUIDELINES

Question 4

$$(a) \quad H'(0) = -\frac{1}{4}(91 - 27) = -16$$

$$H(0) = 91$$

An equation for the tangent line is $y = 91 - 16t$.

The internal temperature of the potato at time $t = 3$ minutes is approximately $91 - 16 \cdot 3 = 43$ degrees Celsius.

$$(b) \quad \frac{d^2H}{dt^2} = -\frac{1}{4} \frac{dH}{dt} = \left(-\frac{1}{4}\right)\left(-\frac{1}{4}\right)(H - 27) = \frac{1}{16}(H - 27)$$

$$H > 27 \text{ for } t > 0 \Rightarrow \frac{d^2H}{dt^2} = \frac{1}{16}(H - 27) > 0 \text{ for } t > 0$$

Therefore, the graph of H is concave up for $t > 0$. Thus, the answer in part (a) is an underestimate.

$$(c) \quad \frac{dG}{(G - 27)^{2/3}} = -dt$$

$$\int \frac{dG}{(G - 27)^{2/3}} = \int (-1) dt$$

$$3(G - 27)^{1/3} = -t + C$$

$$3(91 - 27)^{1/3} = 0 + C \Rightarrow C = 12$$

$$3(G - 27)^{1/3} = 12 - t$$

$$G(t) = 27 + \left(\frac{12 - t}{3}\right)^3 \text{ for } 0 \leq t < 10$$

The internal temperature of the potato at time $t = 3$ minutes is

$$27 + \left(\frac{12 - 3}{3}\right)^3 = 54 \text{ degrees Celsius.}$$

$$3 : \begin{cases} 1 : \text{slope} \\ 1 : \text{tangent line} \\ 1 : \text{approximation} \end{cases}$$

1 : underestimate with reason

$$5 : \begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration and} \\ \quad \text{uses initial condition} \\ 1 : \text{equation involving } G \text{ and } t \\ 1 : G(t) \text{ and } G(3) \end{cases}$$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

2017 SCORING GUIDELINES

Question 5

$$(a) f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2}$$

$$f'(3) = \frac{(-3)(5)}{(18 - 21 + 5)^2} = -\frac{15}{4}$$

$$(b) f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2} = 0 \Rightarrow x = \frac{7}{4}$$

The only critical point in the interval $1 < x < 2.5$ has x -coordinate $\frac{7}{4}$.

f' changes sign from positive to negative at $x = \frac{7}{4}$.

Therefore, f has a relative maximum at $x = \frac{7}{4}$.

$$\begin{aligned} (c) \int_5^\infty f(x) dx &= \lim_{b \rightarrow \infty} \int_5^b \frac{3}{2x^2 - 7x + 5} dx = \lim_{b \rightarrow \infty} \int_5^b \left(\frac{2}{2x - 5} - \frac{1}{x - 1} \right) dx \\ &= \lim_{b \rightarrow \infty} \left[\ln(2x - 5) - \ln(x - 1) \right]_5^b = \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2x - 5}{x - 1} \right) \right]_5^b \\ &= \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2b - 5}{b - 1} \right) - \ln \left(\frac{5}{4} \right) \right] = \ln 2 - \ln \left(\frac{5}{4} \right) = \ln \left(\frac{8}{5} \right) \end{aligned}$$

(d) f is continuous, positive, and decreasing on $[5, \infty)$.

The series converges by the integral test since $\int_5^\infty \frac{3}{2x^2 - 7x + 5} dx$ converges.

— OR —

$$\frac{3}{2n^2 - 7n + 5} > 0 \text{ and } \frac{1}{n^2} > 0 \text{ for } n \geq 5.$$

Since $\lim_{n \rightarrow \infty} \frac{\frac{3}{2n^2 - 7n + 5}}{\frac{1}{n^2}} = \frac{3}{2}$ and the series $\sum_{n=5}^\infty \frac{1}{n^2}$ converges,

the series $\sum_{n=5}^\infty \frac{3}{2n^2 - 7n + 5}$ converges by the limit comparison test.

2 : $f'(3)$

2 : $\begin{cases} 1 : x\text{-coordinate} \\ 1 : \text{relative maximum} \\ \text{with justification} \end{cases}$

3 : $\begin{cases} 1 : \text{antiderivative} \\ 1 : \text{limit expression} \\ 1 : \text{answer} \end{cases}$

2 : answer with conditions

2017 SCORING GUIDELINES

Question 6

(a) $f(0) = 0$

$f'(0) = 1$

$f''(0) = -1(1) = -1$

$f'''(0) = -2(-1) = 2$

$f^{(4)}(0) = -3(2) = -6$

The first four nonzero terms are

$$0 + 1x + \frac{-1}{2!}x^2 + \frac{2}{3!}x^3 + \frac{-6}{4!}x^4 = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4}.$$

The general term is $\frac{(-1)^{n+1}x^n}{n}$.

(b) For $x = 1$, the Maclaurin series becomes $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n}$.

The series does not converge absolutely because the harmonic series diverges.

The series alternates with terms that decrease in magnitude to 0, and therefore the series converges conditionally.

$$\begin{aligned} \text{(c)} \quad \int_0^x f(t) dt &= \int_0^x \left(t - \frac{t^2}{2} + \frac{t^3}{3} - \frac{t^4}{4} + \cdots + \frac{(-1)^{n+1}t^n}{n} + \cdots \right) dt \\ &= \left[\frac{t^2}{2} - \frac{t^3}{3 \cdot 2} + \frac{t^4}{4 \cdot 3} - \frac{t^5}{5 \cdot 4} + \cdots + \frac{(-1)^{n+1}t^{n+1}}{(n+1)n} + \cdots \right]_{t=0}^{t=x} \\ &= \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{12} - \frac{x^5}{20} + \cdots + \frac{(-1)^{n+1}x^{n+1}}{(n+1)n} + \cdots \end{aligned}$$

(d) The terms alternate in sign and decrease in magnitude to 0. By the alternating series error bound, the error $\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right|$ is bounded

by the magnitude of the first unused term, $\left| -\frac{(1/2)^5}{20} \right|$.

$$\text{Thus, } \left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right| \leq \left| -\frac{(1/2)^5}{20} \right| = \frac{1}{32 \cdot 20} < \frac{1}{500}.$$

$$3 : \begin{cases} 1 : f''(0), f'''(0), \text{ and } f^{(4)}(0) \\ 1 : \text{verify terms} \\ 1 : \text{general term} \end{cases}$$

2 : converges conditionally
with reason

$$3 : \begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \\ 1 : \text{general term} \end{cases}$$

1 : error bound

AP[®] Calculus BC

2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2016 SCORING GUIDELINES

Question 1

t (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.

- (a) Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.
- (b) Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
- (c) Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
- (d) For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

(a) $R'(2) \approx \frac{R(3) - R(1)}{3 - 1} = \frac{950 - 1190}{3 - 1} = -120$ liters/hr²

2 : $\begin{cases} 1 : \text{estimate} \\ 1 : \text{units} \end{cases}$

(b) The total amount of water removed is given by $\int_0^8 R(t) dt$.

$$\begin{aligned} \int_0^8 R(t) dt &\approx 1 \cdot R(0) + 2 \cdot R(1) + 3 \cdot R(3) + 2 \cdot R(6) \\ &= 1(1340) + 2(1190) + 3(950) + 2(740) \\ &= 8050 \text{ liters} \end{aligned}$$

This is an overestimate since R is a decreasing function.

3 : $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{estimate} \\ 1 : \text{overestimate with reason} \end{cases}$

(c) $\text{Total} \approx 50000 + \int_0^8 W(t) dt - 8050$
 $= 50000 + 7836.195325 - 8050 \approx 49786$ liters

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{estimate} \end{cases}$

(d) $W(0) - R(0) > 0$, $W(8) - R(8) < 0$, and $W(t) - R(t)$ is continuous.


2 : $\begin{cases} 1 : \text{considers } W(t) - R(t) \\ 1 : \text{answer with explanation} \end{cases}$

Therefore, the Intermediate Value Theorem guarantees at least one time t , $0 < t < 8$, for which $W(t) - R(t) = 0$, or $W(t) = R(t)$.

For this value of t , the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank.

2016 SCORING GUIDELINES

Question 2



At time t , the position of a particle moving in the xy -plane is given by the parametric functions $(x(t), y(t))$, where $\frac{dx}{dt} = t^2 + \sin(3t^2)$. The graph of y , consisting of three line segments, is shown in the figure above. At $t = 0$, the particle is at position $(5, 1)$.

- Find the position of the particle at $t = 3$.
- Find the slope of the line tangent to the path of the particle at $t = 3$.
- Find the speed of the particle at $t = 3$.
- Find the total distance traveled by the particle from $t = 0$ to $t = 2$.

$$(a) \quad x(3) = x(0) + \int_0^3 x'(t) dt = 5 + 9.377035 = 14.377$$

$$y(3) = -\frac{1}{2}$$

The position of the particle at $t = 3$ is $(14.377, -0.5)$.

$$(b) \quad \text{Slope} = \frac{y'(3)}{x'(3)} = \frac{0.5}{9.956376} = 0.05$$

$$(c) \quad \text{Speed} = \sqrt{(x'(3))^2 + (y'(3))^2} = 9.969 \text{ (or } 9.968)$$

$$\begin{aligned} (d) \quad \text{Distance} &= \int_0^2 \sqrt{(x'(t))^2 + (y'(t))^2} dt \\ &= \int_0^1 \sqrt{(x'(t))^2 + (-2)^2} dt + \int_1^2 \sqrt{(x'(t))^2 + 0^2} dt \\ &= 2.237871 + 2.112003 = 4.350 \text{ (or } 4.349) \end{aligned}$$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

1 : slope

2 : $\begin{cases} 1 : \text{expression for speed} \\ 1 : \text{answer} \end{cases}$

3 : $\begin{cases} 1 : \text{expression for distance} \\ 1 : \text{integrals} \\ 1 : \text{answer} \end{cases}$


2016 SCORING GUIDELINES

Question 3

The figure above shows the graph of the piecewise-linear function f . For $-4 \leq x \leq 12$, the function g is defined by

$$g(x) = \int_2^x f(t) \, dt.$$

- (a) Does g have a relative minimum, a relative maximum, or neither at $x = 10$? Justify your answer.
- (b) Does the graph of g have a point of inflection at $x = 4$? Justify your answer.
- (c) Find the absolute minimum value and the absolute maximum value of g on the interval $-4 \leq x \leq 12$. Justify your answers.
- (d) For $-4 \leq x \leq 12$, find all intervals for which $g(x) \leq 0$.

Graph of f

- (a) The function g has neither a relative minimum nor a relative maximum at $x = 10$ since $g'(x) = f(x)$ and $f(x) \leq 0$ for $8 \leq x \leq 12$.
- (b) The graph of g has a point of inflection at $x = 4$ since $g'(x) = f(x)$ is increasing for $2 \leq x \leq 4$ and decreasing for $4 \leq x \leq 8$.
- (c) $g'(x) = f(x)$ changes sign only at $x = -2$ and $x = 6$.

x	$g(x)$
-4	-4
-2	-8
6	8
12	-4

On the interval $-4 \leq x \leq 12$, the absolute minimum value is $g(-2) = -8$ and the absolute maximum value is $g(6) = 8$.

- (d) $g(x) \leq 0$ for $-4 \leq x \leq 2$ and $10 \leq x \leq 12$.

1 : $g'(x) = f(x)$ in (a), (b), (c), or (d)

1 : answer with justification

1 : answer with justification

4 : $\left\{ \begin{array}{l} 1 : \text{considers } x = -2 \text{ and } x = 6 \\ \quad \text{as candidates} \\ 1 : \text{considers } x = -4 \text{ and } x = 12 \\ 2 : \text{answers with justification} \end{array} \right.$

2 : intervals

2016 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = x^2 - \frac{1}{2}y$.

(a) Find $\frac{d^2y}{dx^2}$ in terms of x and y .

(b) Let $y = f(x)$ be the particular solution to the given differential equation whose graph passes through the point $(-2, 8)$. Does the graph of f have a relative minimum, a relative maximum, or neither at the point $(-2, 8)$? Justify your answer.

(c) Let $y = g(x)$ be the particular solution to the given differential equation with $g(-1) = 2$. Find

$$\lim_{x \rightarrow -1} \left(\frac{g(x) - 2}{3(x+1)^2} \right).$$
 Show the work that leads to your answer.

(d) Let $y = h(x)$ be the particular solution to the given differential equation with $h(0) = 2$. Use Euler's method, starting at $x = 0$ with two steps of equal size, to approximate $h(1)$.

$$(a) \quad \frac{d^2y}{dx^2} = 2x - \frac{1}{2} \frac{dy}{dx} = 2x - \frac{1}{2} \left(x^2 - \frac{1}{2}y \right)$$

$$2 : \frac{d^2y}{dx^2} \text{ in terms of } x \text{ and } y$$

$$(b) \quad \left. \frac{dy}{dx} \right|_{(x,y)=(-2,8)} = (-2)^2 - \frac{1}{2} \cdot 8 = 0$$

2 : conclusion with justification

$$\left. \frac{d^2y}{dx^2} \right|_{(x,y)=(-2,8)} = 2(-2) - \frac{1}{2} \left((-2)^2 - \frac{1}{2} \cdot 8 \right) = -4 < 0$$

Thus, the graph of f has a relative maximum at the point $(-2, 8)$.

$$(c) \quad \lim_{x \rightarrow -1} (g(x) - 2) = 0 \text{ and } \lim_{x \rightarrow -1} 3(x+1)^2 = 0$$

$$3 : \begin{cases} 2 : \text{L'Hospital's Rule} \\ 1 : \text{answer} \end{cases}$$

Using L'Hospital's Rule,

$$\lim_{x \rightarrow -1} \left(\frac{g(x) - 2}{3(x+1)^2} \right) = \lim_{x \rightarrow -1} \left(\frac{g'(x)}{6(x+1)} \right)$$

$$\lim_{x \rightarrow -1} g'(x) = 0 \text{ and } \lim_{x \rightarrow -1} 6(x+1) = 0$$

Using L'Hospital's Rule,

$$\lim_{x \rightarrow -1} \left(\frac{g'(x)}{6(x+1)} \right) = \lim_{x \rightarrow -1} \left(\frac{g''(x)}{6} \right) = \frac{-2}{6} = -\frac{1}{3}$$


$$(d) \quad h\left(\frac{1}{2}\right) \approx h(0) + h'(0) \cdot \frac{1}{2} = 2 + (-1) \cdot \frac{1}{2} = \frac{3}{2}$$

$$h(1) \approx h\left(\frac{1}{2}\right) + h'\left(\frac{1}{2}\right) \cdot \frac{1}{2} \approx \frac{3}{2} + \left(-\frac{1}{2}\right) \cdot \frac{1}{2} = \frac{5}{4}$$

$$2 : \begin{cases} 1 : \text{Euler's method} \\ 1 : \text{approximation} \end{cases}$$

2016 SCORING GUIDELINES

Question 5



The inside of a funnel of height 10 inches has circular cross sections, as shown in the figure above. At height h , the radius of the funnel is given by $r = \frac{1}{20}(3 + h^2)$, where $0 \leq h \leq 10$. The units of r and h are inches.

- (a) Find the average value of the radius of the funnel.
- (b) Find the volume of the funnel.
- (c) The funnel contains liquid that is draining from the bottom. At the instant when the height of the liquid is $h = 3$ inches, the radius of the surface of the liquid is decreasing at a rate of $\frac{1}{5}$ inch per second. At this instant, what is the rate of change of the height of the liquid with respect to time?

$$\begin{aligned} \text{(a) Average radius} &= \frac{1}{10} \int_0^{10} \frac{1}{20}(3 + h^2) dh = \frac{1}{200} \left[3h + \frac{h^3}{3} \right]_0^{10} \\ &= \frac{1}{200} \left(\left(30 + \frac{1000}{3} \right) - 0 \right) = \frac{109}{60} \text{ in} \end{aligned}$$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b) Volume} &= \pi \int_0^{10} \left(\left(\frac{1}{20} \right) (3 + h^2) \right)^2 dh = \frac{\pi}{400} \int_0^{10} (9 + 6h^2 + h^4) dh \\ &= \frac{\pi}{400} \left[9h + 2h^3 + \frac{h^5}{5} \right]_0^{10} \\ &= \frac{\pi}{400} \left(\left(90 + 2000 + \frac{100000}{5} \right) - 0 \right) = \frac{2209\pi}{40} \text{ in}^3 \end{aligned}$$

3 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(c) } \frac{dr}{dt} &= \frac{1}{20}(2h) \frac{dh}{dt} \\ -\frac{1}{5} &= \frac{3}{10} \frac{dh}{dt} \\ \frac{dh}{dt} &= -\frac{1}{5} \cdot \frac{10}{3} = -\frac{2}{3} \text{ in/sec} \end{aligned}$$

3 : $\begin{cases} 2 : \text{chain rule} \\ 1 : \text{answer} \end{cases}$

2016 SCORING GUIDELINES

Question 6

The function f has a Taylor series about $x = 1$ that converges to $f(x)$ for all x in the interval of convergence.

It is known that $f(1) = 1$, $f'(1) = -\frac{1}{2}$, and the n th derivative of f at $x = 1$ is given by

$$f^{(n)}(1) = (-1)^n \frac{(n-1)!}{2^n} \text{ for } n \geq 2.$$

- (a) Write the first four nonzero terms and the general term of the Taylor series for f about $x = 1$.
- (b) The Taylor series for f about $x = 1$ has a radius of convergence of 2. Find the interval of convergence. Show the work that leads to your answer.
- (c) The Taylor series for f about $x = 1$ can be used to represent $f(1.2)$ as an alternating series. Use the first three nonzero terms of the alternating series to approximate $f(1.2)$.
- (d) Show that the approximation found in part (c) is within 0.001 of the exact value of $f(1.2)$.

$$\begin{aligned} \text{(a)} \quad f(1) = 1, \quad f'(1) = -\frac{1}{2}, \quad f''(1) = \frac{1}{2^2}, \quad f'''(1) = -\frac{2}{2^3} \\ f(x) = 1 - \frac{1}{2}(x-1) + \frac{1}{2^2 \cdot 2}(x-1)^2 - \frac{1}{2^3 \cdot 3}(x-1)^3 + \cdots \\ + \frac{(-1)^n}{2^n \cdot n}(x-1)^n + \cdots \end{aligned}$$

$$4 : \begin{cases} 1 : \text{first two terms} \\ 1 : \text{third term} \\ 1 : \text{fourth term} \\ 1 : \text{general term} \end{cases}$$

- (b) $R = 2$. The series converges on the interval $(-1, 3)$.

When $x = -1$, the series is $1 + 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots$.

Since the harmonic series diverges, this series diverges.

When $x = 3$, the series is $1 - 1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \cdots$.

Since the alternating harmonic series converges, this series converges.

Therefore, the interval of convergence is $-1 < x \leq 3$.

$$2 : \begin{cases} 1 : \text{identifies both endpoints} \\ 1 : \text{analysis and interval of convergence} \end{cases}$$

$$\text{(c)} \quad f(1.2) \approx 1 - \frac{1}{2}(0.2) + \frac{1}{8}(0.2)^2 = 1 - 0.1 + 0.005 = 0.905$$

1 : approximation

- (d) The series for $f(1.2)$ alternates with terms that decrease in magnitude to 0.

$$2 : \begin{cases} 1 : \text{error form} \\ 1 : \text{analysis} \end{cases}$$

$$|f(1.2) - T_2(1.2)| \leq \left| \frac{-1}{2^3 \cdot 3}(0.2)^3 \right| = \frac{1}{3000} \leq 0.001$$

AP[®] Calculus BC

2015 Scoring Guidelines

© 2015 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

0

2015 SCORING GUIDELINES

Question 1

The rate at which rainwater flows into a drainpipe is modeled by the function R , where $R(t) = 20\sin\left(\frac{t^2}{35}\right)$ cubic feet per hour, t is measured in hours, and $0 \leq t \leq 8$. The pipe is partially blocked, allowing water to drain out the other end of the pipe at a rate modeled by $D(t) = -0.04t^3 + 0.4t^2 + 0.96t$ cubic feet per hour, for $0 \leq t \leq 8$. There are 30 cubic feet of water in the pipe at time $t = 0$.

- (a) How many cubic feet of rainwater flow into the pipe during the 8-hour time interval $0 \leq t \leq 8$?
- (b) Is the amount of water in the pipe increasing or decreasing at time $t = 3$ hours? Give a reason for your answer.
- (c) At what time t , $0 \leq t \leq 8$, is the amount of water in the pipe at a minimum? Justify your answer.
- (d) The pipe can hold 50 cubic feet of water before overflowing. For $t > 8$, water continues to flow into and out of the pipe at the given rates until the pipe begins to overflow. Write, but do not solve, an equation involving one or more integrals that gives the time w when the pipe will begin to overflow.

(a) $\int_0^8 R(t) dt = 76.570$

2 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b) $R(3) - D(3) = -0.313632 < 0$
 Since $R(3) < D(3)$, the amount of water in the pipe is decreasing at time $t = 3$ hours.

2 : $\begin{cases} 1 : \text{considers } R(3) \text{ and } D(3) \\ 1 : \text{answer and reason} \end{cases}$

(c) The amount of water in the pipe at time t , $0 \leq t \leq 8$, is $30 + \int_0^t [R(x) - D(x)] dx$.

3 : $\begin{cases} 1 : \text{considers } R(t) - D(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

$R(t) - D(t) = 0 \Rightarrow t = 0, 3.271658$

t	Amount of water in the pipe
0	30
3.271658	27.964561
8	48.543686

The amount of water in the pipe is a minimum at time $t = 3.272$ (or 3.271) hours.

(d) $30 + \int_0^w [R(t) - D(t)] dt = 50$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{equation} \end{cases}$

2015 SCORING GUIDELINES

Question 2

At time $t \geq 0$, a particle moving along a curve in the xy -plane has position $(x(t), y(t))$ with velocity vector $v(t) = (\cos(t^2), e^{0.5t})$. At $t = 1$, the particle is at the point $(3, 5)$.

- (a) Find the x -coordinate of the position of the particle at time $t = 2$.
- (b) For $0 < t < 1$, there is a point on the curve at which the line tangent to the curve has a slope of 2. At what time is the object at that point?
- (c) Find the time at which the speed of the particle is 3.
- (d) Find the total distance traveled by the particle from time $t = 0$ to time $t = 1$.

(a) $x(2) = 3 + \int_1^2 \cos(t^2) dt = 2.557$ (or 2.556)

$$3 : \begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$$

(b) $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{e^{0.5t}}{\cos(t^2)}$

$$\frac{e^{0.5t}}{\cos(t^2)} = 2$$

$$t = 0.840$$

$$2 : \begin{cases} 1 : \text{slope in terms of } t \\ 1 : \text{answer} \end{cases}$$

(c) Speed = $\sqrt{\cos^2(t^2) + e^t}$

$$\sqrt{\cos^2(t^2) + e^t} = 3$$

$$t = 2.196$$
 (or 2.195)

$$2 : \begin{cases} 1 : \text{speed in terms of } t \\ 1 : \text{answer} \end{cases}$$

(d) Distance = $\int_0^1 \sqrt{\cos^2(t^2) + e^t} dt = 1.595$ (or 1.594)

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

2015 SCORING GUIDELINES

Question 3

t (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	-220	150

Johanna jogs along a straight path. For $0 \leq t \leq 40$, Johanna's velocity is given by a differentiable function v . Selected values of $v(t)$, where t is measured in minutes and $v(t)$ is measured in meters per minute, are given in the table above.

- (a) Use the data in the table to estimate the value of $v'(16)$.
- (b) Using correct units, explain the meaning of the definite integral $\int_0^{40} |v(t)| dt$ in the context of the problem.

Approximate the value of $\int_0^{40} |v(t)| dt$ using a right Riemann sum with the four subintervals indicated in the table.

- (c) Bob is riding his bicycle along the same path. For $0 \leq t \leq 10$, Bob's velocity is modeled by $B(t) = t^3 - 6t^2 + 300$, where t is measured in minutes and $B(t)$ is measured in meters per minute.

Find Bob's acceleration at time $t = 5$.

- (d) Based on the model B from part (c), find Bob's average velocity during the interval $0 \leq t \leq 10$.

(a) $v'(16) \approx \frac{240 - 200}{20 - 12} = 5 \text{ meters/min}^2$

1 : approximation

- (b) $\int_0^{40} |v(t)| dt$ is the total distance Johanna jogs, in meters, over the time interval $0 \leq t \leq 40$ minutes.

3 : $\begin{cases} 1 : \text{explanation} \\ 1 : \text{right Riemann sum} \\ 1 : \text{approximation} \end{cases}$

$$\begin{aligned} \int_0^{40} |v(t)| dt &\approx 12 \cdot |v(12)| + 8 \cdot |v(20)| + 4 \cdot |v(24)| + 16 \cdot |v(40)| \\ &= 12 \cdot 200 + 8 \cdot 240 + 4 \cdot 220 + 16 \cdot 150 \\ &= 2400 + 1920 + 880 + 2400 \\ &= 7600 \text{ meters} \end{aligned}$$

- (c) Bob's acceleration is $B'(t) = 3t^2 - 12t$.
 $B'(5) = 3(25) - 12(5) = 15 \text{ meters/min}^2$

2 : $\begin{cases} 1 : \text{uses } B'(t) \\ 1 : \text{answer} \end{cases}$

(d) Avg vel = $\frac{1}{10} \int_0^{10} (t^3 - 6t^2 + 300) dt$

$$\begin{aligned} &= \frac{1}{10} \left[\frac{t^4}{4} - 2t^3 + 300t \right]_0^{10} \\ &= \frac{1}{10} \left[\frac{10000}{4} - 2000 + 3000 \right] = 350 \text{ meters/min} \end{aligned}$$


3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = 2x - y$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.
- (b) Find $\frac{d^2y}{dx^2}$ in terms of x and y . Determine the concavity of all solution curves for the given differential equation in Quadrant II. Give a reason for your answer.
- (c) Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(2) = 3$. Does f have a relative minimum, a relative maximum, or neither at $x = 2$? Justify your answer.
- (d) Find the values of the constants m and b for which $y = mx + b$ is a solution to the differential equation.



$$2 : \begin{cases} 1 : \text{slopes where } x = 0 \\ 1 : \text{slopes where } x = 1 \end{cases}$$

(b) $\frac{d^2y}{dx^2} = 2 - \frac{dy}{dx} = 2 - (2x - y) = 2 - 2x + y$

In Quadrant II, $x < 0$ and $y > 0$, so $2 - 2x + y > 0$.
Therefore, all solution curves are concave up in Quadrant II.

$$2 : \begin{cases} 1 : \frac{d^2y}{dx^2} \\ 1 : \text{concave up with reason} \end{cases}$$

(c) $\left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} = 2(2) - 3 = 1 \neq 0$

Therefore, f has neither a relative minimum nor a relative maximum at $x = 2$.

$$2 : \begin{cases} 1 : \text{considers } \left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} \\ 1 : \text{conclusion with justification} \end{cases}$$

(d) $y = mx + b \Rightarrow \frac{dy}{dx} = \frac{d}{dx}(mx + b) = m$

$$2x - y = m$$

$$2x - (mx + b) = m$$

$$(2 - m)x - (m + b) = 0$$

$$2 - m = 0 \Rightarrow m = 2$$

$$b = -m \Rightarrow b = -2$$

$$3 : \begin{cases} 1 : \frac{d}{dx}(mx + b) = m \\ 1 : 2x - y = m \\ 1 : \text{answer} \end{cases}$$

2015 SCORING GUIDELINES

Question 5

Consider the function $f(x) = \frac{1}{x^2 - kx}$, where k is a nonzero constant. The derivative of f is given by

$$f'(x) = \frac{k - 2x}{(x^2 - kx)^2}.$$

- (a) Let $k = 3$, so that $f(x) = \frac{1}{x^2 - 3x}$. Write an equation for the line tangent to the graph of f at the point whose x -coordinate is 4.
- (b) Let $k = 4$, so that $f(x) = \frac{1}{x^2 - 4x}$. Determine whether f has a relative minimum, a relative maximum, or neither at $x = 2$. Justify your answer.
- (c) Find the value of k for which f has a critical point at $x = -5$.
- (d) Let $k = 6$, so that $f(x) = \frac{1}{x^2 - 6x}$. Find the partial fraction decomposition for the function f .

Find $\int f(x) dx$.

$$(a) \quad f(4) = \frac{1}{4^2 - 3 \cdot 4} = \frac{1}{4} \quad f'(4) = \frac{3 - 2 \cdot 4}{(4^2 - 3 \cdot 4)^2} = -\frac{5}{16}$$

An equation for the line tangent to the graph of f at the point whose x -coordinate is 4 is $y = -\frac{5}{16}(x - 4) + \frac{1}{4}$.

$$(b) \quad f'(x) = \frac{4 - 2x}{(x^2 - 4x)^2} \quad f'(2) = \frac{4 - 2 \cdot 2}{(2^2 - 4 \cdot 2)^2} = 0$$

$f'(x)$ changes sign from positive to negative at $x = 2$.
Therefore, f has a relative maximum at $x = 2$.

$$(c) \quad f'(-5) = \frac{k - 2 \cdot (-5)}{((-5)^2 - k \cdot (-5))^2} = 0 \Rightarrow k = -10$$

$$(d) \quad \frac{1}{x^2 - 6x} = \frac{1}{x(x - 6)} = \frac{A}{x} + \frac{B}{x - 6} \Rightarrow 1 = A(x - 6) + Bx$$

$$x = 0 \Rightarrow 1 = A \cdot (-6) \Rightarrow A = -\frac{1}{6}$$

$$x = 6 \Rightarrow 1 = B \cdot (6) \Rightarrow B = \frac{1}{6}$$

$$\frac{1}{x(x - 6)} = \frac{-1/6}{x} + \frac{1/6}{x - 6}$$

$$\int f(x) dx = \int \left(\frac{-1/6}{x} + \frac{1/6}{x - 6} \right) dx$$

$$= -\frac{1}{6} \ln|x| + \frac{1}{6} \ln|x - 6| + C = \frac{1}{6} \ln \left| \frac{x - 6}{x} \right| + C$$

2 : $\begin{cases} 1 : \text{slope} \\ 1 : \text{tangent line equation} \end{cases}$

2 : $\begin{cases} 1 : \text{considers } f'(2) \\ 1 : \text{answer with justification} \end{cases}$

1 : answer

4 : $\begin{cases} 2 : \text{partial fraction decomposition} \\ 2 : \text{general antiderivative} \end{cases}$

2015 SCORING GUIDELINES

Question 6

The Maclaurin series for a function f is given by $\sum_{n=1}^{\infty} \frac{(-3)^{n-1}}{n} x^n = x - \frac{3}{2}x^2 + 3x^3 - \cdots + \frac{(-3)^{n-1}}{n} x^n + \cdots$ and converges to $f(x)$ for $|x| < R$, where R is the radius of convergence of the Maclaurin series.

- (a) Use the ratio test to find R .
- (b) Write the first four nonzero terms of the Maclaurin series for f' , the derivative of f . Express f' as a rational function for $|x| < R$.
- (c) Write the first four nonzero terms of the Maclaurin series for e^x . Use the Maclaurin series for e^x to write the third-degree Taylor polynomial for $g(x) = e^x f(x)$ about $x = 0$.

- (a) Let a_n be the n th term of the Maclaurin series.

$$\frac{a_{n+1}}{a_n} = \frac{(-3)^n x^{n+1}}{n+1} \cdot \frac{n}{(-3)^{n-1} x^n} = \frac{-3n}{n+1} \cdot x$$

$$\lim_{n \rightarrow \infty} \left| \frac{-3n}{n+1} \cdot x \right| = 3|x|$$

$$3|x| < 1 \Rightarrow |x| < \frac{1}{3}$$

The radius of convergence is $R = \frac{1}{3}$.

- (b) The first four nonzero terms of the Maclaurin series for f' are $1 - 3x + 9x^2 - 27x^3$.

$$f'(x) = \frac{1}{1 - (-3x)} = \frac{1}{1 + 3x}$$

- (c) The first four nonzero terms of the Maclaurin series for e^x are

$$1 + x + \frac{x^2}{2!} + \frac{x^3}{3!}.$$

The product of the Maclaurin series for e^x and the Maclaurin series for f is

$$\begin{aligned} & \left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots \right) \left(x - \frac{3}{2}x^2 + 3x^3 - \cdots \right) \\ &= x - \frac{1}{2}x^2 + 2x^3 + \cdots \end{aligned}$$

The third-degree Taylor polynomial for $g(x) = e^x f(x)$

about $x = 0$ is $T_3(x) = x - \frac{1}{2}x^2 + 2x^3$.

$$3 : \begin{cases} 1 : \text{sets up ratio} \\ 1 : \text{computes limit of ratio} \\ 1 : \text{determines radius of convergence} \end{cases}$$

$$3 : \begin{cases} 2 : \text{first four nonzero terms} \\ 1 : \text{rational function} \end{cases}$$

$$3 : \begin{cases} 1 : \text{first four nonzero terms} \\ \quad \text{of the Maclaurin series for } e^x \\ 2 : \text{Taylor polynomial} \end{cases}$$