

**Part A (AB or BC): Graphing calculator required****Question 1****9 points****General Scoring Notes**

- The model solution is presented using standard mathematical notation.
- Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be accurate to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

An invasive species of plant appears in a fruit grove at time  $t = 0$  and begins to spread. The function  $C$  defined by  $C(t) = 7.6 \arctan(0.2t)$  models the number of acres in the fruit grove affected by the species  $t$  weeks after the species appears. It can be shown that  $C'(t) = \frac{38}{25 + t^2}$ .

(Note: Your calculator should be in radian mode.)

|          | Model Solution                                                                                                                                   | Scoring                                   |
|----------|--------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------|
| <b>A</b> | Find the average number of acres affected by the invasive species from time $t = 0$ to time $t = 4$ weeks. Show the setup for your calculations. |                                           |
|          | $\frac{1}{4 - 0} \int_0^4 C(t) dt$                                                                                                               | Average value formula <b>Point 1 (P1)</b> |
|          | $= \frac{1}{4}(11.112896) = 2.778224$                                                                                                            | Answer <b>Point 2 (P2)</b>                |
|          | From time $t = 0$ to $t = 4$ weeks, the average number of acres affected by the invasive species was 2.778 acres.                                |                                           |

**Scoring Notes for Part A**

- P1** is earned for the correct integral, with or without the differential, along with evidence of division by 4. In the presence of the correct integral, the correct answer will suffice as evidence of division by 4. These may appear all in one step, as in the model solution, or in multiple steps.
- P2** is earned for the correct answer, with or without supporting work. A reported answer should be accurate to three places after the decimal point, rounded or truncated. An inappropriately rounded answer does not earn the point.
- Incorrect or unclear communication between the correct integral and the correct answer is treated as scratch work and is not considered in scoring. For example:
  - $\int_0^4 C(t) dt = 11.112896$  so the average velocity is 2.778224.  
Note: This response earns **P1** for the correct integral with the correct answer as evidence of division by 4. It also earns **P2** for the correct answer.
  - $\int_0^4 C(t) dt = \frac{11.112896}{4} = 2.778224$   
Note: This response earns **P1** for the correct integral with the correct answer as evidence of division by 4. It also earns **P2** for the correct answer. (In this instance, incorrect linkage is not considered in scoring.)
  - $\int_0^4 C(t) dt = 2.778224$   
Note: This response earns **P1** for the correct integral with the correct answer as evidence of division by 4. It also earns **P2** for the correct answer. (In this instance, incorrect linkage is not considered in scoring.)
- Note that the values  $\frac{1}{4}(11.112)$  and  $\frac{1}{4}(11.113)$  are accurate to three digits after the decimal and therefore earn **P2**.

- B** Find the time  $t$  when the instantaneous rate of change of  $C$  equals the average rate of change of  $C$  over the time interval  $0 \leq t \leq 4$ . Show the setup for your calculations.

$$\frac{C(4) - C(0)}{4 - 0} = 1.282008$$

Uses average rate of change **Point 3 (P3)**

$$C'(t) = \frac{38}{25 + t^2} = 1.282008 \Rightarrow t = 2.154298$$

Answer with supporting work **Point 4 (P4)**

The instantaneous rate of change of  $C$  equals the average rate of change of  $C$  over the interval  $0 \leq t \leq 4$  at time  $t = 2.154$ .

#### Scoring Notes for Part B

- P3** may be earned by presenting the expression or value for the average rate of change. Note that because  $C(0) = 0$  and the interval is  $0 \leq t \leq 4$ , any of the following will earn **P3**:  $\frac{\int_0^4 C'(t) dt}{4}$ ,  $\frac{C(4) - C(0)}{4 - 0}$ ,  $\frac{C(4)}{4}$ ,  $\frac{5.128 - 0}{4 - 0}$ ,  $\frac{5.128}{4}$ , or 1.282. However, neither **P3** nor **P4** is earned by just presenting  $t = 1.282$ .
  - P4** is earned for the correct answer supported by the appropriate equation. A reported answer should be accurate to three places after the decimal point, rounded or truncated. An inappropriately rounded answer does not earn the point, unless an earlier point was not earned due to inappropriate rounding.
- The following response, for example, earns both **P3** and **P4**:  $C'(t) = \frac{C(4) - C(0)}{4}$  when  $t = 2.154$ .

- C** Assume that the invasive species continues to spread according to the given model for all times  $t > 0$ . Write a limit expression that describes the end behavior of the rate of change in the number of acres affected by the species. Evaluate this limit expression.

$$\lim_{t \rightarrow \infty} C'(t) = \lim_{t \rightarrow \infty} \frac{38}{25 + t^2}$$

Limit expression **Point 5 (P5)**

$$= 0$$

Value **Point 6 (P6)**

#### Scoring Notes for Part C

- P5** can be earned for either  $\lim_{t \rightarrow \infty} C'(t)$  or  $\lim_{t \rightarrow \infty} C(t)$ .
- A response that includes  $\lim_{t \rightarrow \infty} C(t)$  is not eligible to earn **P6**.
- For **P6**, arithmetic with infinity, e.g.,  $\frac{38}{25 + \infty^2} = 0$ , will be considered as scratch work and will not be considered in scoring.

- D** At time  $t = 4$  weeks after the invasive species appears in the fruit grove, measures are taken to counter the spread of the species. The function  $A$ , defined by  $A(t) = C(t) - \int_4^t 0.1 \cdot \ln(x) dx$ , models the number of acres affected by the species over the time interval  $4 \leq t \leq 36$ . At what time  $t$ , for  $4 \leq t \leq 36$ , does  $A$  attain its maximum value? Justify your answer.

$$A'(t) = C'(t) - 0.1 \cdot \ln t$$

Considers  $A'(t) = 0$  **Point 7 (P7)**

For  $4 \leq t \leq 36$ , the maximum value of  $A(t)$  occurs when  $A'(t) = 0$  or at an endpoint.

$$A'(t) = C'(t) - 0.1 \cdot \ln t = 0 \Rightarrow C'(t) = 0.1 \cdot \ln t$$

$$\Rightarrow t = 11.441700$$

Justification **Point 8 (P8)**

| $t$       | $A(t)$   |
|-----------|----------|
| 4         | 5.128031 |
| 11.441700 | 7.316978 |
| 36        | 1.743056 |

Therefore, the number of acres affected by the species is a maximum at time  $t = 11.442$  (or 11.441) weeks.

Answer with supporting work **Point 9 (P9)**


#### Scoring Notes for Part D

- P7** is earned for considering  $A'(t) = 0$ ,  $C'(t) - 0.1 \cdot \ln t = 0$ , or  $C'(t) = 0.1 \cdot \ln t$ . **P7** is not earned by just presenting  $t = 11.441700$ . A response that discusses the sign of  $A'(t)$  changing or uses the phrase “critical points of  $A$ ” also earns **P7**.
- To earn **P8** using a candidates test, a response must make a global argument by correctly evaluating  $A(t)$  at  $t = 4$ ,  $t = 11.441700$ , and  $t = 36$ . The evaluations must be correct to the first digit after the decimal, rounded or truncated.
- Alternate justifications:
  - $A'(t) > 0$  for  $4 < t < 11.442$ , and  $A'(t) < 0$  for  $11.442 < t < 36$ . Therefore,  $t = 11.442$  is the location of the absolute maximum for  $A$  on the interval  $4 \leq t \leq 36$ .
  - Because  $A'(t)$  changes sign from positive to negative at  $t = 11.442$  (this might be presented as “ $A'(t) > 0$  for  $t < 11.442$ , and  $A'(t) < 0$  for  $t > 11.442$ ”), it is the location of a relative maximum for  $A$ . And because  $t = 11.442$  is the only critical point of  $A$  in the interval  $4 \leq t \leq 36$ , it is the location of the absolute maximum for  $A$  on the interval.
- A response that presents a local argument (such as a First Derivative Test or a Second Derivative Test) or an incorrect global argument does not earn **P8** but is eligible for **P9** with the correct answer. A reported answer should be accurate to three places after the decimal point, rounded or truncated. An inappropriately rounded answer does not earn the point, unless an earlier point was not earned due to inappropriate rounding.

**Part B (AB or BC): Graphing calculator not allowed****Question 4****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.



The graph of the differentiable function  $f$ , shown for  $-6 \leq x \leq 7$ , has a horizontal tangent at  $x = -2$  and is linear for  $0 \leq x \leq 7$ . Let  $R$  be the region in the second quadrant bounded by the graph of  $f$ , the vertical line  $x = -6$ , and the  $x$ - and  $y$ -axes. Region  $R$  has area 12.

**Model Solution****Scoring**

- (a) The function  $g$  is defined by  $g(x) = \int_0^x f(t) dt$ . Find the values of  $g(-6)$ ,  $g(4)$ , and  $g(6)$ .

|                                                                                           |         |                |
|-------------------------------------------------------------------------------------------|---------|----------------|
| $g(-6) = \int_0^{-6} f(t) dt = -\int_{-6}^0 f(t) dt = -12$                                | $g(-6)$ | <b>1 point</b> |
| $g(4) = \int_0^4 f(t) dt = \frac{1}{2} \cdot 4 \cdot 2 = 4$                               | $g(4)$  | <b>1 point</b> |
| $g(6) = \int_0^6 f(t) dt = \frac{1}{2} \cdot 4 \cdot 2 - \frac{1}{2} \cdot 2 \cdot 1 = 3$ | $g(6)$  | <b>1 point</b> |

**Scoring notes:**

- Supporting work is not required for any of these values. However, any supporting work that is shown must be correct to earn the corresponding point.
- Special case: A response that explicitly presents  $g(x) = \int_{-6}^x f(t) dt$  does not earn the first point it would have otherwise earned. The response is eligible for all subsequent points for correct answers, or for consistent answers with supporting work.
  - Note:  $\int_{-6}^{-6} f(t) dt = 0$ ,  $\int_{-6}^4 f(t) dt = 16$ ,  $\int_{-6}^6 f(t) dt = 15$
- Labeled values may be presented in any order. Unlabeled values are read from left to right and from top to bottom as  $g(-6)$ ,  $g(4)$ , and  $g(6)$ , respectively. A response that presents only 1 or 2 values must label them to earn any points.

**Total for part (a) 3 points**

- (b) For the function  $g$  defined in part (a), find all values of  $x$  in the interval  $0 \leq x \leq 6$  at which the graph of  $g$  has a critical point. Give a reason for your answer.

|                                      |                                 |                |
|--------------------------------------|---------------------------------|----------------|
| $g'(x) = f(x)$                       | Fundamental Theorem of Calculus | <b>1 point</b> |
| $g'(x) = f(x) = 0 \Rightarrow x = 4$ | Answer with reason              | <b>1 point</b> |

Therefore, the graph of  $g$  has a critical point at  $x = 4$ .

**Scoring notes:**

- The first point is earned for explicitly making the connection  $g' = f$  in this part.
  - A response that writes  $g'' = f'$  earns the first point but can only earn the second point by reasoning from  $f = 0$ .
- A response that does not earn the first point is eligible to earn the second point with an implied application of the FTC (e.g., "Because  $g'(4) = 0$ ,  $x = 4$  is a critical point").
- A response that reports any additional critical points in  $0 < x < 6$  does not earn the second point.
  - Any presented critical point outside the interval  $0 < x < 6$  will not affect scoring.

**Total for part (b) 2 points**

- (c) The function  $h$  is defined by  $h(x) = \int_{-6}^x f'(t) dt$ . Find the values of  $h(6)$ ,  $h'(6)$ , and  $h''(6)$ . Show the work that leads to your answers.

|                                                                |                                      |                |
|----------------------------------------------------------------|--------------------------------------|----------------|
| $h(6) = \int_{-6}^6 f'(t) dt = f(6) - f(-6) = -1 - 0.5 = -1.5$ | Uses Fundamental Theorem of Calculus | <b>1 point</b> |
|                                                                | $h(6)$ with supporting work          | <b>1 point</b> |
| $h'(x) = f'(x)$ , so $h'(6) = f'(6) = -\frac{1}{2}$ .          | $h'(6)$                              | <b>1 point</b> |
| $h''(x) = f''(x)$ , so $h''(6) = f''(6) = 0$ .                 | $h''(6)$                             | <b>1 point</b> |

**Scoring notes:**

- Labeled values may be presented in any order.
- Unlabeled values are read from left to right and from top to bottom as  $h(6)$ ,  $h'(6)$ , and  $h''(6)$ , respectively. A response that presents only 1 or 2 values must label them in order to earn any points.
- A response of  $h(6) = -1.5$  does not earn either of the first 2 points. A response of  $h(6) = f(6) - f(-6)$  earns the first point but not yet the second point.
- A response of  $h(6) = -1 - 0.5$  is the minimum work required to earn both of the first 2 points.
- To earn the third point a response must state either  $h'(x) = f'(x)$  or  $h'(6) = f'(6)$ , and provide an answer of  $-\frac{1}{2}$ .
- The fourth point is earned for a response of  $h''(6) = 0$ , with or without supporting work.
- A response that has one or more linkage errors does not earn the first point it would have otherwise earned. For example,  $h'(x) = f'(6) = -\frac{1}{2}$  does not earn the third point but is eligible for the fourth point even in the presence of another linkage error, such as  $h''(x) = f''(6) = 0$ .

**Total for part (c) 4 points****Total for question 4 9 points****Part B (BC): Graphing calculator not allowed****Question 5****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

|         |   |       |        |
|---------|---|-------|--------|
| $x$     | 0 | $\pi$ | $2\pi$ |
| $f'(x)$ | 5 | 6     | 0      |

The function  $f$  is twice differentiable for all  $x$  with  $f(0) = 0$ . Values of  $f'$ , the derivative of  $f$ , are given in the table for selected values of  $x$ .

| Model Solution                                                                                                                                                         | Scoring                                        |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------|
| (a) For $x \geq 0$ , the function $h$ is defined by $h(x) = \int_0^x \sqrt{1 + (f'(t))^2} dt$ . Find the value of $h'(\pi)$ . Show the work that leads to your answer. |                                                |
| $h'(x) = \sqrt{1 + (f'(x))^2}$                                                                                                                                         | Fundamental Theorem of Calculus <b>1 point</b> |
| $h'(\pi) = \sqrt{1 + (f'(\pi))^2} = \sqrt{1 + 6^2} = \sqrt{37}$                                                                                                        | Answer <b>1 point</b>                          |

**Scoring notes:**

- A response of  $\sqrt{1 + (f'(\pi))^2}$  earns the first point.
- A response of  $\sqrt{1 + 6^2}$  alone earns both points.
- A response such as  $h'(x) = \sqrt{1 + (f'(x))^2} = \sqrt{37}$ , that equates a variable expression to a numeric value, earns at most 1 of the 2 points.
- A response that equates  $h'(x)$  or  $h'(\pi)$  to a derivative of a constant, such as

$$h'(x) = \frac{d}{dx} \int_0^x \sqrt{1 + (f'(t))^2} dt, \text{ earns at most 1 of the 2 points.}$$

**Total for part (a) 2 points**

- (b) What information does  $\int_0^\pi \sqrt{1 + (f'(x))^2} dx$  provide about the graph of  $f$ ?

|                                                                                            |                     |                |
|--------------------------------------------------------------------------------------------|---------------------|----------------|
| $\int_0^\pi \sqrt{1 + (f'(x))^2} dx$ is the arc length of the graph of $f$ on $[0, \pi]$ . | Arc length of $f$   | <b>1 point</b> |
|                                                                                            | Interval $[0, \pi]$ | <b>1 point</b> |

**Scoring notes:**

- A response of “arc length” or “length” earns the first point. Such a response does not need to reference  $f$ . However, if the response references a different function, the response does not earn the first point and is eligible to earn the second point.
- A response referring to distance explicitly connected to the graph or  $f$  (or equivalent) earns the first point. For example, a response of “distance along the curve” or “distance traveled by a particle moving along  $f$ ” earns the first point and is eligible to earn the second point.
- A response referring to distance that is not explicitly connected to the graph of  $f$  does not earn the first point but is eligible to earn the second point. For example, a response of “distance” or “distance traveled” does not earn the first point but is eligible to earn the second point.
- To earn the second point a response must connect the interval  $[0, \pi]$  to arc length, length, or distance.

**Total for part (b) 2 points**

- (c) Use Euler’s method, starting at  $x = 0$  with two steps of equal size, to approximate  $f(2\pi)$ . Show the computations that lead to your answer.

|                                                     |                |                |
|-----------------------------------------------------|----------------|----------------|
| $f(\pi) \approx f(0) + \pi f'(0) = 0 + 5\pi = 5\pi$ | Euler’s method | <b>1 point</b> |
| $f(2\pi) \approx f(\pi) + \pi f'(\pi)$              |                |                |
| $\approx 5\pi + 6\pi = 11\pi$                       | Answer         | <b>1 point</b> |

**Scoring notes:**

- To earn the first point a response must demonstrate two Euler’s steps, with use of the correct expression for  $\frac{dy}{dx}$ , and at most one error. If there is an error, the second point is not earned.
- In order to earn the first point, a response that presents a single error in computing the approximation of  $f(\pi)$  must import the incorrect value in computing the approximation of  $f(2\pi)$ .
- The two Euler’s steps may be explicit expressions or may be presented in a table. For example:

| $x$    | $y$     | $\frac{dy}{dx} \cdot \Delta x$ (or $\frac{dy}{dx} \cdot \pi$ ) |
|--------|---------|----------------------------------------------------------------|
| 0      | 0       | $5\pi$                                                         |
| $\pi$  | $5\pi$  | $6\pi$                                                         |
| $2\pi$ | $11\pi$ |                                                                |

- In the presence of a correct answer, a table does not need to be labeled in order to earn both points. In the presence of no answer or an incorrect answer, such a table must be correctly labeled in order to earn the first point.
- Both points are earned for  $5\pi + 6\pi$ .
- The response may report the final answer as  $(2\pi, 11\pi)$ .

**Total for part (c) 2 points**

- (d) Find  $\int (t+5)\cos\left(\frac{t}{4}\right) dt$ . Show the work that leads to your answer.

|                                                                                                                          |                  |                |
|--------------------------------------------------------------------------------------------------------------------------|------------------|----------------|
| $u = t + 5 \quad dv = \cos\left(\frac{t}{4}\right) dt$<br>$du = dt \quad v = 4\sin\left(\frac{t}{4}\right)$              | $u$ and $dv$     | <b>1 point</b> |
| $\int (t+5)\cos\left(\frac{t}{4}\right) dt = 4(t+5)\sin\left(\frac{t}{4}\right) - \int 4\sin\left(\frac{t}{4}\right) dt$ | $uv - \int v du$ | <b>1 point</b> |
| $= 4(t+5)\sin\left(\frac{t}{4}\right) + 16\cos\left(\frac{t}{4}\right) + C$                                              | Answer           | <b>1 point</b> |

**Scoring notes:**

- The first and second points are earned with an implied  $u$  and  $dv$  in the presence of  $4(t+5)\sin\left(\frac{t}{4}\right) - \int 4\sin\left(\frac{t}{4}\right) dt$  or a mathematically equivalent expression.
- The tabular method may be used to show integration by parts. In this case, the first point is earned by columns (labeled or unlabeled) that begin with  $t+5$  and  $\cos\left(\frac{t}{4}\right)$ . The second point is earned for  $4(t+5)\sin\left(\frac{t}{4}\right) - \int 4\sin\left(\frac{t}{4}\right) dt$  or a mathematically equivalent expression.
- The third point is earned only for an expression mathematically equivalent to  $4(t+5)\sin\left(\frac{t}{4}\right) + 16\cos\left(\frac{t}{4}\right) + C$  (such as  $4t\sin\left(\frac{t}{4}\right) + 20\sin\left(\frac{t}{4}\right) + 16\cos\left(\frac{t}{4}\right) + C$ ) in the presence of correct supporting work.
- To earn the third point a response must have a final answer that includes a constant of integration.
- Alternate solution:

$$\int (t+5)\cos\left(\frac{t}{4}\right) dt = \int t\cos\left(\frac{t}{4}\right) dt + \int 5\cos\left(\frac{t}{4}\right) dt$$

$$u = t \quad dv = \cos\left(\frac{t}{4}\right) dt$$

$$du = dt \quad v = 4\sin\left(\frac{t}{4}\right)$$

$$\int t\cos\left(\frac{t}{4}\right) dt + \int 5\cos\left(\frac{t}{4}\right) dt = 4t\sin\left(\frac{t}{4}\right) - \int 4\sin\left(\frac{t}{4}\right) dt + \int 5\cos\left(\frac{t}{4}\right) dt$$


$$= 4t\sin\left(\frac{t}{4}\right) + 16\cos\left(\frac{t}{4}\right) + 20\sin\left(\frac{t}{4}\right) + C$$

- A response can earn the first and second points for correctly applying integration by parts to  $\int t\cos\left(\frac{t}{4}\right) dt$ . The tabular method may be used to show integration by parts. The third point is earned for the correct answer.

**Total for part (d) 3 points****Total for question 5 9 points****Part A (BC): Graphing calculator required****Question 2****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.



For  $0 \leq t \leq \pi$ , a particle is moving along the curve shown so that its position at time  $t$  is  $(x(t), y(t))$ , where  $x(t)$  is not explicitly given and  $y(t) = 2\sin t$ . It is known that  $\frac{dx}{dt} = e^{\cos t}$ . At time  $t = 0$ , the particle is at position  $(1, 0)$ .

| Model Solution                                                                                               | Scoring                            |
|--------------------------------------------------------------------------------------------------------------|------------------------------------|
| (a) Find the acceleration vector of the particle at time $t = 1$ . Show the setup for your calculations.     |                                    |
| $x''(1) = \frac{d}{dt}(e^{\cos t})\Big _{t=1} = -1.444407$                                                   | $x''(1)$ with setup <b>1 point</b> |
| $y(t) = 2\sin t \Rightarrow y'(t) = 2\cos t$<br>$y''(1) = \frac{d}{dt}(2\cos t)\Big _{t=1} = -1.682942$      | $y''(1)$ with setup <b>1 point</b> |
| The acceleration vector at time $t = 1$ is<br>$a(1) = \langle -1.444, -1.683 \text{ (or } -1.682) \rangle$ . |                                    |

**Scoring notes:**

- The exact answer is  $\langle x''(1), y''(1) \rangle = \langle -e^{\cos 1} \sin 1, -2 \sin 1 \rangle$ .
- $\langle -e^{\cos t} \sin t, -2 \sin t \rangle$  together with an incorrect or missing evaluation at  $t = 1$  earns 1 of the 2 points.

- A response of  $\langle -e^{\cos t} \sin t, -2 \sin t \rangle = \langle -e^{\cos 1} \sin 1, -2 \sin 1 \rangle$  or equivalent earns only 1 of the 2 points because it equates an expression to a numerical value.
- An unsupported correct acceleration vector earns 1 of the 2 points.
- The acceleration vector may be presented with other symbols, for example  $(\ , \ )$  or  $[ \ , \ ]$ .
- The components may be listed separately, as long as they are labeled.
- Degree mode: A response that presents answers obtained by using a calculator in degree mode does not earn the first point it would have otherwise earned. The response is generally eligible for all subsequent points (unless no answer is possible in degree mode or the question is made simpler by using degree mode). In degree mode,  $x''(1) = -0.000828$  or  $-0.047433$  and  $y''(1) = -0.000609$  or  $-0.034905$ . A response that presents one of these values with correct setups earns 1 of the 2 points.

**Total for part (a) 2 points**

- (b) For  $0 \leq t \leq \pi$ , find the first time  $t$  at which the speed of the particle is 1.5. Show the work that leads to your answer.

|                                                                                                                            |                                                                             |
|----------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|
| $\text{Speed} = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} = \sqrt{(e^{\cos t})^2 + (2 \cos t)^2}$ | $\sqrt{(e^{\cos t})^2 + (2 \cos t)^2} = 1.5 \quad \mathbf{1 \text{ point}}$ |
|----------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|

$$0 \leq t \leq \pi \text{ and } \sqrt{(e^{\cos t})^2 + (2 \cos t)^2} = 1.5$$

$$\Rightarrow t = 1.254472, t = 2.358077$$

The first time at which the speed of the particle is 1.5 is  $t = 1.254$ .

Answer **1 point**

**Scoring notes:**

- A response with an implied equation is eligible for both points. For example, a response of “Speed =  $\sqrt{(e^{\cos t})^2 + (2 \cos t)^2}$  and is first equal to 1.5 at  $t = 1.254$ ” earns both points.
- $\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} = 1.5$  earns the first point. Speed = 1.5 by itself does not earn the first point. Both of these responses are eligible to earn the second point.
- A response need not consider the value  $t = 2.358077$ .
- A response of  $t = 1.254$  alone does not earn either point.
- A response with a parenthesis error(s) in either  $(e^{\cos t})^2$  or  $(2 \cos t)^2$  does not earn the first point but does earn the second point for the correct answer. Note:  $\sqrt{\frac{dx^2}{dt} + \frac{dy^2}{dt}}$  is not considered a parenthesis error.

- Degree mode: In degree mode,  $\sqrt{(e^{\cos t})^2 + (2 \cos t)^2} = 1.5$  has no solution for  $0 \leq t \leq \pi$ .

A response that finds no time  $t$  at which the speed of the particle is 1.5 cannot be assumed to be working in degree mode.

**Total for part (b) 2 points**

- (c) Find the slope of the line tangent to the path of the particle at time  $t = 1$ . Find the  $x$ -coordinate of the position of the particle at time  $t = 1$ . Show the work that leads to your answers.

|                                                                                                                                          |                                                                              |
|------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------|
| $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2 \cos t}{e^{\cos t}}$ $\frac{dy}{dx} \Big _{t=1} = \frac{2 \cos 1}{e^{\cos 1}} = 0.629530$ | Slope with supporting work <span style="float: right;"><b>1 point</b></span> |
|------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------|

The slope of the line tangent to the curve at  $t = 1$  is 0.630 (or 0.629).

|                                                                                   |                                                         |
|-----------------------------------------------------------------------------------|---------------------------------------------------------|
| $x(1) = x(0) + \int_0^1 \frac{dx}{dt} dt = 1 + \int_0^1 e^{\cos t} dt = 3.341575$ | $\int_0^1 e^{\cos t} dt \quad \mathbf{1 \text{ point}}$ |
|-----------------------------------------------------------------------------------|---------------------------------------------------------|

|                                                                     |                                       |
|---------------------------------------------------------------------|---------------------------------------|
| The $x$ -coordinate of the position at $t = 1$ is 3.342 (or 3.341). | $x(1) \quad \mathbf{1 \text{ point}}$ |
|---------------------------------------------------------------------|---------------------------------------|

**Scoring notes:**

- To earn the first point, the response must communicate  $\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$ ; for example:
  - $\frac{dy}{dx} = \frac{2 \cos 1}{e^{\cos 1}}$
  - $\frac{dy/dt}{dx/dt} = 0.63$
  - $x'(1) = 1.716526, y'(1) = 1.080605, \text{ slope} = 0.63$
  - $\frac{dy}{dt} = 2 \cos t, \text{ slope} = 0.63$
- A response may import an incorrect expression for  $y'(t)$  or value of  $y'(1)$  from part (a), provided it was declared in part (a).
- The second point is earned for a response that presents the definite integral  $\int_0^1 e^{\cos t} dt$  or  $\int_0^1 \frac{dx}{dt} dt$  with or without the initial condition.

- For the second point, if the differential is missing:
  - $\int_0^1 e^{\cos t}$  earns the second point and is eligible for the third point.
  - $x(1) = \int_0^1 e^{\cos t}$  earns the second point but is not eligible for the third point.
  - $x(1) = 1 + \int_0^1 e^{\cos t}$  earns the second point and is eligible for the third point.
  - $x(1) = \int_0^1 e^{\cos t} + 1$  does not earn the second point but earns the third point for the correct answer.
- The third point is not earned for a response that presents an incorrect statement, such as  $x(1) = \int_0^1 e^{\cos t} dt = 1 + 2.342$ .
- Degree mode: In degree mode,  $\frac{dy}{dx} = 0.735759$  or  $0.012841$  and  $1 + \int_0^1 e^{\cos t} dt = 3.718144$ .

**Total for part (c) 3 points**

- (d) Find the total distance traveled by the particle over the time interval  $0 \leq t \leq \pi$ . Show the setup for your calculations.

|                                                      |          |         |
|------------------------------------------------------|----------|---------|
| $\int_0^\pi \sqrt{(e^{\cos t})^2 + (2 \cos t)^2} dt$ | Integral | 1 point |
| $= 6.034611$                                         | Answer   | 1 point |

The total distance traveled by the particle over  $0 \leq t \leq \pi$  is 6.035 (or 6.034).

**Scoring notes:**

- The first point is earned for presenting the correct integrand in a definite integral.
- Parentheses errors were assessed in part (b) and, therefore, will not affect the scoring in part (d).
- If the integrand is an incorrect speed function imported from part (b), the response earns the first point and does not earn the second point.
- An unsupported answer of 6.035 (or 6.034) does not earn either point.
- Degree mode: In degree mode, the total distance is 10.596835 or 8.536161.

**Total for part (d) 2 points**

**Total for question 2 9 points**

**Part A (AB or BC): Graphing calculator required**

**Question 1**

**9 points**

**General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

From 5 A.M. to 10 A.M., the rate at which vehicles arrive at a certain toll plaza is given by  $A(t) = 450\sqrt{\sin(0.62t)}$ , where  $t$  is the number of hours after 5 A.M. and  $A(t)$  is measured in vehicles per hour. Traffic is flowing smoothly at 5 A.M. with no vehicles waiting in line.

| Model Solution                                                                                                                                                               | Scoring        |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| (a) Write, but do not evaluate, an integral expression that gives the total number of vehicles that arrive at the toll plaza from 6 A.M. ( $t = 1$ ) to 10 A.M. ( $t = 5$ ). |                |
| The total number of vehicles that arrive at the toll plaza from 6 A.M. to 10 A.M. is given by $\int_1^5 A(t) dt$ .                                                           | Answer 1 point |

**Scoring notes:**

- The response must be a definite integral with correct lower and upper limits to earn this point.
- Because  $|A(t)| = A(t)$  for  $1 \leq t \leq 5$ , a response of  $\int_1^5 |450\sqrt{\sin(0.62t)}| dt$  or  $\int_1^5 |A(t)| dt$  earns the point.
- A response missing  $dt$  or using  $dx$  is eligible to earn the point.
- A response with a copy error in the expression for  $A(t)$  will earn the point only in the presence of  $\int_1^5 A(t) dt$ .

**Total for part (a) 1 point**

- (b) Find the average value of the rate, in vehicles per hour, at which vehicles arrive at the toll plaza from 6 A.M. ( $t = 1$ ) to 10 A.M. ( $t = 5$ ).

$$\text{Average} = \frac{1}{5-1} \int_1^5 A(t) dt = 375.536966$$

Uses average value formula: **1 point**

$$\frac{1}{b-a} \int_a^b A(t) dt$$

The average rate at which vehicles arrive at the toll plaza from 6 A.M. to 10 A.M. is 375.537 (or 375.536) vehicles per hour.

Answer **1 point**

**Scoring notes:**

- The use of the average value formula, indicating that  $a = 1$  and  $b = 5$ , can be presented in single or multiple steps to earn the first point. For example, the following response earns both points:

$$\int_1^5 A(t) dt = 1502.147865, \text{ so the average value is } 375.536966.$$

- A response that presents a correct integral along with the correct average value, but provides incorrect or incomplete communication, earns 1 out of 2 points. For example, the following response earns 1 out of 2 points:  $\int_1^5 A(t) dt = 1502.147865 = 375.536966$ .

- The answer must be correct to three decimal places. For example,

$$\frac{1}{5-1} \int_1^5 A(t) dt = 375.536966 \approx 376 \text{ earns only the first point.}$$

- Degree mode: A response that presents answers obtained by using a calculator in degree mode does not earn the first point it would have otherwise earned. The response is generally eligible for all subsequent points (unless no answer is possible in degree mode or the question is made simpler by using degree mode). In degree mode,  $\frac{1}{4} \int_1^5 A(t) dt = 79.416068$ .

- Special case:  $\frac{1}{5} \int_1^5 A(t) dt = 300.429573$  earns 1 out of 2 points.

**Total for part (b) 2 points**

- (c) Is the rate at which vehicles arrive at the toll plaza at 6 A.M. ( $t = 1$ ) increasing or decreasing? Give a reason for your answer.

$$A'(1) = 148.947272$$

Considers  $A'(1)$  **1 point**

Because  $A'(1) > 0$ , the rate at which the vehicles arrive at the toll plaza is increasing.

Answer with reason **1 point**

**Scoring notes:**

- The response need not present the value of  $A'(1)$ . The second line of the model solution earns both points.
- An incorrect value assigned to  $A'(1)$  earns the first point (but will not earn the second point).
- Without a reference to  $t = 1$ , the first point is earned by any of the following:
  - 148.947 accurate to the number of decimals presented, with zero up to three decimal places (i.e., 149, 148, 148.9, 148.95, or 148.94)
  - $A'(t) = 148.947$  by itself
- To be eligible for the second point, the first point must be earned.
- To earn the second point, there must be a reference to  $t = 1$ .
- Degree mode:  $A'(1) = 23.404311$

**Total for part (c) 2 points**

- (d) A line forms whenever  $A(t) \geq 400$ . The number of vehicles in line at time  $t$ , for  $a \leq t \leq 4$ , is given by

$N(t) = \int_a^t (A(x) - 400) dx$ , where  $a$  is the time when a line first begins to form. To the nearest whole number, find the greatest number of vehicles in line at the toll plaza in the time interval  $a \leq t \leq 4$ . Justify your answer.

$$N'(t) = A(t) - 400 = 0$$

Considers  $N'(t) = 0$  **1 point**

$$\Rightarrow A(t) = 400 \Rightarrow t = 1.469372, t = 3.597713$$

$t = a$  and  $t = b$  **1 point**

$$a = 1.469372$$

$$b = 3.597713$$

| $t$ | $N(t) = \int_a^t (A(x) - 400) dx$ |
|-----|-----------------------------------|
| $a$ | 0                                 |
| $b$ | 71.254129                         |
| 4   | 62.338346                         |

Answer **1 point**

Justification **1 point**

The greatest number of vehicles in line is 71.

**Scoring notes:**

- It is not necessary to indicate that  $A(t) = 400$  to earn the first point, although this statement alone would earn the first point.
- A response of “ $A(t) \geq 400$  when  $1.469372 \leq t \leq 3.597713$ ” will earn the first 2 points. A response of “ $A(t) \geq 400$ ” along with the presence of exactly one of the two numbers above will earn the first point, but not the second. A response of “ $A(t) \geq 400$ ” by itself will not earn either of the first 2 points.
- To earn the second point the values for  $a$  and  $b$  must be accurate to the number of decimals presented, with at least one and up to three decimal places. These may appear only in a candidates table, as limits of integration, or on a number line.
- A response with incorrect notation involving  $t$  or  $x$  is eligible to earn all 4 points.
- A response that does not earn the first point is still eligible for the remaining 3 points.
- To earn the third point, a response must present the greatest number of vehicles. This point is earned for answers of either 71 or 71.254\*\*\* only.
- A correct justification earns the fourth point, even if the third point is not earned because of a decimal presentation error.
- When using a Candidates Test, the response must include the values for  $N(a)$ ,  $N(b)$ , and  $N(4)$  to earn the fourth point. These values must be correct to the number of decimals presented, with up to three decimal places. (Correctly rounded integer values are acceptable.)
- Alternate solution for the third and fourth points:

For  $a \leq t \leq b$ ,  $A(t) \geq 400$ . For  $b \leq t \leq 4$ ,  $A(t) \leq 400$ .

Thus,  $N(t) = \int_a^t (A(x) - 400) dx$  is greatest at  $t = b$ .

$N(b) = 71.254129$ , and the greatest number of vehicles in line is 71.

- Degree mode: The response is only eligible to earn the first point because in degree mode  $A(t) < 400$ .


**Total for part (d) 4 points**

**Total for question 1 9 points**

**Part B (AB or BC): Graphing calculator not allowed****Question 3****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.



Let  $f$  be a differentiable function with  $f(4) = 3$ . On the interval  $0 \leq x \leq 7$ , the graph of  $f'$ , the derivative of  $f$ , consists of a semicircle and two line segments, as shown in the figure above.

| Model Solution                                                       | Scoring                                                            |
|----------------------------------------------------------------------|--------------------------------------------------------------------|
| (a) Find $f(0)$ and $f(5)$ .                                         |                                                                    |
| $f(0) = f(4) + \int_4^0 f'(x) dx = 3 - \int_0^4 f'(x) dx = 3 + 2\pi$ | Area of either region <b>1 point</b><br>– OR – $\int_0^4 f'(x) dx$ |
| $f(5) = f(4) + \int_4^5 f'(x) dx = 3 + \frac{1}{2} = \frac{7}{2}$    | – OR – $\int_4^5 f'(x) dx$                                         |
|                                                                      | $f(0)$ <b>1 point</b>                                              |
|                                                                      | $f(5)$ <b>1 point</b>                                              |

**Scoring notes:**

- A response with answers of only  $f(0) = \pm 2\pi$ , or only  $f(5) = \frac{1}{2}$ , or both earns 1 of the 3 points.
- A response displaying  $f(5) = \frac{7}{2}$  and a missing or incorrect value for  $f(0)$  earns 2 of the 3 points.
- The second and third points can be earned in either order.
- Read unlabeled values from left to right and from top to bottom as  $f(0)$  and  $f(5)$ . A single value must be labeled as either  $f(0)$  or  $f(5)$  in order to earn any points.

**Total for part (a) 3 points**

- (b) Find the  $x$ -coordinates of all points of inflection of the graph of  $f$  for  $0 < x < 7$ . Justify your answer.

|                                                                                                                                                                                             |               |         |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|---------|
| The graph of $f$ has a point of inflection at each of $x = 2$ and $x = 6$ , because $f'(x)$ changes from decreasing to increasing at $x = 2$ and from increasing to decreasing at $x = 6$ . | Answer        | 1 point |
|                                                                                                                                                                                             | Justification | 1 point |

**Scoring notes:**

- A response that gives only one of  $x = 2$  or  $x = 6$ , along with a correct justification, earns 1 of the 2 points.
- A response that claims that there is a point of inflection at any value other than  $x = 2$  or  $x = 6$  earns neither point.
- To earn the second point a response must use correct reasoning based on the graph of  $f'$ . Examples of correct reasoning include:
  - Correctly discussing the signs of the slopes of the graph of  $f'$
  - Citing  $x = 2$  and  $x = 6$  as the locations of local extrema on the graph of  $f'$
- Examples of reasoning not (sufficiently) connected to the graph of  $f'$  include:
  - Reasoning based on sign changes in  $f''$  unless the connection is made between the sign of  $f''$  and the slopes of the graph of  $f'$
  - Reasoning based only on the concavity of the graph of  $f$
- The second point cannot be earned by use of vague or undefined terms such as “it” or “the function” or “the derivative.”
- Responses that report inflection points as ordered pairs must report the points  $(2, 3 + \pi)$  and  $(6, 5)$  in order to earn the first point. If the  $y$ -coordinates are reported incorrectly, the response remains eligible for the second point.

**Total for part (b) 2 points**

- (c) Let  $g$  be the function defined by  $g(x) = f(x) - x$ . On what intervals, if any, is  $g$  decreasing for  $0 \leq x \leq 7$ ? Show the analysis that leads to your answer.

|                                                                                                           |                      |         |
|-----------------------------------------------------------------------------------------------------------|----------------------|---------|
| $g'(x) = f'(x) - 1$                                                                                       | $g'(x) = f'(x) - 1$  | 1 point |
| $f'(x) - 1 \leq 0 \Rightarrow f'(x) \leq 1$                                                               | Interval with reason | 1 point |
| The graph of $g$ is decreasing on the interval $0 \leq x \leq 5$ because $g'(x) \leq 0$ on this interval. |                      |         |

**Scoring notes:**

- The first point can be earned for  $f'(x) \leq 1$  or the equivalent, in words or symbols.
- Endpoints do not need to be included in the interval to be eligible for the second point.

**Total for part (c) 2 points**

- (d) For the function  $g$  defined in part (c), find the absolute minimum value on the interval  $0 \leq x \leq 7$ . Justify your answer.

|                                                                                    |                           |         |
|------------------------------------------------------------------------------------|---------------------------|---------|
| $g$ is continuous, $g'(x) < 0$ for $0 < x < 5$ , and $g'(x) > 0$ for $5 < x < 7$ . | Considers $g'(x) = 0$     | 1 point |
|                                                                                    | Answer with justification | 1 point |

Therefore, the absolute minimum occurs at  $x = 5$ , and

$g(5) = f(5) - 5 = \frac{7}{2} - 5 = -\frac{3}{2}$  is the absolute minimum value of  $g$ .

**Scoring notes:**

- A justification that uses a local argument, such as “ $g'$  changes from negative to positive (or  $g$  changes from decreasing to increasing) at  $x = 5$ ” must also state that  $x = 5$  is the only critical point.
- If  $g'(x) = 0$  (or equivalent) is not declared explicitly, a response that isolates  $x = 5$  as the only critical number belonging to  $(0, 7)$  earns the first point.
- A response that imports  $g'(x) = f'(x)$  from part (c) is eligible for the first point but not the second.
  - In this case, consideration of  $x = 4$  as the only critical number on  $(0, 7)$  earns the first point.
- Solution using Candidates Test:

$$g'(x) = f'(x) - 1 = 0 \Rightarrow x = 5, x = 7$$

| $x$ | $g(x)$         |
|-----|----------------|
| 0   | $3 + 2\pi$     |
| 5   | $-\frac{3}{2}$ |
| 7   | $-\frac{1}{2}$ |

The absolute minimum value of  $g$  on the interval  $0 \leq x \leq 7$  is  $-\frac{3}{2}$ .

- When using a Candidates Test, a response may import an incorrect value of  $f(0) = g(0) > -\frac{3}{2}$  from part (a). The second point can only be earned for an answer of  $-\frac{3}{2}$ .


**Total for part (d) 2 points**

**Total for question 3 9 points**

**Part B (AB or BC): Graphing calculator not allowed****Question 4****9 points****General Scoring Notes**

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Scoring guidelines and notes contain examples of the most common approaches seen in student responses. These guidelines can be applied to alternate approaches to ensure that these alternate approaches are scored appropriately.

Graph of  $f$ 

Let  $f$  be a continuous function defined on the closed interval  $-4 \leq x \leq 6$ . The graph of  $f$ , consisting of four line segments, is shown above. Let  $G$  be the function defined by  $G(x) = \int_0^x f(t) dt$ .

| Model Solution                                                                                                                                                                                                                                                                                                                                                                    | Scoring                       |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------|
| $G'(x) = f(x)$ in any part of the response.                                                                                                                                                                                                                                                                                                                                       | $G'(x) = f(x)$ <b>1 point</b> |
| <b>Scoring notes:</b> <ul style="list-style-type: none"> <li>This “global point” can be earned in any one part. Expressions that show this connection and therefore earn this point include: <math>G' = f</math>, <math>G'(x) = f(x)</math>, <math>G''(x) = f'(x)</math> in part (a), <math>G'(3) = f(3)</math> in part (b), or <math>G'(2) = f(2)</math> in part (c).</li> </ul> |                               |
| <b>Total</b>                                                                                                                                                                                                                                                                                                                                                                      | <b>1 point</b>                |

- (a) On what open intervals is the graph of  $G$  concave up? Give a reason for your answer.

$$G'(x) = f(x)$$

Answer with reason

**1 point**

The graph of  $G$  is concave up for  $-4 < x < -2$  and  $2 < x < 6$ , because  $G' = f$  is increasing on these intervals.

**Scoring notes:**

- Intervals may also include one or both endpoints.

**Total for part (a) 1 point**

- (b) Let  $P$  be the function defined by  $P(x) = G(x) \cdot f(x)$ . Find  $P'(3)$ .

$$P'(x) = G(x) \cdot f'(x) + f(x) \cdot G'(x)$$

Product rule

**1 point**

$$P'(3) = G(3) \cdot f'(3) + f(3) \cdot G'(3)$$

$$\text{Substituting } G(3) = \int_0^3 f(t) dt = -3.5 \text{ and } G'(3) = f(3) = -3$$

 $G(3)$  or  $G'(3)$ **1 point**

into the above expression for  $P'(3)$  gives the following:

$$P'(3) = -3.5 \cdot 1 + (-3) \cdot (-3) = 5.5$$

Answer

**1 point****Scoring notes:**

- The first point is earned for the correct application of the product rule in terms of  $x$  or in the evaluation of  $P'(3)$ . Once earned, this point cannot be lost.
- The second point is earned by correctly evaluating  $G(3) = -3.5$ ,  $G'(3) = -3$ , or  $f(3) = -3$ .
- To be eligible to earn the third point a response must have earned the first two points.
- Simplification of the numerical value is not required to earn the third point.

**Total for part (b) 3 points**

(c) Find  $\lim_{x \rightarrow 2} \frac{G(x)}{x^2 - 2x}$ .

$$\lim_{x \rightarrow 2} (x^2 - 2x) = 0$$

Because  $G$  is continuous for  $-4 \leq x \leq 6$ ,

$$\lim_{x \rightarrow 2} G(x) = \int_0^2 f(t) dt = 0.$$

Therefore, the limit  $\lim_{x \rightarrow 2} \frac{G(x)}{x^2 - 2x}$  is an indeterminate form of type  $\frac{0}{0}$ .

Using L'Hospital's Rule,

$$\begin{aligned} \lim_{x \rightarrow 2} \frac{G(x)}{x^2 - 2x} &= \lim_{x \rightarrow 2} \frac{G'(x)}{2x - 2} \\ &= \lim_{x \rightarrow 2} \frac{f(x)}{2x - 2} = \frac{f(2)}{2} = \frac{-4}{2} = -2 \end{aligned}$$

**Scoring notes:**

- To earn the first point the response must show  $\lim_{x \rightarrow 2} (x^2 - 2x) = 0$  and  $\lim_{x \rightarrow 2} G(x) = 0$  and must show a ratio of the two derivatives,  $G'(x)$  and  $2x - 2$ . The ratio may be shown as evaluations of the derivatives at  $x = 2$ , such as  $\frac{G'(2)}{2}$ .
- To earn the second point the response must evaluate correctly with appropriate limit notation. In particular the response must include  $\lim_{x \rightarrow 2} \frac{G'(x)}{2x - 2}$  or  $\lim_{x \rightarrow 2} \frac{f(x)}{2x - 2}$ .
- With any linkage errors (such as  $\frac{G'(x)}{2x - 2} = \frac{f(2)}{2}$ ), the response does not earn the second point.

**Total for part (c) 2 points**

(d) Find the average rate of change of  $G$  on the interval  $[-4, 2]$ . Does the Mean Value Theorem guarantee a value  $c$ ,  $-4 < c < 2$ , for which  $G'(c)$  is equal to this average rate of change? Justify your answer.

$$G(2) = \int_0^2 f(t) dt = 0 \text{ and } G(-4) = \int_0^{-4} f(t) dt = -16$$

Average rate of change

**1 point**

$$\text{Average rate of change} = \frac{G(2) - G(-4)}{2 - (-4)} = \frac{0 - (-16)}{6} = \frac{8}{3}$$

Yes,  $G'(x) = f(x)$  so  $G$  is differentiable on  $(-4, 2)$  and continuous on  $[-4, 2]$ . Therefore, the Mean Value Theorem applies and guarantees a value  $c$ ,  $-4 < c < 2$ , such that

Answer with justification

**1 point**

$$G'(c) = \frac{8}{3}.$$

**Scoring notes:**

- To earn the first point a response must present at least a difference and a quotient and a correct evaluation. For example,  $\frac{0 + 16}{6}$  or  $\frac{G(2) - G(-4)}{6} = \frac{16}{6}$ .
- Simplification of the numerical value is not required to earn the first point, but any simplification must be correct.
- The second point can be earned without the first point if the response has the correct setup but an incorrect or no evaluation of the average rate of change. The Mean Value Theorem need not be explicitly stated provided the conditions and conclusion are stated.

**Total for part (d) 2 points**

**Total for question 4 9 points**

2019

AP<sup>®</sup> CollegeBoard

# AP<sup>®</sup> Calculus BC

## Scoring Guidelines

## 2019 SCORING GUIDELINES

## Question 1

(a)  $\int_0^5 E(t) dt = 153.457690$

To the nearest whole number, 153 fish enter the lake from midnight to 5 A.M.

(b)  $\frac{1}{5-0} \int_0^5 L(t) dt = 6.059038$

The average number of fish that leave the lake per hour from midnight to 5 A.M. is 6.059 fish per hour.

- (c) The rate of change in the number of fish in the lake at time  $t$  is given by  $E(t) - L(t)$ .

$$E(t) - L(t) = 0 \Rightarrow t = 6.20356$$

$E(t) - L(t) > 0$  for  $0 \leq t < 6.20356$ , and  $E(t) - L(t) < 0$  for  $6.20356 < t \leq 8$ . Therefore the greatest number of fish in the lake is at time  $t = 6.204$  (or 6.203).

— OR —

Let  $A(t)$  be the change in the number of fish in the lake from midnight to  $t$  hours after midnight.

$$A(t) = \int_0^t (E(s) - L(s)) ds$$

$$A'(t) = E(t) - L(t) = 0 \Rightarrow t = C = 6.20356$$

| $t$ | $A(t)$    |
|-----|-----------|
| 0   | 0         |
| $C$ | 135.01492 |
| 8   | 80.91998  |

Therefore the greatest number of fish in the lake is at time  $t = 6.204$  (or 6.203).

(d)  $E'(5) - L'(5) = -10.7228 < 0$

Because  $E'(5) - L'(5) < 0$ , the rate of change in the number of fish is decreasing at time  $t = 5$ .

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

3 :  $\begin{cases} 1 : \text{sets } E(t) - L(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

2 :  $\begin{cases} 1 : \text{considers } E'(5) \text{ and } L'(5) \\ 1 : \text{answer with explanation} \end{cases}$

## 2019 SCORING GUIDELINES

## Question 2

(a)  $\frac{1}{2} \int_0^{\sqrt{\pi}} (r(\theta))^2 d\theta = 3.534292$

The area of  $S$  is 3.534.

(b)  $\frac{1}{\sqrt{\pi}-0} \int_0^{\sqrt{\pi}} r(\theta) d\theta = 1.579933$

The average distance from the origin to a point on the curve  $r = r(\theta)$  for  $0 \leq \theta \leq \sqrt{\pi}$  is 1.580 (or 1.579).

(c)  $\tan \theta = \frac{y}{x} = m \Rightarrow \theta = \tan^{-1} m$

$$\frac{1}{2} \int_0^{\tan^{-1} m} (r(\theta))^2 d\theta = \frac{1}{2} \left( \frac{1}{2} \int_0^{\sqrt{\pi}} (r(\theta))^2 d\theta \right)$$

- (d) As  $k \rightarrow \infty$ , the circle  $r = k \cos \theta$  grows to enclose all points to the right of the  $y$ -axis.

$$\begin{aligned} \lim_{k \rightarrow \infty} A(k) &= \frac{1}{2} \int_0^{\pi/2} (r(\theta))^2 d\theta \\ &= \frac{1}{2} \int_0^{\pi/2} (3\sqrt{\theta} \sin(\theta^2))^2 d\theta = 3.324 \end{aligned}$$

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

3 :  $\begin{cases} 1 : \text{equates polar areas} \\ 1 : \text{inverse trigonometric function} \\ \text{applied to } m \text{ as limit of} \\ \text{integration} \\ 1 : \text{equation} \end{cases}$

2 :  $\begin{cases} 1 : \text{limits of integration} \\ 1 : \text{answer with integral} \end{cases}$

## 2019 SCORING GUIDELINES

## Question 3

$$\begin{aligned} \text{(a)} \quad \int_{-6}^5 f(x) \, dx &= \int_{-6}^{-2} f(x) \, dx + \int_{-2}^5 f(x) \, dx \\ &\Rightarrow 7 = \int_{-6}^{-2} f(x) \, dx + 2 + \left(9 - \frac{9\pi}{4}\right) \\ &\Rightarrow \int_{-6}^{-2} f(x) \, dx = 7 - \left(11 - \frac{9\pi}{4}\right) = \frac{9\pi}{4} - 4 \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \int_3^5 (2f'(x) + 4) \, dx &= 2 \int_3^5 f'(x) \, dx + \int_3^5 4 \, dx \\ &= 2(f(5) - f(3)) + 4(5 - 3) \\ &= 2(0 - (3 - \sqrt{5})) + 8 \\ &= 2(-3 + \sqrt{5}) + 8 = 2 + 2\sqrt{5} \end{aligned}$$

— OR —

$$\begin{aligned} \int_3^5 (2f'(x) + 4) \, dx &= [2f(x) + 4x]_{x=3}^{x=5} \\ &= (2f(5) + 20) - (2f(3) + 12) \\ &= (2 \cdot 0 + 20) - (2(3 - \sqrt{5}) + 12) \\ &= 2 + 2\sqrt{5} \end{aligned}$$

$$\text{(c)} \quad g'(x) = f(x) = 0 \Rightarrow x = -1, x = \frac{1}{2}, x = 5$$

| $x$           | $g(x)$                |
|---------------|-----------------------|
| -2            | 0                     |
| -1            | $\frac{1}{2}$         |
| $\frac{1}{2}$ | $-\frac{1}{4}$        |
| 5             | $11 - \frac{9\pi}{4}$ |

On the interval  $-2 \leq x \leq 5$ , the absolute maximum value of  $g$  is  $g(5) = 11 - \frac{9\pi}{4}$ .

$$\begin{aligned} \text{(d)} \quad \lim_{x \rightarrow 1} \frac{10^x - 3f'(x)}{f(x) - \arctan x} &= \frac{10^1 - 3f'(1)}{f(1) - \arctan 1} \\ &= \frac{10 - 3 \cdot 2}{1 - \arctan 1} = \frac{4}{1 - \frac{\pi}{4}} \end{aligned}$$

$$3 : \begin{cases} 1 : \int_{-6}^5 f(x) \, dx = \int_{-6}^{-2} f(x) \, dx + \int_{-2}^5 f(x) \, dx \\ 1 : \int_{-2}^5 f(x) \, dx \\ 1 : \text{answer} \end{cases}$$

$$2 : \begin{cases} 1 : \text{Fundamental Theorem of Calculus} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : g'(x) = f(x) \\ 1 : \text{identifies } x = -1 \text{ as a candidate} \\ 1 : \text{answer with justification} \end{cases}$$

1 : answer

## 2019 SCORING GUIDELINES

## Question 4

$$\begin{aligned} \text{(a)} \quad V &= \pi r^2 h = \pi(1)^2 h = \pi h \\ \frac{dV}{dt} \Big|_{h=4} &= \pi \frac{dh}{dt} \Big|_{h=4} = \pi \left( -\frac{1}{10} \sqrt{4} \right) = -\frac{\pi}{5} \text{ cubic feet per second} \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \frac{d^2 h}{dt^2} &= -\frac{1}{20\sqrt{h}} \cdot \frac{dh}{dt} = -\frac{1}{20\sqrt{h}} \cdot \left( -\frac{1}{10} \sqrt{h} \right) = \frac{1}{200} \\ \text{Because } \frac{d^2 h}{dt^2} &= \frac{1}{200} > 0 \text{ for } h > 0, \text{ the rate of change of the} \\ &\text{height is increasing when the height of the water is 3 feet.} \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad \frac{dh}{\sqrt{h}} &= -\frac{1}{10} dt \\ \int \frac{dh}{\sqrt{h}} &= \int -\frac{1}{10} dt \\ 2\sqrt{h} &= -\frac{1}{10} t + C \\ 2\sqrt{5} &= -\frac{1}{10} \cdot 0 + C \Rightarrow C = 2\sqrt{5} \\ 2\sqrt{h} &= -\frac{1}{10} t + 2\sqrt{5} \\ h(t) &= \left( -\frac{1}{20} t + \sqrt{5} \right)^2 \end{aligned}$$

$$2 : \begin{cases} 1 : \frac{dV}{dt} = \pi \frac{dh}{dt} \\ 1 : \text{answer with units} \end{cases}$$

$$3 : \begin{cases} 1 : \frac{d}{dh} \left( -\frac{1}{10} \sqrt{h} \right) = -\frac{1}{20\sqrt{h}} \\ 1 : \frac{d^2 h}{dt^2} = -\frac{1}{20\sqrt{h}} \cdot \frac{dh}{dt} \\ 1 : \text{answer with explanation} \end{cases}$$

$$4 : \begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ \text{and uses initial condition} \\ 1 : h(t) \end{cases}$$

Note: 0/4 if no separation of variables

Note: max 2/4 [1-1-0-0] if no constant of integration

## 2019 SCORING GUIDELINES

## Question 5

$$(a) f'(x) = \frac{-(2x-2)}{(x^2-2x+k)^2}$$

$$f'(0) = \frac{2}{k^2} = 6 \Rightarrow k^2 = \frac{1}{3} \Rightarrow k = \frac{1}{\sqrt{3}}$$

$$3 : \begin{cases} 1 : \text{denominator of } f'(x) \\ 1 : f'(x) \\ 1 : \text{answer} \end{cases}$$

$$(b) \frac{1}{x^2-2x-8} = \frac{1}{(x-4)(x+2)} = \frac{A}{x-4} + \frac{B}{x+2}$$

$$\Rightarrow 1 = A(x+2) + B(x-4)$$

$$\Rightarrow A = \frac{1}{6}, B = -\frac{1}{6}$$

$$3 : \begin{cases} 1 : \text{partial fraction decomposition} \\ 1 : \text{antiderivatives} \\ 1 : \text{answer} \end{cases}$$

$$\begin{aligned} \int_0^1 f(x) dx &= \int_0^1 \left( \frac{1}{6} \frac{1}{x-4} - \frac{1}{6} \frac{1}{x+2} \right) dx \\ &= \left[ \frac{1}{6} \ln|x-4| - \frac{1}{6} \ln|x+2| \right]_{x=0}^{x=1} \\ &= \left( \frac{1}{6} \ln 3 - \frac{1}{6} \ln 3 \right) - \left( \frac{1}{6} \ln 4 - \frac{1}{6} \ln 2 \right) = -\frac{1}{6} \ln 2 \end{aligned}$$

$$\begin{aligned} (c) \int_0^2 \frac{1}{x^2-2x+1} dx &= \int_0^2 \frac{1}{(x-1)^2} dx = \int_0^1 \frac{1}{(x-1)^2} dx + \int_1^2 \frac{1}{(x-1)^2} dx \\ &= \lim_{b \rightarrow 1^-} \int_0^b \frac{1}{(x-1)^2} dx + \lim_{b \rightarrow 1^+} \int_b^2 \frac{1}{(x-1)^2} dx \\ &= \lim_{b \rightarrow 1^-} \left( -\frac{1}{x-1} \Big|_{x=0}^{x=b} \right) + \lim_{b \rightarrow 1^+} \left( -\frac{1}{x-1} \Big|_{x=b}^{x=2} \right) \\ &= \lim_{b \rightarrow 1^-} \left( -\frac{1}{b-1} - 1 \right) + \lim_{b \rightarrow 1^+} \left( -1 + \frac{1}{b-1} \right) \end{aligned}$$

Because  $\lim_{b \rightarrow 1^-} \left( -\frac{1}{b-1} \right)$  does not exist, the integral diverges.

$$3 : \begin{cases} 1 : \text{improper integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer with reason} \end{cases}$$

## 2019 SCORING GUIDELINES

## Question 6

$$(a) f(0) = 3 \text{ and } f'(0) = -2$$

The third-degree Taylor polynomial for  $f$  about  $x = 0$  is

$$3 - 2x + \frac{3}{2!}x^2 + \frac{-2}{3!}x^3 = 3 - 2x + \frac{3}{2}x^2 - \frac{23}{12}x^3.$$

$$2 : \begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \end{cases}$$

$$(b) \text{ The first three nonzero terms of the Maclaurin series for } e^x \text{ are}$$

$$1 + x + \frac{1}{2!}x^2.$$

$$2 : \begin{cases} 1 : \text{three terms for } e^x \\ 1 : \text{three terms for } e^x f(x) \end{cases}$$

The second-degree Taylor polynomial for  $e^x f(x)$  about  $x = 0$  is

$$\begin{aligned} 3 \left( 1 + x + \frac{1}{2!}x^2 \right) - 2x(1+x) + \frac{3}{2}x^2(1) \\ = 3 + (3-2)x + \left( \frac{3}{2} - 2 + \frac{3}{2} \right)x^2 \\ = 3 + x + x^2. \end{aligned}$$

$$(c) h(1) = \int_0^1 f(t) dt$$

$$\begin{aligned} &\approx \int_0^1 \left( 3 - 2t + \frac{3}{2}t^2 - \frac{23}{12}t^3 \right) dt \\ &= \left[ 3t - t^2 + \frac{1}{2}t^3 - \frac{23}{48}t^4 \right]_{t=0}^{t=1} \\ &= 3 - 1 + \frac{1}{2} - \frac{23}{48} = \frac{97}{48} \end{aligned}$$

$$2 : \begin{cases} 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$$

$$(d) \text{ The alternating series error bound is the absolute value of the first omitted term of the series for } h(1).$$

$$\int_0^1 \left( \frac{54}{4!}t^4 \right) dt = \left[ \frac{9}{20}t^5 \right]_{t=0}^{t=1} = \frac{9}{20}$$

$$\text{Error} \leq \left| \frac{9}{20} \right| = 0.45$$

$$3 : \begin{cases} 1 : \text{uses fourth-degree term of Maclaurin series for } f \\ 1 : \text{uses first omitted term of series for } h(1) \\ 1 : \text{error bound} \end{cases}$$

# AP Calculus BC

## Scoring Guidelines

### 2018 SCORING GUIDELINES

#### Question 1

(a)  $\int_0^{300} r(t) \, dt = 270$

According to the model, 270 people enter the line for the escalator during the time interval  $0 \leq t \leq 300$ .

(b)  $20 + \int_0^{300} (r(t) - 0.7) \, dt = 20 + \int_0^{300} r(t) \, dt - 0.7 \cdot 300 = 80$

According to the model, 80 people are in line at time  $t = 300$ .

(c) Based on part (b), the number of people in line at time  $t = 300$  is 80.

The first time  $t$  that there are no people in line is

$$300 + \frac{80}{0.7} = 414.286 \text{ (or 414.285) seconds.}$$

(d) The total number of people in line at time  $t$ ,  $0 \leq t \leq 300$ , is modeled by  $20 + \int_0^t r(x) \, dx - 0.7t$ .

$$r(t) - 0.7 = 0 \Rightarrow t_1 = 33.013298, t_2 = 166.574719$$

| $t$   | People in line for escalator |
|-------|------------------------------|
| 0     | 20                           |
| $t_1$ | 3.803                        |
| $t_2$ | 158.070                      |
| 300   | 80                           |

The number of people in line is a minimum at time  $t = 33.013$  seconds, when there are 4 people in line.

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

2 :  $\begin{cases} 1 : \text{considers rate out} \\ 1 : \text{answer} \end{cases}$

1 : answer

4 :  $\begin{cases} 1 : \text{considers } r(t) - 0.7 = 0 \\ 1 : \text{identifies } t = 33.013 \\ 1 : \text{answers} \\ 1 : \text{justification} \end{cases}$

## 2018 SCORING GUIDELINES

## Question 2

(a)  $p'(25) = -1.179$

At a depth of 25 meters, the density of plankton cells is changing at a rate of  $-1.179$  million cells per cubic meter per meter.

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{meaning with units} \end{cases}$$

(b)  $\int_0^{30} 3p(h) \, dh = 1675.414936$

There are 1675 million plankton cells in the column of water between  $h = 0$  and  $h = 30$  meters.

$$2 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$$

(c)  $\int_{30}^K 3f(h) \, dh$  represents the number of plankton cells, in millions, in the column of water from a depth of 30 meters to a depth of  $K$  meters.

The number of plankton cells, in millions, in the entire column of water is given by  $\int_0^{30} 3p(h) \, dh + \int_{30}^K 3f(h) \, dh$ .

Because  $0 \leq f(h) \leq u(h)$  for all  $h \geq 30$ ,

$$3 \int_{30}^K f(h) \, dh \leq 3 \int_{30}^K u(h) \, dh \leq 3 \int_{30}^{\infty} u(h) \, dh = 3 \cdot 105 = 315.$$

The total number of plankton cells in the column of water is bounded by  $1675.415 + 315 = 1990.415 \leq 2000$  million.

$$3 : \begin{cases} 1 : \text{integral expression} \\ 1 : \text{compares improper integral} \\ 1 : \text{explanation} \end{cases}$$

(d)  $\int_0^1 \sqrt{(x'(t))^2 + (y'(t))^2} \, dt = 757.455862$

The total distance traveled by the boat over the time interval  $0 \leq t \leq 1$  is 757.456 (or 757.455) meters.

$$2 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{total distance} \end{cases}$$

## 2018 SCORING GUIDELINES

## Question 3

$$\begin{aligned} \text{(a)} \quad f(-5) &= f(1) + \int_1^{-5} g(x) \, dx = f(1) - \int_{-5}^1 g(x) \, dx \\ &= 3 - \left(-9 - \frac{3}{2} + 1\right) = 3 - \left(-\frac{19}{2}\right) = \frac{25}{2} \end{aligned}$$

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

$$\begin{aligned} \text{(b)} \quad \int_1^6 g(x) \, dx &= \int_1^3 g(x) \, dx + \int_3^6 g(x) \, dx \\ &= \int_1^3 2 \, dx + \int_3^6 2(x-4)^2 \, dx \\ &= 4 + \left[\frac{2}{3}(x-4)^3\right]_{x=3}^{x=6} = 4 + \frac{16}{3} - \left(-\frac{2}{3}\right) = 10 \end{aligned}$$

$$3 : \begin{cases} 1 : \text{split at } x = 3 \\ 1 : \text{antiderivative of } 2(x-4)^2 \\ 1 : \text{answer} \end{cases}$$

(c) The graph of  $f$  is increasing and concave up on  $0 < x < 1$  and  $4 < x < 6$  because  $f'(x) = g(x) > 0$  and  $f''(x) = g(x)$  is increasing on those intervals.

$$2 : \begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$$

(d) The graph of  $f$  has a point of inflection at  $x = 4$  because  $f''(x) = g(x)$  changes from decreasing to increasing at  $x = 4$ .

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$$

## 2018 SCORING GUIDELINES

## Question 4

$$(a) \quad H'(6) \approx \frac{H(7) - H(5)}{7 - 5} = \frac{11 - 6}{2} = \frac{5}{2}$$

$H'(6)$  is the rate at which the height of the tree is changing, in meters per year, at time  $t = 6$  years.

$$(b) \quad \frac{H(5) - H(3)}{5 - 3} = \frac{6 - 2}{2} = 2$$

Because  $H$  is differentiable on  $3 \leq t \leq 5$ ,  $H$  is continuous on  $3 \leq t \leq 5$ .

By the Mean Value Theorem, there exists a value  $c$ ,  $3 < c < 5$ , such that  $H'(c) = 2$ .

$$(c) \quad \text{The average height of the tree over the time interval } 2 \leq t \leq 10 \text{ is given by } \frac{1}{10 - 2} \int_2^{10} H(t) \, dt.$$

$$\begin{aligned} \frac{1}{8} \int_2^{10} H(t) \, dt &\approx \frac{1}{8} \left( \frac{1.5 + 2}{2} \cdot 1 + \frac{2 + 6}{2} \cdot 2 + \frac{6 + 11}{2} \cdot 2 + \frac{11 + 15}{2} \cdot 3 \right) \\ &= \frac{1}{8} (65.75) = \frac{263}{32} \end{aligned}$$

The average height of the tree over the time interval  $2 \leq t \leq 10$  is  $\frac{263}{32}$  meters.

$$(d) \quad G(x) = 50 \Rightarrow x = 1$$

$$\frac{d}{dt}(G(x)) = \frac{d}{dx}(G(x)) \cdot \frac{dx}{dt} = \frac{(1+x)100 - 100x \cdot 1}{(1+x)^2} \cdot \frac{dx}{dt} = \frac{100}{(1+x)^2} \cdot \frac{dx}{dt}$$

$$\left. \frac{d}{dt}(G(x)) \right|_{x=1} = \frac{100}{(1+1)^2} \cdot 0.03 = \frac{3}{4}$$

According to the model, the rate of change of the height of the tree with respect to time when the tree is 50 meters tall is  $\frac{3}{4}$  meter per year.

$$2 : \begin{cases} 1 : \text{estimate} \\ 1 : \text{interpretation with units} \end{cases}$$

$$2 : \begin{cases} 1 : \frac{H(5) - H(3)}{5 - 3} \\ 1 : \text{conclusion using Mean Value Theorem} \end{cases}$$

$$2 : \begin{cases} 1 : \text{trapezoidal sum} \\ 1 : \text{approximation} \end{cases}$$

$$3 : \begin{cases} 2 : \frac{d}{dt}(G(x)) \\ 1 : \text{answer} \end{cases}$$

Note: max 1/3 [1-0] if no chain rule

## 2018 SCORING GUIDELINES

## Question 5

$$(a) \quad \text{Area} = \frac{1}{2} \int_{\pi/3}^{5\pi/3} (4^2 - (3 + 2\cos\theta)^2) \, d\theta$$

$$(b) \quad \frac{dr}{d\theta} = -2\sin\theta \Rightarrow \left. \frac{dr}{d\theta} \right|_{\theta=\pi/2} = -2$$

$$r\left(\frac{\pi}{2}\right) = 3 + 2\cos\left(\frac{\pi}{2}\right) = 3$$

$$y = r\sin\theta \Rightarrow \frac{dy}{d\theta} = \frac{dr}{d\theta}\sin\theta + r\cos\theta$$

$$x = r\cos\theta \Rightarrow \frac{dx}{d\theta} = \frac{dr}{d\theta}\cos\theta - r\sin\theta$$

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/2} = \frac{dy/d\theta}{dx/d\theta} \Big|_{\theta=\pi/2} = \frac{-2\sin(\frac{\pi}{2}) + 3\cos(\frac{\pi}{2})}{-2\cos(\frac{\pi}{2}) - 3\sin(\frac{\pi}{2})} = \frac{2}{3}$$

The slope of the line tangent to the graph of  $r = 3 + 2\cos\theta$

at  $\theta = \frac{\pi}{2}$  is  $\frac{2}{3}$ .

— OR —

$$y = r\sin\theta = (3 + 2\cos\theta)\sin\theta \Rightarrow \frac{dy}{d\theta} = 3\cos\theta + 2\cos^2\theta - 2\sin^2\theta$$

$$x = r\cos\theta = (3 + 2\cos\theta)\cos\theta \Rightarrow \frac{dx}{d\theta} = -3\sin\theta - 4\sin\theta\cos\theta$$

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/2} = \frac{dy/d\theta}{dx/d\theta} \Big|_{\theta=\pi/2} = \frac{3\cos(\frac{\pi}{2}) + 2\cos^2(\frac{\pi}{2}) - 2\sin^2(\frac{\pi}{2})}{-3\sin(\frac{\pi}{2}) - 4\sin(\frac{\pi}{2})\cos(\frac{\pi}{2})} = \frac{2}{3}$$

The slope of the line tangent to the graph of  $r = 3 + 2\cos\theta$

at  $\theta = \frac{\pi}{2}$  is  $\frac{2}{3}$ .

$$(c) \quad \frac{dr}{dt} = \frac{dr}{d\theta} \cdot \frac{d\theta}{dt} = -2\sin\theta \cdot \frac{d\theta}{dt} \Rightarrow \frac{d\theta}{dt} = \frac{dr}{dt} \cdot \frac{1}{-2\sin\theta}$$

$$\left. \frac{d\theta}{dt} \right|_{\theta=\pi/3} = 3 \cdot \frac{1}{-2\sin(\frac{\pi}{3})} = \frac{3}{-\sqrt{3}} = -\sqrt{3} \text{ radians per second}$$

$$3 : \begin{cases} 1 : \text{constant and limits} \\ 2 : \text{integrand} \end{cases}$$

$$3 : \begin{cases} 1 : \frac{dy}{d\theta} = \frac{dr}{d\theta}\sin\theta + r\cos\theta \\ \quad \text{or } \frac{dx}{d\theta} = \frac{dr}{d\theta}\cos\theta - r\sin\theta \\ 1 : \frac{dy}{dx} = \frac{\frac{dr}{d\theta}\sin\theta + r\cos\theta}{\frac{dr}{d\theta}\cos\theta - r\sin\theta} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : \frac{dr}{dt} = \frac{dr}{d\theta} \cdot \frac{d\theta}{dt} \\ 1 : \frac{d\theta}{dt} = \frac{dr}{dt} \cdot \frac{1}{-2\sin\theta} \\ 1 : \text{answer with units} \end{cases}$$

## 2018 SCORING GUIDELINES

## Question 6

(a) The first four nonzero terms are  $\frac{x^2}{3} - \frac{x^3}{2 \cdot 3^2} + \frac{x^4}{3 \cdot 3^3} - \frac{x^5}{4 \cdot 3^4}$ .

The general term is  $(-1)^{n+1} \frac{x^{n+1}}{n \cdot 3^n}$ .

$$(b) \lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+2} x^{n+2}}{(n+1)(3^{n+1})}}{\frac{(-1)^{n+1} x^{n+1}}{n \cdot 3^n}} \right| = \lim_{n \rightarrow \infty} \left| \frac{-x}{3} \cdot \frac{n}{(n+1)} \right| = \left| \frac{x}{3} \right|$$

$$\left| \frac{x}{3} \right| < 1 \text{ for } |x| < 3$$

Therefore, the radius of convergence of the Maclaurin series for  $f$  is 3.

— OR —

The radius of convergence of the Maclaurin series for  $\ln(1+x)$  is 1, so the series for  $f(x) = x \ln\left(1 + \frac{x}{3}\right)$  converges absolutely for  $\left|\frac{x}{3}\right| < 1$ .

$$\left| \frac{x}{3} \right| < 1 \Rightarrow |x| < 3$$

Therefore, the radius of convergence of the Maclaurin series for  $f$  is 3.

When  $x = -3$ , the series is  $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{(-3)^{n+1}}{n \cdot 3^n} = \sum_{n=1}^{\infty} \frac{3}{n}$ , which

diverges by comparison to the harmonic series.

When  $x = 3$ , the series is  $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{3^{n+1}}{n \cdot 3^n} = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{3}{n}$ , which

converges by the alternating series test.

The interval of convergence of the Maclaurin series for  $f$  is  $-3 < x \leq 3$ .

(c) By the alternating series error bound, an upper bound for  $|P_4(2) - f(2)|$  is the magnitude of the next term of the alternating series.

$$|P_4(2) - f(2)| < \left| -\frac{2^5}{4 \cdot 3^4} \right| = \frac{8}{81}$$

2 :  $\begin{cases} 1 : \text{first four terms} \\ 1 : \text{general term} \end{cases}$

5 :  $\begin{cases} 1 : \text{sets up ratio} \\ 1 : \text{computes limit of ratio} \\ 1 : \text{radius of convergence} \\ 1 : \text{considers both endpoints} \\ 1 : \text{analysis and interval of convergence} \end{cases}$

— OR —

5 :  $\begin{cases} 1 : \text{radius for } \ln(1+x) \text{ series} \\ 1 : \text{substitutes } \frac{x}{3} \\ 1 : \text{radius of convergence} \\ 1 : \text{considers both endpoints} \\ 1 : \text{analysis and interval of convergence} \end{cases}$

2 :  $\begin{cases} 1 : \text{uses fifth-degree term as error bound} \\ 1 : \text{answer} \end{cases}$