


Part A (AB or BC): Graphing calculator not allowed**Question 4****9 points****General Scoring Notes**

- The model solution is presented using standard mathematical notation.
- Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be accurate to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

The continuous function f is defined on the closed interval $-6 \leq x \leq 12$. The graph of f , consisting of two semicircles and one line segment, is shown in the figure.

Graph of f

Let g be the function defined by $g(x) = \int_6^x f(t) dt$.

	Model Solution	Scoring
A	Find $g'(8)$. Give a reason for your answer.	
	$g'(x) = f(x)$	Considers $g'(x) = f(x)$ Point 1 (P1)
	$g'(8) = f(8) = 1$	Answer Point 2 (P2)

Scoring Notes for Part A

- P1** is earned for $g' = f$, $g'(x) = f(x)$, or $g'(8) = f(8)$ in part A.
- A response of $g'(8) = f(8) = 1$ earns both **P1** and **P2**.
- A response that does not earn **P1** can earn **P2** with an implied application of the Fundamental Theorem of Calculus (e.g., $g'(8) = 1$ or $f(8) = 1$).
- A response of $g'(8) = f(8) - f(6) = 1$ earns **P2** but not **P1**.

- B** Find all values of x in the open interval $-6 < x < 12$ at which the graph of g has a point of inflection. Give a reason for your answer.

The graph of g has a point of inflection where $g'' = f'$ changes sign, which is where $g' = f$ changes from decreasing to increasing or vice versa.

The graph of g has points of inflection at $x = -3$ and $x = 6$ because f changes from decreasing to increasing there.

The graph of g also has a point of inflection at $x = 3$ because f changes from increasing to decreasing there.

Answer

Point 3 (P3)

Reason

Point 4 (P4)

Scoring Notes for Part B

- **P3** is earned only for an answer of $x = -3$, $x = 3$, and $x = 6$. If any other/additional values of x in $-6 < x < 12$ are declared to be points of inflection, the response does not earn either **P3** or **P4**. Consideration of $x = -6$ or of $x = 12$ does not impact scoring.
- To earn **P4**, a response must tie the reason to the given graph of f .
 - A response of “ g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because f changes from increasing to decreasing or decreasing to increasing there” earns both **P3** and **P4**.
 - A response of “ g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because the slope of f changes sign there” earns both **P3** and **P4**.
 - A response of “ g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because f attains relative extrema there” earns both **P3** and **P4**.
 - A response of “ g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because g changes concavity there” earns **P3** but not **P4**.
 - A response of “ g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because $g'' = f'$ changes sign there” earns **P3** but not **P4**.
 - A response that relies upon an ambiguous term such as “the function” or “the graph” does not earn **P4**.
- **Special case:** A response with two of the three correct x -values with correct reasoning and no other/additional values of x declared to be points of inflection earns **P4** but not **P3**.

C Find $g(12)$ and $g(0)$. Label your answers.

$$g(12) = \int_6^{12} f(t) \, dt = \frac{1}{2} \cdot 6 \cdot 3 = 9$$

$g(12)$

Point 5 (P5)

$$g(0) = \int_6^0 f(x) \, dx = -\int_0^6 f(x) \, dx = -\frac{\pi}{2} 3^2 = -\frac{9\pi}{2}$$

$g(0)$

Point 6 (P6)

Scoring Notes for Part C

- Unlabeled values do not earn either **P5** or **P6**.
- **P5** is earned for a response of $g(12) = 9$, with or without supporting work.
- **P6** is earned for a response of $g(0) = -\frac{9\pi}{2}$, with or without supporting work.

Note: Incorrect communication between the label “ $g(0)$ ” and the answer will be treated as scratch work and will not impact scoring. For example, $g(0) = \int_0^6 f(x) \, dx = -\frac{9\pi}{2}$ earns **P6**.

D Find the value of x at which g attains an absolute minimum on the closed interval $-6 \leq x \leq 12$. Justify your answer.											
For $-6 \leq x \leq 12$, g attains a minimum either when $g'(x) = f(x) = 0$ or at an endpoint. $g'(x) = f(x) = 0$	Considers $g'(x) = 0$ Point 7 (P7)										
$\Rightarrow x = 0, x = 6$ <table border="1" data-bbox="183 560 343 795"> <thead> <tr> <th>x</th> <th>$g(x)$</th> </tr> </thead> <tbody> <tr> <td>-6</td> <td>0</td> </tr> <tr> <td>0</td> <td>$-\frac{9\pi}{2}$</td> </tr> <tr> <td>6</td> <td>0</td> </tr> <tr> <td>12</td> <td>9</td> </tr> </tbody> </table>	x	$g(x)$	-6	0	0	$-\frac{9\pi}{2}$	6	0	12	9	Justification Point 8 (P8)
x	$g(x)$										
-6	0										
0	$-\frac{9\pi}{2}$										
6	0										
12	9										
Therefore, on the closed interval $-6 \leq x \leq 12$, g attains an absolute minimum value at $x = 0$.	Answer Point 9 (P9)										

Scoring Notes for Part D

- **P7** is earned for considering $g'(x) = 0$ or $f(x) = 0$. **P7** is not earned by just presenting $x = 0$ and $x = 6$.
A response that discusses the sign of $g'(x)$ or $f(x)$ changing OR uses the phrase “critical points of g ” also earns **P7**.
- To earn **P8** using a candidates test, a response must make a global argument by providing evaluations or reasoning for each of $g(-6)$, $g(0)$, $g(6)$, and $g(12)$ (and no other x -values).
- Alternate justification and answer:
Because $g'(x) \leq 0$ (or $f(x) \leq 0$) for $-6 \leq x < 0$ and $g'(x) \geq 0$ (or $f(x) \geq 0$) for $0 < x \leq 12$, the absolute minimum of g occurs at $x = 0$.
- A response that presents a local argument (such as a First Derivative Test or a Second Derivative Test) or an incorrect global argument does not earn **P8** but is eligible for **P9** with the correct answer of $x = 0$.
- For **P8**, values of $g(0)$ and $g(12)$ can be imported from part C. A response can earn **P9** with an answer that is consistent with the imported values.

Part A (AB or BC): Graphing calculator required**Question 1****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

t (minutes)	0	3	7	12
$C(t)$ (degrees Celsius)	100	85	69	55

The temperature of coffee in a cup at time t minutes is modeled by a decreasing differentiable function C , where $C(t)$ is measured in degrees Celsius. For $0 \leq t \leq 12$, selected values of $C(t)$ are given in the table shown.

Model Solution**Scoring**

- (a) Approximate $C'(5)$ using the average rate of change of C over the interval $3 \leq t \leq 7$. Show the work that leads to your answer and include units of measure.

$$C'(5) \approx \frac{C(7) - C(3)}{7 - 3} = \frac{69 - 85}{4} = -4 \text{ degrees Celsius per minute}$$

Estimate with supporting work **1 point**

Units **1 point**

Scoring notes:

- To earn the first point a response must include a difference and a quotient as the supporting work.
- $\frac{-16}{7-3}$, $\frac{69-85}{7-3}$, or $\frac{69-85}{4}$ is sufficient to earn the first point.
- A response that presents only units without a numerical approximation for $C'(5)$ does not earn the second point.
- The second point is also earned for “degrees per minute” attached to a numerical value.

Total for part (a) 2 points

- (b) Use a left Riemann sum with the three subintervals indicated by the data in the table to approximate the value of $\int_0^{12} C(t) dt$. Interpret the meaning of $\frac{1}{12} \int_0^{12} C(t) dt$ in the context of the problem.

$\int_0^{12} C(t) dt \approx (3 - 0) \cdot C(0) + (7 - 3) \cdot C(3) + (12 - 7) \cdot C(7)$ $= 3 \cdot 100 + 4 \cdot 85 + 5 \cdot 69 = 985$	Form of left Riemann sum	1 point
	Estimate	1 point
$\frac{1}{12} \int_0^{12} C(t) dt$ is the average temperature of the coffee (in degrees Celsius) over the interval from $t = 0$ to $t = 12$.	Interpretation	1 point

Scoring notes:

- Read “=” as “ $\approx$ ” for the first point.
- To earn the first point at least five of the six factors in the Riemann sum must be correct. If any of the six factors is incorrect, the response does not earn the second point.
- A response of $(3 - 0) \cdot C(0) + (7 - 3) \cdot C(3) + (12 - 7) \cdot C(7)$ earns the first point. Values must be pulled from the table to earn the second point.
- A response of $3 \cdot 100 + 4 \cdot 85 + 5 \cdot 69$ earns both the first and second points, unless there is a subsequent error in simplification, in which case the response would earn only the first point.
- A completely correct right Riemann sum (e.g., $3 \cdot 85 + 4 \cdot 69 + 5 \cdot 55$) earns 1 of the first 2 points. An unsupported answer of 806 does not earn either of the first 2 points.
- Units will not affect scoring for the second point.
- To earn the third point the interpretation must include both “average temperature” and the time interval. The response need not include a reference to units. However, if incorrect units are given in the interpretation, the response does not earn the third point.

Total for part (b) 3 points

- (c) For $12 \leq t \leq 20$, the rate of change of the temperature of the coffee is modeled by $C'(t) = \frac{-24.55e^{0.01t}}{t}$, where $C'(t)$ is measured in degrees Celsius per minute. Find the temperature of the coffee at time $t = 20$. Show the setup for your calculations.

$C(20) = C(12) + \int_{12}^{20} C'(t) dt$	Integral	1 point
	Uses initial condition	1 point
$= 55 - 14.670812 = 40.329188$	Answer	1 point
The temperature of the coffee at time $t = 20$ is 40.329 degrees Celsius.		

Scoring notes:

- The first point is earned for a definite integral with integrand $C'(t)$. If the limits of integration are incorrect, the response does not earn the third point.
- A linkage error such as $C(20) = \int_{12}^{20} C'(t) dt = 55 - 14.670812$ or $\int_{12}^{20} C'(t) dt = -14.670812 = 40.329188$ earns the first 2 points but does not earn the third point.
- Missing differential (dt):
 - Unambiguous responses of $C(20) = C(12) + \int_{12}^{20} C'(t)$ or $C(20) = 55 + \int_{12}^{20} C'(t)$ earn the first 2 points and are eligible for the third point.
 - Ambiguous responses of $C(20) = \int_{12}^{20} C'(t) + C(12)$ or $C(20) = \int_{12}^{20} C'(t) + 55$ do not earn the first point, earn the second point, and earn the third point if the given numeric answer is correct. If there is no numeric answer given, these responses do not earn the third point.
- The second point is earned for adding $C(12)$ or 55 to a definite integral with a lower limit of 12, either symbolically or numerically.
- The third point is earned for an answer of $55 - 14.671$ or $-14.671 + 55$ with no additional simplification, provided there is some supporting work for these values.
- An answer of just 40.329 with no supporting work does not earn any points.

Total for part (c) 3 points

(d) For the model defined in part (c), it can be shown that $C''(t) = \frac{0.2455e^{0.01t}(100-t)}{t^2}$. For $12 < t < 20$, determine whether the temperature of the coffee is changing at a decreasing rate or at an increasing rate. Give a reason for your answer.		
Because $C''(t) > 0$ on the interval $12 < t < 20$, the rate of change in the temperature of the coffee, $C'(t)$, is increasing on this interval. That is, on the interval $12 < t < 20$, the temperature of the coffee is changing at an increasing rate.	Answer with reason	1 point

Scoring notes:

- This point is earned only for a correct answer with a correct reason that references the sign of the second derivative of C .
- A response that provides a reason based on the evaluation of $C''(t)$ at a single point does not earn this point.
- A response that uses ambiguous pronouns (such as “It is positive, so increasing”) does not earn this point.
- A response does not need to reference the interval $12 < t < 20$ to earn the point.

Total for part (d) 1 point**Total for question 1 9 points**

Part A (BC): Graphing calculator required**Question 2****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

A particle moving along a curve in the xy -plane has position $(x(t), y(t))$ at time t seconds, where $x(t)$ and $y(t)$ are measured in centimeters. It is known that $x'(t) = 8t - t^2$ and $y'(t) = -t + \sqrt{t^{1.2} + 20}$. At time $t = 2$ seconds, the particle is at the point $(3, 6)$.

	Model Solution	Scoring
(a)	Find the speed of the particle at time $t = 2$ seconds. Show the setup for your calculations.	
	$\sqrt{(x'(2))^2 + (y'(2))^2}$	Setup for speed 1 point
	$= 12.3048506$	Answer 1 point
	The speed of the particle at time $t = 2$ seconds is 12.305 (or 12.304) centimeters per second.	

Scoring notes:

- The first point is earned for the expression $\sqrt{(x'(2))^2 + (y'(2))^2}$, $\sqrt{(x'(t))^2 + (y'(t))^2}$, or equivalent.
- A response that presents just the exact answer, $\sqrt{144 + (-2 + \sqrt{2^{1.2} + 20})^2}$, earns both points.
- The second point is earned only for the answer 12.305 (or 12.304) regardless of whether the first point is earned.
- A response that includes a linkage error, such as $\sqrt{(x'(t))^2 + (y'(t))^2} = \sqrt{144 + (-2 + \sqrt{2^{1.2} + 20})^2}$ or $\sqrt{(x'(t))^2 + (y'(t))^2} = 12.305$ (or 12.304), earns at most 1 of the 2 points.
- Missing or incorrect units will not affect scoring in this part.

Total for part (a) 2 points

- (b) Find the total distance traveled by the particle over the time interval $0 \leq t \leq 2$. Show the setup for your calculations.

$\int_0^2 \sqrt{(x'(t))^2 + (y'(t))^2} dt$	Integral	1 point
$= 15.901715$	Answer	1 point
The total distance traveled by the particle over the time interval $0 \leq t \leq 2$ is 15.902 (or 15.901) centimeters.		

Scoring notes:

- The first point is earned only for an integral of $\int_0^2 \sqrt{(x'(t))^2 + (y'(t))^2} dt$ (or the mathematical equivalent), with or without the differential.
 - Note: $\int_0^2 \sqrt{x'(t)^2 + y'(t)^2} dt$ is not read as a parenthesis error.
- The second point is earned only for an answer of 15.902 (or 15.901), regardless of whether the first point is earned.
- Missing or incorrect units will not affect scoring in this part.

Total for part (b) 2 points

- (c) Find the y -coordinate of the position of the particle at the time $t = 0$. Show the setup for your calculations.

$y(0) = 6 + \int_2^0 y'(t) dt = 6 - 7.173613 = -1.173613$	Definite integral	1 point
	Uses initial condition	1 point
The y -coordinate of the position of the particle at time $t = 0$ is -1.174 (or -1.173).	Answer	1 point

Scoring notes:

- An answer of -1.174 (or -1.173) with no supporting work does not earn any points.
- The first point is earned for either of the definite integrals $\int_2^0 y'(t) dt$ or $\int_0^2 y'(t) dt$.
- The second point is earned for any of:
 - $y(2) \pm \int_2^0 y'(t) dt$, $6 \pm \int_2^0 y'(t) dt$,
 - $y(2) \pm \int_0^2 y'(t) dt$, $6 \pm \int_0^2 y'(t) dt$,
 - $y(2) \pm 7.173613$, or 6 ± 7.173613 .
- A response that attempts to evaluate $\int y'(t) dt$ does not earn the first or the third point.
 - Such a response can earn the second point by attempting to solve for the constant of integration by presenting an expression as an antiderivative for $y'(t)$, evaluating this expression at $t = 2$, and setting this expression equal to 6.

- A response that reverses the limits of integration, e.g., $y(0) = 6 + \int_0^2 y'(t) dt$ or $6 + 7.173613$, earns the second point but does not earn the third point.
- In order to earn the third point, a response must have earned at least 1 of the first 2 points.
- A response containing any linkage error can earn at most 2 of the 3 points. For example:
 - Equating two unequal quantities: $\int_2^0 y'(t) dt = -1.174$, $\int_2^0 y'(t) dt = 6 - 7.173613$,
 $6 + \int_0^2 y'(t) dt = 6 - 7.173613$, or $6 + \int_0^2 y'(t) dt = 6 + 7.173613 = -1.173613$
 - Equating an expression to a numerical value: $y(t) = 6 + \int_2^0 y'(t) dt = -1.174$
- Missing differentials (dt):
 - Unambiguous responses of $y(2) + \int_2^0 y'(t)$, $y(2) - \int_0^2 y'(t)$, $6 + \int_2^0 y'(t)$, or $6 - \int_0^2 y'(t)$ earn the first 2 points and would earn the third point for the correct numerical answer.
 - Unambiguous responses of $y(2) + \int_0^2 y'(t)$ or $6 + \int_0^2 y'(t)$ with reversed limits of integration and missing differential earn the first 2 points but cannot earn the third point.
 - Ambiguous responses of $\int_2^0 y'(t) + y(2)$, $-\int_0^2 y'(t) + y(2)$, $\int_2^0 y'(t) + 6$, or $-\int_0^2 y'(t) + 6$ earn the first point, do not earn the second point, but do earn the third point if a correct numeric answer is provided. If no numeric answer is given, none of these responses earn the third point.
 - Ambiguous responses of $\int_0^2 y'(t) + y(2)$ or $\int_0^2 y'(t) + 6$ with reversed limits of integration and no differential earn 1 out of 3 points.
- If a response provides work for both the x - and y -coordinates, the work for the x -coordinate will not affect scoring.
- However, a response that reports only a completely correct x -coordinate of the particle's position at time $t = 0$ with all supporting work, e.g., $x(0) = 3 + \int_2^0 x'(t) dt = -\frac{31}{3} = -10.333$, earns 2 out of 3 points.

Total for part (c) 3 points

- (d) For $2 \leq t \leq 8$, the particle remains in the first quadrant. Find all times t in the interval $2 \leq t \leq 8$ when the particle is moving toward the x -axis. Give a reason for your answer.

Because $y(t) > 0$ when $2 \leq t \leq 8$, the particle will be moving toward the x -axis when $y'(t) < 0$. This occurs when 5.222 (or 5.221) $< t < 8$.	Considers sign of $y'(t)$	1 point
	Answer with reason	1 point

Scoring notes:

- The first point can be earned by stating $y'(t) = 0$, $y'(t) < 0$, $y'(t) > 0$, or $t = 5.222$ (or 5.221).
Note: $y'(t)$ may be written as $\frac{dy}{dt}$.
- The second point cannot be earned without the first.
- To earn the second point, a response must identify the correct interval (and no additional intervals in $[2, 8]$) and explicitly state the need for $y'(t) < 0$. The interval can be open, closed, or half-open.


Total for part (d) 2 points

Total for question 2 9 points

Part B (AB or BC): Graphing calculator not allowed**Question 4****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.



The graph of the differentiable function f , shown for $-6 \leq x \leq 7$, has a horizontal tangent at $x = -2$ and is linear for $0 \leq x \leq 7$. Let R be the region in the second quadrant bounded by the graph of f , the vertical line $x = -6$, and the x - and y -axes. Region R has area 12.

Model Solution**Scoring**

- (a) The function g is defined by $g(x) = \int_0^x f(t) dt$. Find the values of $g(-6)$, $g(4)$, and $g(6)$.

$g(-6) = \int_0^{-6} f(t) dt = -\int_{-6}^0 f(t) dt = -12$	$g(-6)$	1 point
$g(4) = \int_0^4 f(t) dt = \frac{1}{2} \cdot 4 \cdot 2 = 4$	$g(4)$	1 point
$g(6) = \int_0^6 f(t) dt = \frac{1}{2} \cdot 4 \cdot 2 - \frac{1}{2} \cdot 2 \cdot 1 = 3$	$g(6)$	1 point

Scoring notes:

- Supporting work is not required for any of these values. However, any supporting work that is shown must be correct to earn the corresponding point.
- Special case: A response that explicitly presents $g(x) = \int_{-6}^x f(t) dt$ does not earn the first point it would have otherwise earned. The response is eligible for all subsequent points for correct answers, or for consistent answers with supporting work.
 - Note: $\int_{-6}^{-6} f(t) dt = 0$, $\int_{-6}^4 f(t) dt = 16$, $\int_{-6}^6 f(t) dt = 15$
- Labeled values may be presented in any order. Unlabeled values are read from left to right and from top to bottom as $g(-6)$, $g(4)$, and $g(6)$, respectively. A response that presents only 1 or 2 values must label them to earn any points.

Total for part (a) 3 points

- (b) For the function g defined in part (a), find all values of x in the interval $0 \leq x \leq 6$ at which the graph of g has a critical point. Give a reason for your answer.

$g'(x) = f(x)$	Fundamental Theorem of Calculus	1 point
$g'(x) = f(x) = 0 \Rightarrow x = 4$	Answer with reason	1 point
Therefore, the graph of g has a critical point at $x = 4$.		

Scoring notes:

- The first point is earned for explicitly making the connection $g' = f$ in this part.
 - A response that writes $g'' = f'$ earns the first point but can only earn the second point by reasoning from $f = 0$.
- A response that does not earn the first point is eligible to earn the second point with an implied application of the FTC (e.g., “Because $g'(4) = 0$, $x = 4$ is a critical point”).
- A response that reports any additional critical points in $0 < x < 6$ does not earn the second point.
 - Any presented critical point outside the interval $0 < x < 6$ will not affect scoring.

Total for part (b) 2 points

- (c) The function h is defined by $h(x) = \int_{-6}^x f'(t) dt$. Find the values of $h(6)$, $h'(6)$, and $h''(6)$. Show the work that leads to your answers.

$h(6) = \int_{-6}^6 f'(t) dt = f(6) - f(-6) = -1 - 0.5 = -1.5$	Uses Fundamental Theorem of Calculus	1 point
	$h(6)$ with supporting work	1 point
$h'(x) = f'(x)$, so $h'(6) = f'(6) = -\frac{1}{2}$.	$h'(6)$	1 point
$h''(x) = f''(x)$, so $h''(6) = f''(6) = 0$.	$h''(6)$	1 point

Scoring notes:

- Labeled values may be presented in any order.
- Unlabeled values are read from left to right and from top to bottom as $h(6)$, $h'(6)$, and $h''(6)$, respectively. A response that presents only 1 or 2 values must label them in order to earn any points.
- A response of $h(6) = -1.5$ does not earn either of the first 2 points. A response of $h(6) = f(6) - f(-6)$ earns the first point but not yet the second point.
- A response of $h(6) = -1 - 0.5$ is the minimum work required to earn both of the first 2 points.
- To earn the third point a response must state either $h'(x) = f'(x)$ or $h'(6) = f'(6)$, and provide an answer of $-\frac{1}{2}$.
- The fourth point is earned for a response of $h''(6) = 0$, with or without supporting work.
- A response that has one or more linkage errors does not earn the first point it would have otherwise earned. For example, $h'(x) = f'(6) = -\frac{1}{2}$ does not earn the third point but is eligible for the fourth point even in the presence of another linkage error, such as $h''(x) = f''(6) = 0$.


Total for part (c) 4 points

Total for question 4 9 points

Part B (AB or BC): Graphing calculator not allowed**Question 3****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.



Let f be a differentiable function with $f(4) = 3$. On the interval $0 \leq x \leq 7$, the graph of f' , the derivative of f , consists of a semicircle and two line segments, as shown in the figure above.

Model Solution**Scoring**

(a) Find $f(0)$ and $f(5)$.

$$f(0) = f(4) + \int_4^0 f'(x) dx = 3 - \int_0^4 f'(x) dx = 3 + 2\pi$$

$$f(5) = f(4) + \int_4^5 f'(x) dx = 3 + \frac{1}{2} = \frac{7}{2}$$

Area of either region **1 point**

– OR – $\int_0^4 f'(x) dx$

– OR – $\int_4^5 f'(x) dx$

$f(0)$ **1 point**

$f(5)$ **1 point**

Scoring notes:

- A response with answers of only $f(0) = \pm 2\pi$, or only $f(5) = \frac{1}{2}$, or both earns 1 of the 3 points.
- A response displaying $f(5) = \frac{7}{2}$ and a missing or incorrect value for $f(0)$ earns 2 of the 3 points.
- The second and third points can be earned in either order.
- Read unlabeled values from left to right and from top to bottom as $f(0)$ and $f(5)$. A single value must be labeled as either $f(0)$ or $f(5)$ in order to earn any points.

Total for part (a) **2 points**

- (b) Find the x -coordinates of all points of inflection of the graph of f for $0 < x < 7$. Justify your answer.

The graph of f has a point of inflection at each of $x = 2$ and $x = 6$, because $f'(x)$ changes from decreasing to increasing at $x = 2$ and from increasing to decreasing at $x = 6$.	Answer	1 point
	Justification	1 point

Scoring notes:

- A response that gives only one of $x = 2$ or $x = 6$, along with a correct justification, earns 1 of the 2 points.
- A response that claims that there is a point of inflection at any value other than $x = 2$ or $x = 6$ earns neither point.
- To earn the second point a response must use correct reasoning based on the graph of f' . Examples of correct reasoning include:
 - Correctly discussing the signs of the slopes of the graph of f'
 - Citing $x = 2$ and $x = 6$ as the locations of local extrema on the graph of f'
- Examples of reasoning not (sufficiently) connected to the graph of f' include:
 - Reasoning based on sign changes in f'' unless the connection is made between the sign of f'' and the slopes of the graph of f'
 - Reasoning based only on the concavity of the graph of f
- The second point cannot be earned by use of vague or undefined terms such as “it” or “the function” or “the derivative.”
- Responses that report inflection points as ordered pairs must report the points $(2, 3 + \pi)$ and $(6, 5)$ in order to earn the first point. If the y -coordinates are reported incorrectly, the response remains eligible for the second point.

Total for part (b) 2 points

- (c) Let g be the function defined by $g(x) = f(x) - x$. On what intervals, if any, is g decreasing for $0 \leq x \leq 7$? Show the analysis that leads to your answer.

$g'(x) = f'(x) - 1$	$g'(x) = f'(x) - 1$	1 point
$f'(x) - 1 \leq 0 \Rightarrow f'(x) \leq 1$	Interval with reason	1 point
The graph of g is decreasing on the interval $0 \leq x \leq 5$ because $g'(x) \leq 0$ on this interval.		

Scoring notes:

- The first point can be earned for $f'(x) \leq 1$ or the equivalent, in words or symbols.
- Endpoints do not need to be included in the interval to be eligible for the second point.

Total for part (c) 2 points

- (d) For the function g defined in part (c), find the absolute minimum value on the interval $0 \leq x \leq 7$.

Justify your answer.

g is continuous, $g'(x) < 0$ for $0 < x < 5$, and $g'(x) > 0$ for $5 < x < 7$. Therefore, the absolute minimum occurs at $x = 5$, and $g(5) = f(5) - 5 = \frac{7}{2} - 5 = -\frac{3}{2}$ is the absolute minimum value of g .	Considers $g'(x) = 0$	1 point
	Answer with justification	1 point

Scoring notes:

- A justification that uses a local argument, such as “ g' changes from negative to positive (or g changes from decreasing to increasing) at $x = 5$ ” must also state that $x = 5$ is the only critical point.
- If $g'(x) = 0$ (or equivalent) is not declared explicitly, a response that isolates $x = 5$ as the only critical number belonging to $(0, 7)$ earns the first point.
- A response that imports $g'(x) = f'(x)$ from part (c) is eligible for the first point but not the second.
 - In this case, consideration of $x = 4$ as the only critical number on $(0, 7)$ earns the first point.
- Solution using Candidates Test:

$$g'(x) = f'(x) - 1 = 0 \Rightarrow x = 5, x = 7$$

x	$g(x)$
0	$3 + 2\pi$
5	$-\frac{3}{2}$
7	$-\frac{1}{2}$

The absolute minimum value of g on the interval $0 \leq x \leq 7$ is $-\frac{3}{2}$.

- When using a Candidates Test, a response may import an incorrect value of $f(0) = g(0) > -\frac{3}{2}$ from part (a). The second point can only be earned for an answer of $-\frac{3}{2}$.

Total for part (d) 2 points

Total for question 3 9 points

Part B (BC): Graphing calculator not allowed**Question 5****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.




Figure 1




Figure 2

Figures 1 and 2, shown above, illustrate regions in the first quadrant associated with the graphs of $y = \frac{1}{x}$ and $y = \frac{1}{x^2}$, respectively. In Figure 1, let R be the region bounded by the graph of $y = \frac{1}{x}$, the x -axis, and the vertical lines $x = 1$ and $x = 5$. In Figure 2, let W be the unbounded region between the graph of $y = \frac{1}{x^2}$ and the x -axis that lies to the right of the vertical line $x = 3$.

Model Solution**Scoring**

- (a) Find the area of region R .

Area = $\int_1^5 \frac{1}{x} dx$	Integral	1 point
$= \ln x \Big _1^5$	Answer	1 point
$= \ln 5 - \ln 1 = \ln 5$		

Scoring notes:

- A definite integral with incorrect bounds does not earn either point.
- An unevaluated indefinite integral does not earn either point.
- An indefinite integral that is evaluated in a later step may earn one or both points. For example,
 $\int \frac{1}{x} dx = \ln 5 - \ln 1$ (or $\ln 5$) does not earn the first point but does earn the second. However,
 $\int \frac{1}{x} dx = \ln x + C \Rightarrow \text{Area} = \ln 5 - \ln 1$ earns both points.

Total for part (a) 2 points

- (b) Region R is the base of a solid. For the solid, at each x the cross section perpendicular to the x -axis is a rectangle with area given by $xe^{x/5}$. Find the volume of the solid.

Volume = $\int_1^5 xe^{x/5} dx$	Definite integral	1 point
Using integration by parts, $u = x \quad dv = e^{x/5} dx$ $du = dx \quad v = 5e^{x/5}$	u and dv	1 point
$\int xe^{x/5} dx = 5xe^{x/5} - \int 5e^{x/5} dx$ $= 5xe^{x/5} - 25e^{x/5} + C$ $= 5e^{x/5}(x - 5) + C$	$\int xe^{x/5} dx$ $= 5xe^{x/5} - \int 5e^{x/5} dx$	1 point
Volume = $5e^{x/5}(x - 5) \Big _1^5$ $= 5e(0) - 5e^{1/5}(-4) = 20e^{1/5}$	Answer	1 point

Scoring notes:

- The first point is earned for $c \int_1^5 xe^{x/5} dx$, where $c \neq 0$. Errors of $c \neq 1$, for example $c = \pi$, will not earn the fourth point.
- Incorrect integrals that require integration by parts are still eligible for the second and third points. Both of these points will be earned with at least one correct application of integration by parts.
- The second point will be earned with an implied u and dv in the presence of $5xe^{x/5} - \int 5e^{x/5} dx$.
- The tabular method may be used to show integration by parts. In this case, the second point is earned by having columns (labeled or unlabeled) that begin with x and $e^{x/5}$. The third point is earned for either $5xe^{x/5} - \int 5e^{x/5} dx$ or $5xe^{x/5} - 25e^{x/5}$.
- Limits of integration may be present, omitted, or partially present in the work for the second and third points.
- The fourth point is earned only for the correct answer.

Total for part (b) 4 points

- (c) Find the volume of the solid generated when the unbounded region W is revolved about the x -axis.

Volume = $\pi \int_3^{\infty} \left(\frac{1}{x^2}\right)^2 dx = \pi \lim_{b \rightarrow \infty} \int_3^b \frac{1}{x^4} dx$	Improper integral	1 point
$= \pi \lim_{b \rightarrow \infty} \left(\frac{1}{-3x^3} \Big _3^b \right)$	Antiderivative	1 point
$= \pi \lim_{b \rightarrow \infty} \left(\frac{1}{-3} \right) \left[\frac{1}{b^3} - \frac{1}{3^3} \right]$ $= \pi \left(\frac{1}{-3} \right) \left(0 - \frac{1}{3^3} \right) = \frac{\pi}{81}$	Answer	1 point

Scoring notes:

- The first point is earned for either $c \int_3^{\infty} \left(\frac{1}{x^2}\right)^2 dx$ or $\lim_{b \rightarrow \infty} c \int_3^b \frac{1}{x^4} dx$, where $c \neq 0$. Errors of $c \neq \pi$ will not earn the third point.
- The second point is earned for a correct antiderivative of any integrand of the form $\frac{1}{x^n}$, for any integer $n \geq 2$.
- To earn the answer point, a response must use correct limit notation and cannot include arithmetic with infinity, such as $\frac{1}{\infty^3}$.

Total for part (c) 3 points

Total for question 5 9 points

2019

AP

CollegeBoard

AP[®] Calculus BC

Scoring Guidelines

2019 SCORING GUIDELINES

Question 1

(a) $\int_0^5 E(t) dt = 153.457690$

To the nearest whole number, 153 fish enter the lake from midnight to 5 A.M.

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

(b) $\frac{1}{5-0} \int_0^5 L(t) dt = 6.059038$

The average number of fish that leave the lake per hour from midnight to 5 A.M. is 6.059 fish per hour.

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

- (c) The rate of change in the number of fish in the lake at time t is given by $E(t) - L(t)$.

$$E(t) - L(t) = 0 \Rightarrow t = 6.20356$$

$E(t) - L(t) > 0$ for $0 \leq t < 6.20356$, and $E(t) - L(t) < 0$ for $6.20356 < t \leq 8$. Therefore the greatest number of fish in the lake is at time $t = 6.204$ (or 6.203).

$$3 : \begin{cases} 1 : \text{sets } E(t) - L(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

— OR —

Let $A(t)$ be the change in the number of fish in the lake from midnight to t hours after midnight.

$$A(t) = \int_0^t (E(s) - L(s)) ds$$

$$A'(t) = E(t) - L(t) = 0 \Rightarrow t = C = 6.20356$$

t	$A(t)$
0	0
C	135.01492
8	80.91998

Therefore the greatest number of fish in the lake is at time $t = 6.204$ (or 6.203).

(d) $E'(5) - L'(5) = -10.7228 < 0$

Because $E'(5) - L'(5) < 0$, the rate of change in the number of fish is decreasing at time $t = 5$.

$$2 : \begin{cases} 1 : \text{considers } E'(5) \text{ and } L'(5) \\ 1 : \text{answer with explanation} \end{cases}$$

2019 SCORING GUIDELINES

Question 2

$$(a) \quad \frac{1}{2} \int_0^{\sqrt{\pi}} (r(\theta))^2 d\theta = 3.534292$$

The area of S is 3.534.

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

$$(b) \quad \frac{1}{\sqrt{\pi} - 0} \int_0^{\sqrt{\pi}} r(\theta) d\theta = 1.579933$$

The average distance from the origin to a point on the curve $r = r(\theta)$ for $0 \leq \theta \leq \sqrt{\pi}$ is 1.580 (or 1.579).

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

$$(c) \quad \tan \theta = \frac{y}{x} = m \Rightarrow \theta = \tan^{-1} m$$

$$\frac{1}{2} \int_0^{\tan^{-1} m} (r(\theta))^2 d\theta = \frac{1}{2} \left(\frac{1}{2} \int_0^{\sqrt{\pi}} (r(\theta))^2 d\theta \right)$$

$$3 : \begin{cases} 1 : \text{equates polar areas} \\ 1 : \text{inverse trigonometric function} \\ \text{applied to } m \text{ as limit of} \\ \text{integration} \\ 1 : \text{equation} \end{cases}$$

(d) As $k \rightarrow \infty$, the circle $r = k \cos \theta$ grows to enclose all points to the right of the y -axis.

$$2 : \begin{cases} 1 : \text{limits of integration} \\ 1 : \text{answer with integral} \end{cases}$$

$$\begin{aligned} \lim_{k \rightarrow \infty} A(k) &= \frac{1}{2} \int_0^{\pi/2} (r(\theta))^2 d\theta \\ &= \frac{1}{2} \int_0^{\pi/2} (3\sqrt{\theta} \sin(\theta^2))^2 d\theta = 3.324 \end{aligned}$$

2019 SCORING GUIDELINES

Question 3

$$\begin{aligned}
 \text{(a)} \quad \int_{-6}^5 f(x) \, dx &= \int_{-6}^{-2} f(x) \, dx + \int_{-2}^5 f(x) \, dx \\
 &\Rightarrow 7 = \int_{-6}^{-2} f(x) \, dx + 2 + \left(9 - \frac{9\pi}{4}\right) \\
 &\Rightarrow \int_{-6}^{-2} f(x) \, dx = 7 - \left(11 - \frac{9\pi}{4}\right) = \frac{9\pi}{4} - 4
 \end{aligned}$$

$$\begin{aligned}
 \text{(b)} \quad \int_3^5 (2f'(x) + 4) \, dx &= 2 \int_3^5 f'(x) \, dx + \int_3^5 4 \, dx \\
 &= 2(f(5) - f(3)) + 4(5 - 3) \\
 &= 2(0 - (3 - \sqrt{5})) + 8 \\
 &= 2(-3 + \sqrt{5}) + 8 = 2 + 2\sqrt{5}
 \end{aligned}$$

— OR —

$$\begin{aligned}
 \int_3^5 (2f'(x) + 4) \, dx &= [2f(x) + 4x]_{x=3}^{x=5} \\
 &= (2f(5) + 20) - (2f(3) + 12) \\
 &= (2 \cdot 0 + 20) - (2(3 - \sqrt{5}) + 12) \\
 &= 2 + 2\sqrt{5}
 \end{aligned}$$

$$\text{(c)} \quad g'(x) = f(x) = 0 \Rightarrow x = -1, x = \frac{1}{2}, x = 5$$

x	$g(x)$
-2	0
-1	$\frac{1}{2}$
$\frac{1}{2}$	$-\frac{1}{4}$
5	$11 - \frac{9\pi}{4}$

On the interval $-2 \leq x \leq 5$, the absolute maximum value of g is $g(5) = 11 - \frac{9\pi}{4}$.

$$\begin{aligned}
 \text{(d)} \quad \lim_{x \rightarrow 1} \frac{10^x - 3f'(x)}{f(x) - \arctan x} &= \frac{10^1 - 3f'(1)}{f(1) - \arctan 1} \\
 &= \frac{10 - 3 \cdot 2}{1 - \arctan 1} = \frac{4}{1 - \frac{\pi}{4}}
 \end{aligned}$$

$$3 : \begin{cases} 1 : \int_{-6}^5 f(x) \, dx = \int_{-6}^{-2} f(x) \, dx + \int_{-2}^5 f(x) \, dx \\ 1 : \int_{-2}^5 f(x) \, dx \\ 1 : \text{answer} \end{cases}$$

$$2 : \begin{cases} 1 : \text{Fundamental Theorem of Calculus} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : g'(x) = f(x) \\ 1 : \text{identifies } x = -1 \text{ as a candidate} \\ 1 : \text{answer with justification} \end{cases}$$

$$1 : \text{answer}$$

2019 SCORING GUIDELINES

Question 4

(a) $V = \pi r^2 h = \pi(1)^2 h = \pi h$
 $\left. \frac{dV}{dt} \right|_{h=4} = \pi \left. \frac{dh}{dt} \right|_{h=4} = \pi \left(-\frac{1}{10} \sqrt{4} \right) = -\frac{\pi}{5}$ cubic feet per second

$$2 : \begin{cases} 1 : \frac{dV}{dt} = \pi \frac{dh}{dt} \\ 1 : \text{answer with units} \end{cases}$$

(b) $\frac{d^2 h}{dt^2} = -\frac{1}{20\sqrt{h}} \cdot \frac{dh}{dt} = -\frac{1}{20\sqrt{h}} \cdot \left(-\frac{1}{10} \sqrt{h} \right) = \frac{1}{200}$
 Because $\frac{d^2 h}{dt^2} = \frac{1}{200} > 0$ for $h > 0$, the rate of change of the height is increasing when the height of the water is 3 feet.

$$3 : \begin{cases} 1 : \frac{d}{dh} \left(-\frac{1}{10} \sqrt{h} \right) = -\frac{1}{20\sqrt{h}} \\ 1 : \frac{d^2 h}{dt^2} = -\frac{1}{20\sqrt{h}} \cdot \frac{dh}{dt} \\ 1 : \text{answer with explanation} \end{cases}$$

(c) $\frac{dh}{\sqrt{h}} = -\frac{1}{10} dt$
 $\int \frac{dh}{\sqrt{h}} = \int -\frac{1}{10} dt$
 $2\sqrt{h} = -\frac{1}{10} t + C$
 $2\sqrt{5} = -\frac{1}{10} \cdot 0 + C \Rightarrow C = 2\sqrt{5}$
 $2\sqrt{h} = -\frac{1}{10} t + 2\sqrt{5}$
 $h(t) = \left(-\frac{1}{20} t + \sqrt{5} \right)^2$

$$4 : \begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ \text{and uses initial condition} \\ 1 : h(t) \end{cases}$$

Note: 0/4 if no separation of variables

Note: max 2/4 [1-1-0-0] if no constant of integration

2019 SCORING GUIDELINES

Question 5

$$(a) \quad f'(x) = \frac{-(2x-2)}{(x^2-2x+k)^2}$$

$$f'(0) = \frac{2}{k^2} = 6 \Rightarrow k^2 = \frac{1}{3} \Rightarrow k = \frac{1}{\sqrt{3}}$$

3 : $\begin{cases} 1 : \text{denominator of } f'(x) \\ 1 : f'(x) \\ 1 : \text{answer} \end{cases}$

$$(b) \quad \frac{1}{x^2-2x-8} = \frac{1}{(x-4)(x+2)} = \frac{A}{x-4} + \frac{B}{x+2}$$

$$\Rightarrow 1 = A(x+2) + B(x-4)$$

$$\Rightarrow A = \frac{1}{6}, B = -\frac{1}{6}$$

3 : $\begin{cases} 1 : \text{partial fraction decomposition} \\ 1 : \text{antiderivatives} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \int_0^1 f(x) dx &= \int_0^1 \left(\frac{\frac{1}{6}}{x-4} - \frac{\frac{1}{6}}{x+2} \right) dx \\ &= \left[\frac{1}{6} \ln|x-4| - \frac{1}{6} \ln|x+2| \right]_{x=0}^{x=1} \\ &= \left(\frac{1}{6} \ln 3 - \frac{1}{6} \ln 3 \right) - \left(\frac{1}{6} \ln 4 - \frac{1}{6} \ln 2 \right) = -\frac{1}{6} \ln 2 \end{aligned}$$

$$\begin{aligned} (c) \quad \int_0^2 \frac{1}{x^2-2x+1} dx &= \int_0^2 \frac{1}{(x-1)^2} dx = \int_0^1 \frac{1}{(x-1)^2} dx + \int_1^2 \frac{1}{(x-1)^2} dx \\ &= \lim_{b \rightarrow 1^-} \int_0^b \frac{1}{(x-1)^2} dx + \lim_{b \rightarrow 1^+} \int_b^2 \frac{1}{(x-1)^2} dx \\ &= \lim_{b \rightarrow 1^-} \left(-\frac{1}{x-1} \Big|_{x=0}^{x=b} \right) + \lim_{b \rightarrow 1^+} \left(-\frac{1}{x-1} \Big|_{x=b}^{x=2} \right) \\ &= \lim_{b \rightarrow 1^-} \left(-\frac{1}{b-1} - 1 \right) + \lim_{b \rightarrow 1^+} \left(-1 + \frac{1}{b-1} \right) \end{aligned}$$

Because $\lim_{b \rightarrow 1^-} \left(-\frac{1}{b-1} \right)$ does not exist, the integral diverges.

3 : $\begin{cases} 1 : \text{improper integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer with reason} \end{cases}$

2019 SCORING GUIDELINES

Question 6

(a) $f(0) = 3$ and $f'(0) = -2$

The third-degree Taylor polynomial for f about $x = 0$ is

$$3 - 2x + \frac{3}{2!}x^2 + \frac{-2}{3!}x^3 = 3 - 2x + \frac{3}{2}x^2 - \frac{2}{6}x^3.$$

(b) The first three nonzero terms of the Maclaurin series for e^x are

$$1 + x + \frac{1}{2!}x^2.$$

The second-degree Taylor polynomial for $e^x f(x)$ about $x = 0$ is

$$\begin{aligned} 3\left(1 + x + \frac{1}{2!}x^2\right) - 2x(1 + x) + \frac{3}{2}x^2(1) \\ = 3 + (3 - 2)x + \left(\frac{3}{2} - 2 + \frac{3}{2}\right)x^2 \\ = 3 + x + x^2. \end{aligned}$$

(c) $h(1) = \int_0^1 f(t) dt$

$$\begin{aligned} &\approx \int_0^1 \left(3 - 2t + \frac{3}{2}t^2 - \frac{23}{12}t^3\right) dt \\ &= \left[3t - t^2 + \frac{1}{2}t^3 - \frac{23}{48}t^4\right]_{t=0}^{t=1} \\ &= 3 - 1 + \frac{1}{2} - \frac{23}{48} = \frac{97}{48} \end{aligned}$$

(d) The alternating series error bound is the absolute value of the first omitted term of the series for $h(1)$.

$$\int_0^1 \left(\frac{54}{4!}t^4\right) dt = \left[\frac{9}{20}t^5\right]_{t=0}^{t=1} = \frac{9}{20}$$

$$\text{Error} \leq \left|\frac{9}{20}\right| = 0.45$$

$$2 : \begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \end{cases}$$

$$2 : \begin{cases} 1 : \text{three terms for } e^x \\ 1 : \text{three terms for } e^x f(x) \end{cases}$$

$$2 : \begin{cases} 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : \text{uses fourth-degree term} \\ \quad \text{of Maclaurin series for } f \\ 1 : \text{uses first omitted term} \\ \quad \text{of series for } h(1) \\ 1 : \text{error bound} \end{cases}$$

2018**AP[®]** **CollegeBoard**

AP Calculus BC

Scoring Guidelines

2018 SCORING GUIDELINES

Question 1

(a) $\int_0^{300} r(t) dt = 270$

According to the model, 270 people enter the line for the escalator during the time interval $0 \leq t \leq 300$.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) $20 + \int_0^{300} (r(t) - 0.7) dt = 20 + \int_0^{300} r(t) dt - 0.7 \cdot 300 = 80$

According to the model, 80 people are in line at time $t = 300$.

2 : $\begin{cases} 1 : \text{considers rate out} \\ 1 : \text{answer} \end{cases}$

(c) Based on part (b), the number of people in line at time $t = 300$ is 80.

The first time t that there are no people in line is

$$300 + \frac{80}{0.7} = 414.286 \text{ (or 414.285) seconds.}$$

1 : answer

(d) The total number of people in line at time t , $0 \leq t \leq 300$, is modeled by

$$20 + \int_0^t r(x) dx - 0.7t.$$

$$r(t) - 0.7 = 0 \Rightarrow t_1 = 33.013298, t_2 = 166.574719$$

4 : $\begin{cases} 1 : \text{considers } r(t) - 0.7 = 0 \\ 1 : \text{identifies } t = 33.013 \\ 1 : \text{answers} \\ 1 : \text{justification} \end{cases}$

t	People in line for escalator
0	20
t_1	3.803
t_2	158.070
300	80

The number of people in line is a minimum at time $t = 33.013$ seconds, when there are 4 people in line.

2018 SCORING GUIDELINES

Question 2

(a) $p'(25) = -1.179$

At a depth of 25 meters, the density of plankton cells is changing at a rate of -1.179 million cells per cubic meter per meter.

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{meaning with units} \end{cases}$$

(b) $\int_0^{30} 3p(h) \, dh = 1675.414936$

There are 1675 million plankton cells in the column of water between $h = 0$ and $h = 30$ meters.

$$2 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$$

(c) $\int_{30}^K 3f(h) \, dh$ represents the number of plankton cells, in millions, in the column of water from a depth of 30 meters to a depth of K meters.

The number of plankton cells, in millions, in the entire column of water is given by $\int_0^{30} 3p(h) \, dh + \int_{30}^K 3f(h) \, dh$.

Because $0 \leq f(h) \leq u(h)$ for all $h \geq 30$,

$$3 \int_{30}^K f(h) \, dh \leq 3 \int_{30}^K u(h) \, dh \leq 3 \int_{30}^{\infty} u(h) \, dh = 3 \cdot 105 = 315.$$

The total number of plankton cells in the column of water is bounded by $1675.415 + 315 = 1990.415 \leq 2000$ million.

$$3 : \begin{cases} 1 : \text{integral expression} \\ 1 : \text{compares improper integral} \\ 1 : \text{explanation} \end{cases}$$

(d) $\int_0^1 \sqrt{(x'(t))^2 + (y'(t))^2} \, dt = 757.455862$

The total distance traveled by the boat over the time interval $0 \leq t \leq 1$ is 757.456 (or 757.455) meters.

$$2 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{total distance} \end{cases}$$

2018 SCORING GUIDELINES

Question 3

$$\begin{aligned} \text{(a)} \quad f(-5) &= f(1) + \int_1^{-5} g(x) \, dx = f(1) - \int_{-5}^1 g(x) \, dx \\ &= 3 - \left(-9 - \frac{3}{2} + 1 \right) = 3 - \left(-\frac{19}{2} \right) = \frac{25}{2} \end{aligned}$$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b)} \quad \int_1^6 g(x) \, dx &= \int_1^3 g(x) \, dx + \int_3^6 g(x) \, dx \\ &= \int_1^3 2 \, dx + \int_3^6 2(x-4)^2 \, dx \\ &= 4 + \left[\frac{2}{3}(x-4)^3 \right]_{x=3}^{x=6} = 4 + \frac{16}{3} - \left(-\frac{2}{3} \right) = 10 \end{aligned}$$

3 : $\begin{cases} 1 : \text{split at } x = 3 \\ 1 : \text{antiderivative of } 2(x-4)^2 \\ 1 : \text{answer} \end{cases}$

(c) The graph of f is increasing and concave up on $0 < x < 1$ and $4 < x < 6$ because $f'(x) = g(x) > 0$ and $f''(x) = g'(x)$ is increasing on those intervals.

2 : $\begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$

(d) The graph of f has a point of inflection at $x = 4$ because $f'(x) = g(x)$ changes from decreasing to increasing at $x = 4$.

2 : $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

2018 SCORING GUIDELINES

Question 4

$$(a) \quad H'(6) \approx \frac{H(7) - H(5)}{7 - 5} = \frac{11 - 6}{2} = \frac{5}{2}$$

$H'(6)$ is the rate at which the height of the tree is changing, in meters per year, at time $t = 6$ years.

$$(b) \quad \frac{H(5) - H(3)}{5 - 3} = \frac{6 - 2}{2} = 2$$

Because H is differentiable on $3 \leq t \leq 5$, H is continuous on $3 \leq t \leq 5$.

By the Mean Value Theorem, there exists a value c , $3 < c < 5$, such that $H'(c) = 2$.

$$(c) \quad \text{The average height of the tree over the time interval } 2 \leq t \leq 10 \text{ is given by } \frac{1}{10 - 2} \int_2^{10} H(t) \, dt.$$

$$\begin{aligned} \frac{1}{8} \int_2^{10} H(t) \, dt &\approx \frac{1}{8} \left(\frac{1.5 + 2}{2} \cdot 1 + \frac{2 + 6}{2} \cdot 2 + \frac{6 + 11}{2} \cdot 2 + \frac{11 + 15}{2} \cdot 3 \right) \\ &= \frac{1}{8} (65.75) = \frac{263}{32} \end{aligned}$$

The average height of the tree over the time interval $2 \leq t \leq 10$ is $\frac{263}{32}$ meters.

$$(d) \quad G(x) = 50 \Rightarrow x = 1$$

$$\frac{d}{dt}(G(x)) = \frac{d}{dx}(G(x)) \cdot \frac{dx}{dt} = \frac{(1+x)100 - 100x \cdot 1}{(1+x)^2} \cdot \frac{dx}{dt} = \frac{100}{(1+x)^2} \cdot \frac{dx}{dt}$$

$$\left. \frac{d}{dt}(G(x)) \right|_{x=1} = \frac{100}{(1+1)^2} \cdot 0.03 = \frac{3}{4}$$

According to the model, the rate of change of the height of the tree with respect to time when the tree is 50 meters tall is $\frac{3}{4}$ meter per year.

$$2 : \begin{cases} 1 : \text{estimate} \\ 1 : \text{interpretation with units} \end{cases}$$

$$2 : \begin{cases} 1 : \frac{H(5) - H(3)}{5 - 3} \\ 1 : \text{conclusion using Mean Value Theorem} \end{cases}$$

$$2 : \begin{cases} 1 : \text{trapezoidal sum} \\ 1 : \text{approximation} \end{cases}$$

$$3 : \begin{cases} 2 : \frac{d}{dt}(G(x)) \\ 1 : \text{answer} \end{cases}$$

Note: max 1/3 [1-0] if no chain rule

2018 SCORING GUIDELINES

Question 5

$$(a) \text{ Area} = \frac{1}{2} \int_{\pi/3}^{5\pi/3} (4^2 - (3 + 2\cos \theta)^2) d\theta$$

$$(b) \frac{dr}{d\theta} = -2\sin \theta \Rightarrow \left. \frac{dr}{d\theta} \right|_{\theta=\pi/2} = -2$$

$$r\left(\frac{\pi}{2}\right) = 3 + 2\cos\left(\frac{\pi}{2}\right) = 3$$

$$y = r \sin \theta \Rightarrow \frac{dy}{d\theta} = \frac{dr}{d\theta} \sin \theta + r \cos \theta$$

$$x = r \cos \theta \Rightarrow \frac{dx}{d\theta} = \frac{dr}{d\theta} \cos \theta - r \sin \theta$$

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/2} = \left. \frac{dy/d\theta}{dx/d\theta} \right|_{\theta=\pi/2} = \frac{-2\sin\left(\frac{\pi}{2}\right) + 3\cos\left(\frac{\pi}{2}\right)}{-2\cos\left(\frac{\pi}{2}\right) - 3\sin\left(\frac{\pi}{2}\right)} = \frac{2}{3}$$

The slope of the line tangent to the graph of $r = 3 + 2\cos \theta$

at $\theta = \frac{\pi}{2}$ is $\frac{2}{3}$.

— OR —

$$y = r \sin \theta = (3 + 2\cos \theta) \sin \theta \Rightarrow \frac{dy}{d\theta} = 3\cos \theta + 2\cos^2 \theta - 2\sin^2 \theta$$

$$x = r \cos \theta = (3 + 2\cos \theta) \cos \theta \Rightarrow \frac{dx}{d\theta} = -3\sin \theta - 4\sin \theta \cos \theta$$

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/2} = \left. \frac{dy/d\theta}{dx/d\theta} \right|_{\theta=\pi/2} = \frac{3\cos\left(\frac{\pi}{2}\right) + 2\cos^2\left(\frac{\pi}{2}\right) - 2\sin^2\left(\frac{\pi}{2}\right)}{-3\sin\left(\frac{\pi}{2}\right) - 4\sin\left(\frac{\pi}{2}\right)\cos\left(\frac{\pi}{2}\right)} = \frac{2}{3}$$

The slope of the line tangent to the graph of $r = 3 + 2\cos \theta$

at $\theta = \frac{\pi}{2}$ is $\frac{2}{3}$.

$$(c) \frac{dr}{dt} = \frac{dr}{d\theta} \cdot \frac{d\theta}{dt} = -2\sin \theta \cdot \frac{d\theta}{dt} \Rightarrow \frac{d\theta}{dt} = \frac{dr}{dt} \cdot \frac{1}{-2\sin \theta}$$

$$\left. \frac{d\theta}{dt} \right|_{\theta=\pi/3} = 3 \cdot \frac{1}{-2\sin\left(\frac{\pi}{3}\right)} = \frac{3}{-\sqrt{3}} = -\sqrt{3} \text{ radians per second}$$

$$3 : \begin{cases} 1 : \text{constant and limits} \\ 2 : \text{integrand} \end{cases}$$

$$3 : \begin{cases} 1 : \frac{dy}{d\theta} = \frac{dr}{d\theta} \sin \theta + r \cos \theta \\ \text{or } \frac{dx}{d\theta} = \frac{dr}{d\theta} \cos \theta - r \sin \theta \\ 1 : \frac{dy}{dx} = \frac{\frac{dr}{d\theta} \sin \theta + r \cos \theta}{\frac{dr}{d\theta} \cos \theta - r \sin \theta} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : \frac{dr}{dt} = \frac{dr}{d\theta} \cdot \frac{d\theta}{dt} \\ 1 : \frac{d\theta}{dt} = \frac{dr}{dt} \cdot \frac{1}{-2\sin \theta} \\ 1 : \text{answer with units} \end{cases}$$

2018 SCORING GUIDELINES

Question 6

(a) The first four nonzero terms are $\frac{x^2}{3} - \frac{x^3}{2 \cdot 3^2} + \frac{x^4}{3 \cdot 3^3} - \frac{x^5}{4 \cdot 3^4}$.

The general term is $(-1)^{n+1} \frac{x^{n+1}}{n \cdot 3^n}$.

(b)
$$\lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+2} x^{n+2}}{(n+1)(3^{n+1})}}{\frac{(-1)^{n+1} x^{n+1}}{n \cdot 3^n}} \right| = \lim_{n \rightarrow \infty} \left| \frac{-x}{3} \cdot \frac{n}{n+1} \right| = \left| \frac{x}{3} \right|$$

$\left| \frac{x}{3} \right| < 1$ for $|x| < 3$

Therefore, the radius of convergence of the Maclaurin series for f is 3.

— OR —

The radius of convergence of the Maclaurin series for $\ln(1+x)$ is 1, so the series for $f(x) = x \ln\left(1 + \frac{x}{3}\right)$ converges absolutely for $\left|\frac{x}{3}\right| < 1$.

$\left|\frac{x}{3}\right| < 1 \Rightarrow |x| < 3$

Therefore, the radius of convergence of the Maclaurin series for f is 3.

When $x = -3$, the series is $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{(-3)^{n+1}}{n \cdot 3^n} = \sum_{n=1}^{\infty} \frac{3}{n}$, which diverges by comparison to the harmonic series.

When $x = 3$, the series is $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{3^{n+1}}{n \cdot 3^n} = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{3}{n}$, which converges by the alternating series test.

The interval of convergence of the Maclaurin series for f is $-3 < x \leq 3$.

(c) By the alternating series error bound, an upper bound for $|P_4(2) - f(2)|$ is the magnitude of the next term of the alternating series.

$$|P_4(2) - f(2)| < \left| -\frac{2^5}{4 \cdot 3^4} \right| = \frac{8}{81}$$

2 : $\begin{cases} 1 : \text{first four terms} \\ 1 : \text{general term} \end{cases}$

5 : $\begin{cases} 1 : \text{sets up ratio} \\ 1 : \text{computes limit of ratio} \\ 1 : \text{radius of convergence} \\ 1 : \text{considers both endpoints} \\ 1 : \text{analysis and interval of convergence} \end{cases}$

— OR —

5 : $\begin{cases} 1 : \text{radius for } \ln(1+x) \text{ series} \\ 1 : \text{substitutes } \frac{x}{3} \\ 1 : \text{radius of convergence} \\ 1 : \text{considers both endpoints} \\ 1 : \text{analysis and interval of convergence} \end{cases}$

2 : $\begin{cases} 1 : \text{uses fifth-degree term as error bound} \\ 1 : \text{answer} \end{cases}$

2017**AP[®]** **CollegeBoard**

AP Calculus BC

Scoring Guidelines

2017 SCORING GUIDELINES

Question 1

(a) Volume = $\int_0^{10} A(h) \, dh$ $\approx (2 - 0) \cdot A(0) + (5 - 2) \cdot A(2) + (10 - 5) \cdot A(5)$ $= 2 \cdot 50.3 + 3 \cdot 14.4 + 5 \cdot 6.5$ $= 176.3$ cubic feet (b) The approximation in part (a) is an overestimate because a left Riemann sum is used and A is decreasing. (c) $\int_0^{10} f(h) \, dh = 101.325338$ The volume is 101.325 cubic feet. (d) Using the model, $V(h) = \int_0^h f(x) \, dx$. $\begin{aligned} \left. \frac{dV}{dt} \right _{h=5} &= \left[\frac{dV}{dh} \cdot \frac{dh}{dt} \right]_{h=5} \\ &= \left[f(h) \cdot \frac{dh}{dt} \right]_{h=5} \\ &= f(5) \cdot 0.26 = 1.694419 \end{aligned}$ When $h = 5$, the volume of water is changing at a rate of 1.694 cubic feet per minute.	1 : units in parts (a), (c), and (d) 2 : $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{approximation} \end{cases}$ 1 : overestimate with reason 2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$ 3 : $\begin{cases} 2 : \frac{dV}{dt} \\ 1 : \text{answer} \end{cases}$
--	---

2017 SCORING GUIDELINES

Question 2

$$(a) \frac{1}{2} \int_0^{\pi/2} (f(\theta))^2 d\theta = 0.648414$$

The area of R is 0.648.

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

$$(b) \int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \frac{1}{2} \int_0^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

— OR —

$$\int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \int_k^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

$$2 : \begin{cases} 1 : \text{integral expression} \\ \quad \text{for one region} \\ 1 : \text{equation} \end{cases}$$

$$(c) w(\theta) = g(\theta) - f(\theta)$$

$$w_A = \frac{\int_0^{\pi/2} w(\theta) d\theta}{\frac{\pi}{2} - 0} = 0.485446$$

The average value of $w(\theta)$ on the interval $\left[0, \frac{\pi}{2}\right]$ is 0.485.

$$3 : \begin{cases} 1 : w(\theta) \\ 1 : \text{integral} \\ 1 : \text{average value} \end{cases}$$

$$(d) w(\theta) = w_A \text{ for } 0 \leq \theta \leq \frac{\pi}{2} \Rightarrow \theta = 0.517688$$

$$w(\theta) = w_A \text{ at } \theta = 0.518 \text{ (or 0.517).}$$

$$w'(0.518) < 0 \Rightarrow w(\theta) \text{ is decreasing at } \theta = 0.518.$$

$$2 : \begin{cases} 1 : \text{solves } w(\theta) = w_A \\ 1 : \text{answer with reason} \end{cases}$$

2017 SCORING GUIDELINES

Question 3

$$(a) \quad f(-6) = f(-2) + \int_{-2}^{-6} f'(x) \, dx = 7 - \int_{-6}^{-2} f'(x) \, dx = 7 - 4 = 3$$

$$f(5) = f(-2) + \int_{-2}^5 f'(x) \, dx = 7 - 2\pi + 3 = 10 - 2\pi$$

$$3 : \begin{cases} 1 : \text{uses initial condition} \\ 1 : f(-6) \\ 1 : f(5) \end{cases}$$

$$(b) \quad f'(x) > 0 \text{ on the intervals } [-6, -2) \text{ and } (2, 5).$$

Therefore, f is increasing on the intervals $[-6, -2]$ and $[2, 5]$.

2 : answer with justification

$$(c) \quad \text{The absolute minimum will occur at a critical point where } f'(x) = 0 \text{ or at an endpoint.}$$

$$f'(x) = 0 \Rightarrow x = -2, x = 2$$

$$2 : \begin{cases} 1 : \text{considers } x = 2 \\ 1 : \text{answer with justification} \end{cases}$$

x	$f(x)$
-6	3
-2	7
2	$7 - 2\pi$
5	$10 - 2\pi$

The absolute minimum value is $f(2) = 7 - 2\pi$.

$$(d) \quad f''(-5) = \frac{2 - 0}{-6 - (-2)} = -\frac{1}{2}$$

$$2 : \begin{cases} 1 : f''(-5) \\ 1 : f''(3) \text{ does not exist,} \\ \text{with explanation} \end{cases}$$

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} = 2 \text{ and } \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3} = -1$$

$f''(3)$ does not exist because

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} \neq \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3}.$$

2017 SCORING GUIDELINES

Question 4

$$(a) \quad H'(0) = -\frac{1}{4}(91 - 27) = -16$$

$$H(0) = 91$$

An equation for the tangent line is $y = 91 - 16t$.

The internal temperature of the potato at time $t = 3$ minutes is approximately $91 - 16 \cdot 3 = 43$ degrees Celsius.

$$(b) \quad \frac{d^2H}{dt^2} = -\frac{1}{4} \frac{dH}{dt} = \left(-\frac{1}{4}\right)\left(-\frac{1}{4}\right)(H - 27) = \frac{1}{16}(H - 27)$$

$$H > 27 \text{ for } t > 0 \Rightarrow \frac{d^2H}{dt^2} = \frac{1}{16}(H - 27) > 0 \text{ for } t > 0$$

Therefore, the graph of H is concave up for $t > 0$. Thus, the answer in part (a) is an underestimate.

$$(c) \quad \frac{dG}{(G - 27)^{2/3}} = -dt$$

$$\int \frac{dG}{(G - 27)^{2/3}} = \int (-1) dt$$

$$3(G - 27)^{1/3} = -t + C$$

$$3(91 - 27)^{1/3} = 0 + C \Rightarrow C = 12$$

$$3(G - 27)^{1/3} = 12 - t$$

$$G(t) = 27 + \left(\frac{12 - t}{3}\right)^3 \text{ for } 0 \leq t < 10$$

The internal temperature of the potato at time $t = 3$ minutes is

$$27 + \left(\frac{12 - 3}{3}\right)^3 = 54 \text{ degrees Celsius.}$$

$$3 : \begin{cases} 1 : \text{slope} \\ 1 : \text{tangent line} \\ 1 : \text{approximation} \end{cases}$$

1 : underestimate with reason

$$5 : \begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration and} \\ \quad \text{uses initial condition} \\ 1 : \text{equation involving } G \text{ and } t \\ 1 : G(t) \text{ and } G(3) \end{cases}$$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

2017 SCORING GUIDELINES

Question 5

$$(a) f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2}$$

$$f'(3) = \frac{(-3)(5)}{(18 - 21 + 5)^2} = -\frac{15}{4}$$

$$(b) f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2} = 0 \Rightarrow x = \frac{7}{4}$$

The only critical point in the interval $1 < x < 2.5$ has x -coordinate $\frac{7}{4}$.

f' changes sign from positive to negative at $x = \frac{7}{4}$.

Therefore, f has a relative maximum at $x = \frac{7}{4}$.

$$\begin{aligned} (c) \int_5^\infty f(x) dx &= \lim_{b \rightarrow \infty} \int_5^b \frac{3}{2x^2 - 7x + 5} dx = \lim_{b \rightarrow \infty} \int_5^b \left(\frac{2}{2x - 5} - \frac{1}{x - 1} \right) dx \\ &= \lim_{b \rightarrow \infty} \left[\ln(2x - 5) - \ln(x - 1) \right]_5^b = \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2x - 5}{x - 1} \right) \right]_5^b \\ &= \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2b - 5}{b - 1} \right) - \ln \left(\frac{5}{4} \right) \right] = \ln 2 - \ln \left(\frac{5}{4} \right) = \ln \left(\frac{8}{5} \right) \end{aligned}$$

(d) f is continuous, positive, and decreasing on $[5, \infty)$.

The series converges by the integral test since $\int_5^\infty \frac{3}{2x^2 - 7x + 5} dx$ converges.

— OR —

$$\frac{3}{2n^2 - 7n + 5} > 0 \text{ and } \frac{1}{n^2} > 0 \text{ for } n \geq 5.$$

Since $\lim_{n \rightarrow \infty} \frac{\frac{3}{2n^2 - 7n + 5}}{\frac{1}{n^2}} = \frac{3}{2}$ and the series $\sum_{n=5}^\infty \frac{1}{n^2}$ converges,

the series $\sum_{n=5}^\infty \frac{3}{2n^2 - 7n + 5}$ converges by the limit comparison test.

2 : $f'(3)$

2 : $\begin{cases} 1 : x\text{-coordinate} \\ 1 : \text{relative maximum} \\ \text{with justification} \end{cases}$

3 : $\begin{cases} 1 : \text{antiderivative} \\ 1 : \text{limit expression} \\ 1 : \text{answer} \end{cases}$

2 : answer with conditions

2017 SCORING GUIDELINES

Question 6

(a) $f(0) = 0$

$f'(0) = 1$

$f''(0) = -1(1) = -1$

$f'''(0) = -2(-1) = 2$

$f^{(4)}(0) = -3(2) = -6$

The first four nonzero terms are

$$0 + 1x + \frac{-1}{2!}x^2 + \frac{2}{3!}x^3 + \frac{-6}{4!}x^4 = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4}.$$

The general term is $\frac{(-1)^{n+1}x^n}{n}$.

(b) For $x = 1$, the Maclaurin series becomes $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n}$.

The series does not converge absolutely because the harmonic series diverges.

The series alternates with terms that decrease in magnitude to 0, and therefore the series converges conditionally.

$$\begin{aligned}
 \text{(c)} \quad \int_0^x f(t) dt &= \int_0^x \left(t - \frac{t^2}{2} + \frac{t^3}{3} - \frac{t^4}{4} + \cdots + \frac{(-1)^{n+1}t^n}{n} + \cdots \right) dt \\
 &= \left[\frac{t^2}{2} - \frac{t^3}{3 \cdot 2} + \frac{t^4}{4 \cdot 3} - \frac{t^5}{5 \cdot 4} + \cdots + \frac{(-1)^{n+1}t^{n+1}}{(n+1)n} + \cdots \right]_{t=0}^{t=x} \\
 &= \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{12} - \frac{x^5}{20} + \cdots + \frac{(-1)^{n+1}x^{n+1}}{(n+1)n} + \cdots
 \end{aligned}$$

(d) The terms alternate in sign and decrease in magnitude to 0. By the alternating series error bound, the error $\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right|$ is bounded

by the magnitude of the first unused term, $\left| -\frac{(1/2)^5}{20} \right|$.

Thus, $\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right| \leq \left| -\frac{(1/2)^5}{20} \right| = \frac{1}{32 \cdot 20} < \frac{1}{500}.$

$$3 : \begin{cases} 1 : f''(0), f'''(0), \text{ and } f^{(4)}(0) \\ 1 : \text{verify terms} \\ 1 : \text{general term} \end{cases}$$

2 : converges conditionally
with reason

$$3 : \begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \\ 1 : \text{general term} \end{cases}$$

1 : error bound