

Part B (AB or BC): Graphing calculator not allowed**Question 3****9 points****General Scoring Notes**


The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

The depth of seawater at a location can be modeled by the function H that satisfies the differential equation $\frac{dH}{dt} = \frac{1}{2}(H - 1)\cos\left(\frac{t}{2}\right)$, where $H(t)$ is measured in feet and t is measured in hours after noon ($t = 0$). It is known that $H(0) = 4$.

Model Solution**Scoring**

- (a) A portion of the slope field for the differential equation is provided. Sketch the solution curve, $y = H(t)$, through the point $(0, 4)$.



Solution curve

1 point**Scoring notes:**

- The solution curve must pass through the point $(0, 4)$, extend to at least $t = 4.5$, and have no obvious conflicts with the given slope lines.
- Only portions of the solution curve within the given slope field are considered.

Total for part (a) 1 point

- (b) For $0 < t < 5$, it can be shown that $H(t) > 1$. Find the value of t , for $0 < t < 5$, at which H has a critical point. Determine whether the critical point corresponds to a relative minimum, a relative maximum, or neither a relative minimum nor a relative maximum of the depth of seawater at the location. Justify your answer.

Because $H(t) > 1$, then $\frac{dH}{dt} = 0$ implies $\cos\left(\frac{t}{2}\right) = 0$.	Considers sign of $\frac{dH}{dt}$	1 point
This implies that $t = \pi$ is a critical point.	Identifies $t = \pi$	1 point
For $0 < t < \pi$, $\frac{dH}{dt} > 0$ and for $\pi < t < 5$, $\frac{dH}{dt} < 0$. Therefore, $t = \pi$ is the location of a relative maximum value of H .	Answer with justification	1 point

Scoring notes:

- The first point is earned for considering $\frac{dH}{dt} = 0$, $\frac{dH}{dt} > 0$, $\frac{dH}{dt} < 0$, $\cos\left(\frac{t}{2}\right) = 0$, $\cos\left(\frac{t}{2}\right) > 0$, or $\cos\left(\frac{t}{2}\right) < 0$.
- The second point is earned for identifying $t = \pi$, with or without supporting work. A response may consider $H = 1$ or $t = 1$ as potential critical points without penalty.
- The third point cannot be earned without the first point. The third point is earned only for a correct justification and a correct answer of “relative maximum.”
- The justification can be shown by determining the sign of $\frac{dH}{dt}$ (or $\cos\left(\frac{t}{2}\right)$) at a single value in $0 < t < \pi$ and at a single value in $\pi < t < 5$. It is not necessary to state that $\frac{dH}{dt}$ does not change sign on these intervals.
- The third point can also be earned by using the Second Derivative Test. For example:

$$\frac{d^2H}{dt^2} = \frac{1}{2}(H - 1)\left(-\frac{1}{2}\sin\left(\frac{t}{2}\right)\right) + \cos\left(\frac{t}{2}\right) \cdot \frac{1}{2} \cdot \frac{dH}{dt}$$

$$\left.\frac{d^2H}{dt^2}\right|_{t=\pi} < 0$$

Therefore, $t = \pi$ is the location of a relative maximum value of H .

Total for part (b) 3 points

- (c) Use separation of variables to find $y = H(t)$, the particular solution to the differential equation

$$\frac{dH}{dt} = \frac{1}{2}(H - 1)\cos\left(\frac{t}{2}\right) \text{ with initial condition } H(0) = 4.$$

$\frac{dH}{H-1} = \frac{1}{2}\cos\left(\frac{t}{2}\right)dt$	Separation of variables	1 point
$\int \frac{dH}{H-1} = \int \frac{1}{2}\cos\left(\frac{t}{2}\right)dt$ $\Rightarrow \ln H-1 = \sin\left(\frac{t}{2}\right) + C$	One antiderivative	1 point
	Second antiderivative	1 point
$\ln 4-1 = \sin\left(\frac{0}{2}\right) + C \Rightarrow C = \ln 3$ Because $H(0) = 4$, $H > 1$, so $ H-1 = H-1$. $\ln(H-1) = \sin\left(\frac{t}{2}\right) + \ln 3$	Constant of integration and uses initial condition	1 point
$H-1 = e^{\sin(t/2)+\ln 3} = 3e^{\sin(t/2)}$ $H(t) = 1 + 3e^{\sin(t/2)}$	Solves for H	1 point

Scoring notes:

- A response with no separation of variables earns 0 out of 5 points.
- A response that presents $\int \frac{dH}{H-1} = \ln(H-1)$ without absolute value symbols earns that antiderivative point.
- A response with no constant of integration can earn at most the first 3 points.
- A response is eligible for the fourth point only if it has earned the first point and at least 1 of the 2 antiderivative points.
- An eligible response earns the fourth point by correctly including the constant of integration in an equation and substituting 0 for t and 4 for H .
- A response is eligible for the fifth point only if it has earned the first 4 points.
- A response earns the fifth point only for an answer of $H(t) = 1 + 3e^{\sin(t/2)}$ or a mathematically equivalent expression for $H(t)$ such as $H(t) = 1 + e^{\sin(t/2)+\ln 3}$.
- A response does not need to argue that $|H-1| = H-1$ in order to earn the fifth point.
- Special case: A response that presents an incorrect separation of variables of $\frac{1}{2} \cdot \frac{dH}{H-1} = \cos\left(\frac{t}{2}\right)dt$ does not earn the first point or the fifth point but is eligible for the 2 antiderivative points. If the response earns at least 1 of the 2 antiderivative points, then the response is eligible for the fourth point.

Total for part (c) 5 points

Total for question 3 9 points