

Part B (AB or BC): Graphing calculator not allowed**Question 3****9 points****General Scoring Notes**


The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

The depth of seawater at a location can be modeled by the function H that satisfies the differential equation $\frac{dH}{dt} = \frac{1}{2}(H - 1)\cos\left(\frac{t}{2}\right)$, where $H(t)$ is measured in feet and t is measured in hours after noon ($t = 0$). It is known that $H(0) = 4$.

Model Solution**Scoring**

- (a) A portion of the slope field for the differential equation is provided. Sketch the solution curve, $y = H(t)$, through the point $(0, 4)$.



Solution curve

1 point**Scoring notes:**

- The solution curve must pass through the point $(0, 4)$, extend to at least $t = 4.5$, and have no obvious conflicts with the given slope lines.
- Only portions of the solution curve within the given slope field are considered.

Total for part (a) 1 point

- (b) For $0 < t < 5$, it can be shown that $H(t) > 1$. Find the value of t , for $0 < t < 5$, at which H has a critical point. Determine whether the critical point corresponds to a relative minimum, a relative maximum, or neither a relative minimum nor a relative maximum of the depth of seawater at the location. Justify your answer.

Because $H(t) > 1$, then $\frac{dH}{dt} = 0$ implies $\cos\left(\frac{t}{2}\right) = 0$.	Considers sign of $\frac{dH}{dt}$	1 point
This implies that $t = \pi$ is a critical point.	Identifies $t = \pi$	1 point
For $0 < t < \pi$, $\frac{dH}{dt} > 0$ and for $\pi < t < 5$, $\frac{dH}{dt} < 0$. Therefore, $t = \pi$ is the location of a relative maximum value of H .	Answer with justification	1 point

Scoring notes:

- The first point is earned for considering $\frac{dH}{dt} = 0$, $\frac{dH}{dt} > 0$, $\frac{dH}{dt} < 0$, $\cos\left(\frac{t}{2}\right) = 0$, $\cos\left(\frac{t}{2}\right) > 0$, or $\cos\left(\frac{t}{2}\right) < 0$.
- The second point is earned for identifying $t = \pi$, with or without supporting work. A response may consider $H = 1$ or $t = 1$ as potential critical points without penalty.
- The third point cannot be earned without the first point. The third point is earned only for a correct justification and a correct answer of “relative maximum.”
- The justification can be shown by determining the sign of $\frac{dH}{dt}$ (or $\cos\left(\frac{t}{2}\right)$) at a single value in $0 < t < \pi$ and at a single value in $\pi < t < 5$. It is not necessary to state that $\frac{dH}{dt}$ does not change sign on these intervals.
- The third point can also be earned by using the Second Derivative Test. For example:

$$\frac{d^2H}{dt^2} = \frac{1}{2}(H - 1)\left(-\frac{1}{2}\sin\left(\frac{t}{2}\right)\right) + \cos\left(\frac{t}{2}\right) \cdot \frac{1}{2} \cdot \frac{dH}{dt}$$

$$\left.\frac{d^2H}{dt^2}\right|_{t=\pi} < 0$$

Therefore, $t = \pi$ is the location of a relative maximum value of H .

Total for part (b) 3 points

- (c) Use separation of variables to find $y = H(t)$, the particular solution to the differential equation

$$\frac{dH}{dt} = \frac{1}{2}(H - 1)\cos\left(\frac{t}{2}\right) \text{ with initial condition } H(0) = 4.$$

$\frac{dH}{H-1} = \frac{1}{2}\cos\left(\frac{t}{2}\right)dt$	Separation of variables	1 point
$\int \frac{dH}{H-1} = \int \frac{1}{2}\cos\left(\frac{t}{2}\right)dt$ $\Rightarrow \ln H-1 = \sin\left(\frac{t}{2}\right) + C$	One antiderivative	1 point
	Second antiderivative	1 point
$\ln 4-1 = \sin\left(\frac{0}{2}\right) + C \Rightarrow C = \ln 3$ Because $H(0) = 4$, $H > 1$, so $ H-1 = H-1$. $\ln(H-1) = \sin\left(\frac{t}{2}\right) + \ln 3$	Constant of integration and uses initial condition	1 point
$H-1 = e^{\sin(t/2)+\ln 3} = 3e^{\sin(t/2)}$ $H(t) = 1 + 3e^{\sin(t/2)}$	Solves for H	1 point

Scoring notes:

- A response with no separation of variables earns 0 out of 5 points.
- A response that presents $\int \frac{dH}{H-1} = \ln(H-1)$ without absolute value symbols earns that antiderivative point.
- A response with no constant of integration can earn at most the first 3 points.
- A response is eligible for the fourth point only if it has earned the first point and at least 1 of the 2 antiderivative points.
- An eligible response earns the fourth point by correctly including the constant of integration in an equation and substituting 0 for t and 4 for H .
- A response is eligible for the fifth point only if it has earned the first 4 points.
- A response earns the fifth point only for an answer of $H(t) = 1 + 3e^{\sin(t/2)}$ or a mathematically equivalent expression for $H(t)$ such as $H(t) = 1 + e^{\sin(t/2)+\ln 3}$.
- A response does not need to argue that $|H-1| = H-1$ in order to earn the fifth point.
- Special case: A response that presents an incorrect separation of variables of $\frac{1}{2} \cdot \frac{dH}{H-1} = \cos\left(\frac{t}{2}\right)dt$ does not earn the first point or the fifth point but is eligible for the 2 antiderivative points. If the response earns at least 1 of the 2 antiderivative points, then the response is eligible for the fourth point.

Total for part (c) 5 points


Total for question 3 9 points

Part B (AB or BC): Graphing calculator not allowed**Question 3****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

A bottle of milk is taken out of a refrigerator and placed in a pan of hot water to be warmed. The increasing function M models the temperature of the milk at time t , where $M(t)$ is measured in degrees Celsius ($^{\circ}\text{C}$) and t is the number of minutes since the bottle was placed in the pan. M satisfies the differential equation $\frac{dM}{dt} = \frac{1}{4}(40 - M)$. At time $t = 0$, the temperature of the milk is 5°C . It can be shown that $M(t) < 40$ for all values of t .

Model Solution	Scoring
(a) A slope field for the differential equation $\frac{dM}{dt} = \frac{1}{4}(40 - M)$ is shown. Sketch the solution curve through the point $(0, 5)$. 	Solution curve 1 point

Scoring notes:

- The solution curve must pass through the point $(0, 5)$, extend reasonably close to the left and right edges of the rectangle, and have no obvious conflicts with the given slope lines.
- Only portions of the solution curve within the given slope field are considered.
- The solution curve must lie entirely below the horizontal line segments at $M = 40$.

Total Score (a) 1 point

- (b) Use the line tangent to the graph of M at $t = 0$ to approximate $M(2)$, the temperature of the milk at time $t = 2$ minutes.

$\left. \frac{dM}{dt} \right _{t=0} = \frac{1}{4}(40 - 5) = \frac{35}{4}$	$\left. \frac{dM}{dt} \right _{t=0}$ 1 point
The tangent line equation is $y = 5 + \frac{35}{4}(t - 0)$. $M(2) \approx 5 + \frac{35}{4} \cdot 2 = 22.5$ The temperature of the milk at time $t = 2$ minutes is approximately 22.5° Celsius.	Approximation 1 point

Scoring notes:

- The value of the slope may appear in a tangent line equation or approximation.
- A response of $5 + \frac{35}{4} \cdot 2$ is the minimal response to earn both points.
- A response of $\frac{1}{4}(40 - 5)$ earns the first point. If there are any subsequent errors in simplification, the response does not earn the second point.
- In order to earn the second point the response must present an approximation found by using a tangent line that:
 - passes through the point $(0, 5)$ and
 - has slope $\frac{35}{4}$ or a nonzero slope that is declared to be the value of $\frac{dM}{dt}$.
- An unsupported approximation does not earn the second point.
- The approximation need not be simplified, but the response does not earn the second point if the approximation is simplified incorrectly.

Total for part (b) 2 points

- (c) Write an expression for $\frac{d^2M}{dt^2}$ in terms of M . Use $\frac{d^2M}{dt^2}$ to determine whether the approximation from part (b) is an underestimate or an overestimate for the actual value of $M(2)$. Give a reason for your answer.

$\frac{d^2M}{dt^2} = -\frac{1}{4} \frac{dM}{dt} = -\frac{1}{4} \left(\frac{1}{4}(40 - M) \right) = -\frac{1}{16}(40 - M)$	$\frac{d^2M}{dt^2}$ 1 point
Because $M(t) < 40$, $\frac{d^2M}{dt^2} < 0$, so the graph of M is concave down. Therefore, the tangent line approximation of $M(2)$ is an overestimate.	Overestimate with reason 1 point

Scoring notes:

- The first point is earned for either $\frac{d^2M}{dt^2} = -\frac{1}{4}\left(\frac{1}{4}(40 - M)\right)$ or $\frac{d^2M}{dt^2} = -\frac{1}{16}(40 - M)$ (or equivalent). A response that presents any subsequent simplification error does not earn the second point.
- A response that presents an expression for $\frac{d^2M}{dt^2}$ in terms of $\frac{dM}{dt}$ but fails to continue to an expression in terms of M (i.e., $\frac{d^2M}{dt^2} = -\frac{1}{4}\frac{dM}{dt}$) does not earn the first point but is eligible for the second point.
- If the response presents an expression for $\frac{d^2M}{dt^2}$ that is incorrect, the response is eligible for the second point only if the expression is a nonconstant linear function that is negative for $5 < M < 40$.
 - Special case: A response that presents $\frac{d^2M}{dt^2} = \frac{1}{16}(40 - M)$ does not earn the first point but is eligible to earn the second point for a consistent answer and reason.
- To earn the second point a response must include $\frac{d^2M}{dt^2} < 0$, or $\frac{dM}{dt}$ is decreasing, or the graph of M is concave down, as well as the conclusion that the approximation is an overestimate.
- A response that presents an argument based on $\frac{d^2M}{dt^2}$ or concavity at a single point does not earn the second point.

Total for part (c) 2 points

- (d) Use separation of variables to find an expression for $M(t)$, the particular solution to the differential equation $\frac{dM}{dt} = \frac{1}{4}(40 - M)$ with initial condition $M(0) = 5$.

$\frac{dM}{40 - M} = \frac{1}{4} dt$ $\int \frac{dM}{40 - M} = \int \frac{1}{4} dt$	Separates variables	1 point
$-\ln 40 - M = \frac{1}{4}t + C$	Finds antiderivatives	1 point
$-\ln 40 - 5 = 0 + C \Rightarrow C = -\ln 35$ $M(t) < 40 \Rightarrow 40 - M > 0 \Rightarrow 40 - M = 40 - M$ $-\ln(40 - M) = \frac{1}{4}t - \ln 35$ $\ln(40 - M) = -\frac{1}{4}t + \ln 35$	Constant of integration and uses initial condition	1 point

$$40 - M = 35e^{-t/4}$$

$$M = 40 - 35e^{-t/4}$$

Solves for M **1 point****Scoring notes:**

- A response with no separation of variables earns 0 out of 4 points.
- A response that presents an antiderivative of $-\ln(40 - M)$ without absolute value symbols is eligible for all 4 points.
- A response with no constant of integration can earn at most the first 2 points.
- A response is eligible for the third point only if it has earned the first 2 points.
 - Special Case: A response that presents $+\ln(40 - M) = \frac{t}{4} + C$ (or equivalent) does not earn the second point, is eligible for the third point, but not eligible for the fourth.
- An eligible response earns the third point by correctly including the constant of integration in an equation and substituting 0 for t and 5 for M .
- A response is eligible for the fourth point only if it has earned the first 3 points.
- A response earns the fourth point only for an answer of $M = 40 - 35e^{-t/4}$ or equivalent.

Total for part (d) 4 points**Total for question 3 9 points**

Part B (BC): Graphing calculator not allowed**Question 5****9 points****General Scoring Notes**

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Scoring guidelines and notes contain examples of the most common approaches seen in student responses. These guidelines can be applied to alternate approaches to ensure that these alternate approaches are scored appropriately.

Let $y = f(x)$ be the particular solution to the differential equation $\frac{dy}{dx} = y \cdot (x \ln x)$ with initial condition $f(1) = 4$. It can be shown that $f''(1) = 4$.

Model Solution	Scoring
(a) Write the second-degree Taylor polynomial for f about $x = 1$. Use the Taylor polynomial to approximate $f(2)$. $f'(1) = \left. \frac{dy}{dx} \right _{(x,y)=(1,4)} = 4 \cdot (1 \ln 1) = 0$ The second-degree Taylor polynomial for f about $x = 1$ is $f(1) + f'(1)(x-1) + \frac{f''(1)}{2!}(x-1)^2 = 4 + 0(x-1) + \frac{4}{2}(x-1)^2.$	Polynomial 1 point
$f(2) \approx 4 + 2(2-1)^2 = 6$	Approximation 1 point

Scoring notes:

- The first point is earned for $4 + \frac{4 \cdot \ln 1}{1!}(x-1)^1 + \frac{4}{2!}(x-1)^2$ or any correctly simplified equivalent expression. A term involving $(x-1)$ is not necessary. The polynomial must be written about (centered at) $x = 1$.
- If the first point is earned, the second point is earned for just “6” with no additional supporting work.
- If the polynomial is never explicitly written, the first point is not earned. In this case, to earn the second point supporting work of at least “ $4 + 2(1)$ ” is required.

Total for part (a) 2 points

- (b) Use Euler's method, starting at $x = 1$ with two steps of equal size, to approximate $f(2)$. Show the work that leads to your answer.

$f(1.5) \approx f(1) + 0.5 \cdot \left. \frac{dy}{dx} \right _{(x,y)=(1,4)} = 4 + 0.5 \cdot 0 = 4$ $f(2) \approx f(1.5) + 0.5 \cdot \left. \frac{dy}{dx} \right _{(x,y)=(1.5,4)}$	Euler's method with two steps	1 point
$\approx 4 + 0.5 \cdot 4 \cdot (1.5 \ln 1.5) = 4 + 3 \ln 1.5$	Answer	1 point

Scoring notes:

- The first point is earned for two steps (of size 0.5) of Euler's method, with at most one error. If there is any error, the second point is not earned.
- To earn the first point a response must contain two Euler steps, $\Delta x = 0.5$, use of the correct expression for $\frac{dy}{dx}$, and use of the initial condition $f(1) = 4$.
 - The two Euler steps may be explicit expressions or may be presented in a table. Here is a minimal example of a (correctly labeled) table.

x	y	$\Delta y = \frac{dy}{dx} \cdot \Delta x$ or $\Delta y = \frac{dy}{dx} \cdot (0.5)$
1	4	0
1.5	4	$3 \ln 1.5$
2	$4 + 3 \ln 1.5$	

- Note: In the presence of the correct answer, such a table does not need to be labeled in order to earn both points. In the presence of an incorrect answer, the table must be correctly labeled for the response to earn the first point.
- A single error in computing the approximation of $f(1.5)$ is not considered a second error if that incorrect value is imported correctly into an approximation of $f(2)$.
- Both points are earned for " $4 + 0.5 \cdot 0 + 0.5 \cdot 4 \cdot (1.5 \ln 1.5)$ " or " $4 + 0.5 \cdot 4 \cdot (1.5 \ln 1.5)$ ".
- Both points are earned for presenting the ordered pair $(2, 4 + 3 \ln 1.5)$ with sufficient supporting work.

Total for part (b) 2 points

- (c) Find the particular solution $y = f(x)$ to the differential equation $\frac{dy}{dx} = y \cdot (x \ln x)$ with initial condition $f(1) = 4$.

$\frac{1}{y} dy = x \ln x dx$	Separation of variables	1 point
Using integration by parts, $u = \ln x \quad du = \frac{1}{x} dx$ $dv = x dx \quad v = \frac{x^2}{2}$ $\int x \ln x dx = \frac{x^2}{2} \cdot \ln x - \int \frac{x^2}{2} \cdot \frac{1}{x} dx = \frac{x^2 \ln x}{2} - \frac{x^2}{4} + C$	Antiderivative for $x \ln x$	1 point
$\ln y = \frac{x^2 \ln x}{2} - \frac{x^2}{4} + C$	Antiderivative for $\frac{1}{y}$	1 point
$\ln 4 = 0 - \frac{1}{4} + C \Rightarrow C = \ln 4 + \frac{1}{4}$	Constant of integration and uses initial condition	1 point
$y = e^{\left(\frac{x^2 \ln x}{2} - \frac{x^2}{4} + \ln 4 + \frac{1}{4}\right)}$ Note: This solution is valid for $x > 0$.	Solves for y	1 point

Scoring notes:

- A response with no separation of variables earns 0 out of 5 points. If an error in separation results in one side being correct ($\frac{1}{y} dy$ or $x \ln x dx$), the response is only eligible to earn the corresponding antiderivative point.
- The third point (antiderivative of $\frac{1}{y}$) can be earned for either $\ln y$ or $\ln|y|$.
- A response with no constant of integration can earn at most 3 out of 5 points.
- A response is eligible for the fourth point if it has earned the first point and at least 1 of the 2 antiderivative points.
- A response earns the fourth point by correctly including the constant of integration in an equation and then replacing x with 1 and y with 4.
- A response is eligible for the fifth point only if it has earned the first 4 points.

Total for part (c) 5 points

Total for question 5 9 points

2019**AP[®]** **CollegeBoard**

AP[®] Calculus BC

Scoring Guidelines

2019 SCORING GUIDELINES

Question 1

(a) $\int_0^5 E(t) dt = 153.457690$

To the nearest whole number, 153 fish enter the lake from midnight to 5 A.M.

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

(b) $\frac{1}{5-0} \int_0^5 L(t) dt = 6.059038$

The average number of fish that leave the lake per hour from midnight to 5 A.M. is 6.059 fish per hour.

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

- (c) The rate of change in the number of fish in the lake at time t is given by $E(t) - L(t)$.

$$E(t) - L(t) = 0 \Rightarrow t = 6.20356$$

$E(t) - L(t) > 0$ for $0 \leq t < 6.20356$, and $E(t) - L(t) < 0$ for $6.20356 < t \leq 8$. Therefore the greatest number of fish in the lake is at time $t = 6.204$ (or 6.203).

$$3 : \begin{cases} 1 : \text{sets } E(t) - L(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

— OR —

Let $A(t)$ be the change in the number of fish in the lake from midnight to t hours after midnight.

$$A(t) = \int_0^t (E(s) - L(s)) ds$$

$$A'(t) = E(t) - L(t) = 0 \Rightarrow t = C = 6.20356$$

t	$A(t)$
0	0
C	135.01492
8	80.91998

Therefore the greatest number of fish in the lake is at time $t = 6.204$ (or 6.203).

(d) $E'(5) - L'(5) = -10.7228 < 0$

Because $E'(5) - L'(5) < 0$, the rate of change in the number of fish is decreasing at time $t = 5$.

$$2 : \begin{cases} 1 : \text{considers } E'(5) \text{ and } L'(5) \\ 1 : \text{answer with explanation} \end{cases}$$

2019 SCORING GUIDELINES

Question 2

$$(a) \quad \frac{1}{2} \int_0^{\sqrt{\pi}} (r(\theta))^2 d\theta = 3.534292$$

The area of S is 3.534.

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

$$(b) \quad \frac{1}{\sqrt{\pi} - 0} \int_0^{\sqrt{\pi}} r(\theta) d\theta = 1.579933$$

The average distance from the origin to a point on the curve $r = r(\theta)$ for $0 \leq \theta \leq \sqrt{\pi}$ is 1.580 (or 1.579).

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

$$(c) \quad \tan \theta = \frac{y}{x} = m \Rightarrow \theta = \tan^{-1} m$$

$$\frac{1}{2} \int_0^{\tan^{-1} m} (r(\theta))^2 d\theta = \frac{1}{2} \left(\frac{1}{2} \int_0^{\sqrt{\pi}} (r(\theta))^2 d\theta \right)$$

$$3 : \begin{cases} 1 : \text{equates polar areas} \\ 1 : \text{inverse trigonometric function} \\ \text{applied to } m \text{ as limit of} \\ \text{integration} \\ 1 : \text{equation} \end{cases}$$

(d) As $k \rightarrow \infty$, the circle $r = k \cos \theta$ grows to enclose all points to the right of the y -axis.

$$2 : \begin{cases} 1 : \text{limits of integration} \\ 1 : \text{answer with integral} \end{cases}$$

$$\begin{aligned} \lim_{k \rightarrow \infty} A(k) &= \frac{1}{2} \int_0^{\pi/2} (r(\theta))^2 d\theta \\ &= \frac{1}{2} \int_0^{\pi/2} (3\sqrt{\theta} \sin(\theta^2))^2 d\theta = 3.324 \end{aligned}$$

2019 SCORING GUIDELINES

Question 3

$$\begin{aligned}
 \text{(a)} \quad \int_{-6}^5 f(x) \, dx &= \int_{-6}^{-2} f(x) \, dx + \int_{-2}^5 f(x) \, dx \\
 &\Rightarrow 7 = \int_{-6}^{-2} f(x) \, dx + 2 + \left(9 - \frac{9\pi}{4}\right) \\
 &\Rightarrow \int_{-6}^{-2} f(x) \, dx = 7 - \left(11 - \frac{9\pi}{4}\right) = \frac{9\pi}{4} - 4
 \end{aligned}$$

$$\begin{aligned}
 \text{(b)} \quad \int_3^5 (2f'(x) + 4) \, dx &= 2 \int_3^5 f'(x) \, dx + \int_3^5 4 \, dx \\
 &= 2(f(5) - f(3)) + 4(5 - 3) \\
 &= 2(0 - (3 - \sqrt{5})) + 8 \\
 &= 2(-3 + \sqrt{5}) + 8 = 2 + 2\sqrt{5}
 \end{aligned}$$

— OR —

$$\begin{aligned}
 \int_3^5 (2f'(x) + 4) \, dx &= [2f(x) + 4x]_{x=3}^{x=5} \\
 &= (2f(5) + 20) - (2f(3) + 12) \\
 &= (2 \cdot 0 + 20) - (2(3 - \sqrt{5}) + 12) \\
 &= 2 + 2\sqrt{5}
 \end{aligned}$$

$$\text{(c)} \quad g'(x) = f(x) = 0 \Rightarrow x = -1, x = \frac{1}{2}, x = 5$$

x	$g(x)$
-2	0
-1	$\frac{1}{2}$
$\frac{1}{2}$	$-\frac{1}{4}$
5	$11 - \frac{9\pi}{4}$

On the interval $-2 \leq x \leq 5$, the absolute maximum value of g is $g(5) = 11 - \frac{9\pi}{4}$.

$$\begin{aligned}
 \text{(d)} \quad \lim_{x \rightarrow 1} \frac{10^x - 3f'(x)}{f(x) - \arctan x} &= \frac{10^1 - 3f'(1)}{f(1) - \arctan 1} \\
 &= \frac{10 - 3 \cdot 2}{1 - \arctan 1} = \frac{4}{1 - \frac{\pi}{4}}
 \end{aligned}$$

$$3 : \begin{cases} 1 : \int_{-6}^5 f(x) \, dx = \int_{-6}^{-2} f(x) \, dx + \int_{-2}^5 f(x) \, dx \\ 1 : \int_{-2}^5 f(x) \, dx \\ 1 : \text{answer} \end{cases}$$

$$2 : \begin{cases} 1 : \text{Fundamental Theorem of Calculus} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : g'(x) = f(x) \\ 1 : \text{identifies } x = -1 \text{ as a candidate} \\ 1 : \text{answer with justification} \end{cases}$$

$$1 : \text{answer}$$

2019 SCORING GUIDELINES

Question 4

(a) $V = \pi r^2 h = \pi(1)^2 h = \pi h$
 $\left. \frac{dV}{dt} \right|_{h=4} = \pi \left. \frac{dh}{dt} \right|_{h=4} = \pi \left(-\frac{1}{10} \sqrt{4} \right) = -\frac{\pi}{5}$ cubic feet per second

$$2 : \begin{cases} 1 : \frac{dV}{dt} = \pi \frac{dh}{dt} \\ 1 : \text{answer with units} \end{cases}$$

(b) $\frac{d^2 h}{dt^2} = -\frac{1}{20\sqrt{h}} \cdot \frac{dh}{dt} = -\frac{1}{20\sqrt{h}} \cdot \left(-\frac{1}{10} \sqrt{h} \right) = \frac{1}{200}$
 Because $\frac{d^2 h}{dt^2} = \frac{1}{200} > 0$ for $h > 0$, the rate of change of the height is increasing when the height of the water is 3 feet.

$$3 : \begin{cases} 1 : \frac{d}{dh} \left(-\frac{1}{10} \sqrt{h} \right) = -\frac{1}{20\sqrt{h}} \\ 1 : \frac{d^2 h}{dt^2} = -\frac{1}{20\sqrt{h}} \cdot \frac{dh}{dt} \\ 1 : \text{answer with explanation} \end{cases}$$

(c) $\frac{dh}{\sqrt{h}} = -\frac{1}{10} dt$
 $\int \frac{dh}{\sqrt{h}} = \int -\frac{1}{10} dt$
 $2\sqrt{h} = -\frac{1}{10} t + C$
 $2\sqrt{5} = -\frac{1}{10} \cdot 0 + C \Rightarrow C = 2\sqrt{5}$
 $2\sqrt{h} = -\frac{1}{10} t + 2\sqrt{5}$
 $h(t) = \left(-\frac{1}{20} t + \sqrt{5} \right)^2$

$$4 : \begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ \text{and uses initial condition} \\ 1 : h(t) \end{cases}$$

Note: 0/4 if no separation of variables

Note: max 2/4 [1-1-0-0] if no constant of integration

2019 SCORING GUIDELINES

Question 5

$$(a) \quad f'(x) = \frac{-(2x-2)}{(x^2-2x+k)^2}$$

$$f'(0) = \frac{2}{k^2} = 6 \Rightarrow k^2 = \frac{1}{3} \Rightarrow k = \frac{1}{\sqrt{3}}$$

3 : $\begin{cases} 1 : \text{denominator of } f'(x) \\ 1 : f'(x) \\ 1 : \text{answer} \end{cases}$

$$(b) \quad \frac{1}{x^2-2x-8} = \frac{1}{(x-4)(x+2)} = \frac{A}{x-4} + \frac{B}{x+2}$$

$$\Rightarrow 1 = A(x+2) + B(x-4)$$

$$\Rightarrow A = \frac{1}{6}, B = -\frac{1}{6}$$

3 : $\begin{cases} 1 : \text{partial fraction decomposition} \\ 1 : \text{antiderivatives} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \int_0^1 f(x) dx &= \int_0^1 \left(\frac{\frac{1}{6}}{x-4} - \frac{\frac{1}{6}}{x+2} \right) dx \\ &= \left[\frac{1}{6} \ln|x-4| - \frac{1}{6} \ln|x+2| \right]_{x=0}^{x=1} \\ &= \left(\frac{1}{6} \ln 3 - \frac{1}{6} \ln 3 \right) - \left(\frac{1}{6} \ln 4 - \frac{1}{6} \ln 2 \right) = -\frac{1}{6} \ln 2 \end{aligned}$$

$$\begin{aligned} (c) \quad \int_0^2 \frac{1}{x^2-2x+1} dx &= \int_0^2 \frac{1}{(x-1)^2} dx = \int_0^1 \frac{1}{(x-1)^2} dx + \int_1^2 \frac{1}{(x-1)^2} dx \\ &= \lim_{b \rightarrow 1^-} \int_0^b \frac{1}{(x-1)^2} dx + \lim_{b \rightarrow 1^+} \int_b^2 \frac{1}{(x-1)^2} dx \\ &= \lim_{b \rightarrow 1^-} \left(-\frac{1}{x-1} \Big|_{x=0}^{x=b} \right) + \lim_{b \rightarrow 1^+} \left(-\frac{1}{x-1} \Big|_{x=b}^{x=2} \right) \\ &= \lim_{b \rightarrow 1^-} \left(-\frac{1}{b-1} - 1 \right) + \lim_{b \rightarrow 1^+} \left(-1 + \frac{1}{b-1} \right) \end{aligned}$$

Because $\lim_{b \rightarrow 1^-} \left(-\frac{1}{b-1} \right)$ does not exist, the integral diverges.

3 : $\begin{cases} 1 : \text{improper integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer with reason} \end{cases}$

2019 SCORING GUIDELINES

Question 6

(a) $f(0) = 3$ and $f'(0) = -2$

The third-degree Taylor polynomial for f about $x = 0$ is

$$3 - 2x + \frac{3}{2!}x^2 + \frac{-2}{3!}x^3 = 3 - 2x + \frac{3}{2}x^2 - \frac{2}{6}x^3.$$

(b) The first three nonzero terms of the Maclaurin series for e^x are

$$1 + x + \frac{1}{2!}x^2.$$

The second-degree Taylor polynomial for $e^x f(x)$ about $x = 0$ is

$$\begin{aligned} 3\left(1 + x + \frac{1}{2!}x^2\right) - 2x(1 + x) + \frac{3}{2}x^2(1) \\ = 3 + (3 - 2)x + \left(\frac{3}{2} - 2 + \frac{3}{2}\right)x^2 \\ = 3 + x + x^2. \end{aligned}$$

(c) $h(1) = \int_0^1 f(t) dt$

$$\begin{aligned} &\approx \int_0^1 \left(3 - 2t + \frac{3}{2}t^2 - \frac{23}{12}t^3\right) dt \\ &= \left[3t - t^2 + \frac{1}{2}t^3 - \frac{23}{48}t^4\right]_{t=0}^{t=1} \\ &= 3 - 1 + \frac{1}{2} - \frac{23}{48} = \frac{97}{48} \end{aligned}$$

(d) The alternating series error bound is the absolute value of the first omitted term of the series for $h(1)$.

$$\int_0^1 \left(\frac{54}{4!}t^4\right) dt = \left[\frac{9}{20}t^5\right]_{t=0}^{t=1} = \frac{9}{20}$$

$$\text{Error} \leq \left|\frac{9}{20}\right| = 0.45$$

$$2 : \begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \end{cases}$$

$$2 : \begin{cases} 1 : \text{three terms for } e^x \\ 1 : \text{three terms for } e^x f(x) \end{cases}$$

$$2 : \begin{cases} 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : \text{uses fourth-degree term} \\ \quad \text{of Maclaurin series for } f \\ 1 : \text{uses first omitted term} \\ \quad \text{of series for } h(1) \\ 1 : \text{error bound} \end{cases}$$

2017**AP[®]** **CollegeBoard**

AP Calculus BC

Scoring Guidelines

2017 SCORING GUIDELINES

Question 1

(a) Volume = $\int_0^{10} A(h) \, dh$ $\approx (2 - 0) \cdot A(0) + (5 - 2) \cdot A(2) + (10 - 5) \cdot A(5)$ $= 2 \cdot 50.3 + 3 \cdot 14.4 + 5 \cdot 6.5$ $= 176.3$ cubic feet (b) The approximation in part (a) is an overestimate because a left Riemann sum is used and A is decreasing. (c) $\int_0^{10} f(h) \, dh = 101.325338$ The volume is 101.325 cubic feet. (d) Using the model, $V(h) = \int_0^h f(x) \, dx$. $\begin{aligned} \left. \frac{dV}{dt} \right _{h=5} &= \left[\frac{dV}{dh} \cdot \frac{dh}{dt} \right]_{h=5} \\ &= \left[f(h) \cdot \frac{dh}{dt} \right]_{h=5} \\ &= f(5) \cdot 0.26 = 1.694419 \end{aligned}$ When $h = 5$, the volume of water is changing at a rate of 1.694 cubic feet per minute.	1 : units in parts (a), (c), and (d) 2 : $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{approximation} \end{cases}$ 1 : overestimate with reason 2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$ 3 : $\begin{cases} 2 : \frac{dV}{dt} \\ 1 : \text{answer} \end{cases}$
--	---

2017 SCORING GUIDELINES

Question 2

$$(a) \frac{1}{2} \int_0^{\pi/2} (f(\theta))^2 d\theta = 0.648414$$

The area of R is 0.648.

$$(b) \int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \frac{1}{2} \int_0^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

— OR —

$$\int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \int_k^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

$$(c) w(\theta) = g(\theta) - f(\theta)$$

$$w_A = \frac{\int_0^{\pi/2} w(\theta) d\theta}{\frac{\pi}{2} - 0} = 0.485446$$

The average value of $w(\theta)$ on the interval $\left[0, \frac{\pi}{2}\right]$ is 0.485.

$$(d) w(\theta) = w_A \text{ for } 0 \leq \theta \leq \frac{\pi}{2} \Rightarrow \theta = 0.517688$$

$$w(\theta) = w_A \text{ at } \theta = 0.518 \text{ (or } 0.517\text{)}.$$

$$w'(0.518) < 0 \Rightarrow w(\theta) \text{ is decreasing at } \theta = 0.518.$$

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

$$2 : \begin{cases} 1 : \text{integral expression} \\ \quad \text{for one region} \\ 1 : \text{equation} \end{cases}$$

$$3 : \begin{cases} 1 : w(\theta) \\ 1 : \text{integral} \\ 1 : \text{average value} \end{cases}$$

$$2 : \begin{cases} 1 : \text{solves } w(\theta) = w_A \\ 1 : \text{answer with reason} \end{cases}$$

2017 SCORING GUIDELINES

Question 3

$$(a) \quad f(-6) = f(-2) + \int_{-2}^{-6} f'(x) \, dx = 7 - \int_{-6}^{-2} f'(x) \, dx = 7 - 4 = 3$$

$$f(5) = f(-2) + \int_{-2}^5 f'(x) \, dx = 7 - 2\pi + 3 = 10 - 2\pi$$

$$3 : \begin{cases} 1 : \text{uses initial condition} \\ 1 : f(-6) \\ 1 : f(5) \end{cases}$$

$$(b) \quad f'(x) > 0 \text{ on the intervals } [-6, -2) \text{ and } (2, 5).$$

Therefore, f is increasing on the intervals $[-6, -2]$ and $[2, 5]$.

$$2 : \text{answer with justification}$$

$$(c) \quad \text{The absolute minimum will occur at a critical point where } f'(x) = 0 \text{ or at an endpoint.}$$

$$f'(x) = 0 \Rightarrow x = -2, x = 2$$

$$2 : \begin{cases} 1 : \text{considers } x = 2 \\ 1 : \text{answer with justification} \end{cases}$$

x	$f(x)$
-6	3
-2	7
2	$7 - 2\pi$
5	$10 - 2\pi$

The absolute minimum value is $f(2) = 7 - 2\pi$.

$$(d) \quad f''(-5) = \frac{2 - 0}{-6 - (-2)} = -\frac{1}{2}$$

$$2 : \begin{cases} 1 : f''(-5) \\ 1 : f''(3) \text{ does not exist,} \\ \text{with explanation} \end{cases}$$

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} = 2 \text{ and } \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3} = -1$$

$f''(3)$ does not exist because

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} \neq \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3}.$$

2017 SCORING GUIDELINES

Question 4

$$(a) \quad H'(0) = -\frac{1}{4}(91 - 27) = -16$$

$$H(0) = 91$$

An equation for the tangent line is $y = 91 - 16t$.

The internal temperature of the potato at time $t = 3$ minutes is approximately $91 - 16 \cdot 3 = 43$ degrees Celsius.

$$(b) \quad \frac{d^2H}{dt^2} = -\frac{1}{4} \frac{dH}{dt} = \left(-\frac{1}{4}\right)\left(-\frac{1}{4}\right)(H - 27) = \frac{1}{16}(H - 27)$$

$$H > 27 \text{ for } t > 0 \Rightarrow \frac{d^2H}{dt^2} = \frac{1}{16}(H - 27) > 0 \text{ for } t > 0$$

Therefore, the graph of H is concave up for $t > 0$. Thus, the answer in part (a) is an underestimate.

$$(c) \quad \frac{dG}{(G - 27)^{2/3}} = -dt$$

$$\int \frac{dG}{(G - 27)^{2/3}} = \int (-1) dt$$

$$3(G - 27)^{1/3} = -t + C$$

$$3(91 - 27)^{1/3} = 0 + C \Rightarrow C = 12$$

$$3(G - 27)^{1/3} = 12 - t$$

$$G(t) = 27 + \left(\frac{12 - t}{3}\right)^3 \text{ for } 0 \leq t < 10$$

The internal temperature of the potato at time $t = 3$ minutes is

$$27 + \left(\frac{12 - 3}{3}\right)^3 = 54 \text{ degrees Celsius.}$$

$$3 : \begin{cases} 1 : \text{slope} \\ 1 : \text{tangent line} \\ 1 : \text{approximation} \end{cases}$$

1 : underestimate with reason

$$5 : \begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration and} \\ \quad \text{uses initial condition} \\ 1 : \text{equation involving } G \text{ and } t \\ 1 : G(t) \text{ and } G(3) \end{cases}$$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

2017 SCORING GUIDELINES

Question 5

$$(a) f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2}$$

$$f'(3) = \frac{(-3)(5)}{(18 - 21 + 5)^2} = -\frac{15}{4}$$

$$(b) f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2} = 0 \Rightarrow x = \frac{7}{4}$$

The only critical point in the interval $1 < x < 2.5$ has x -coordinate $\frac{7}{4}$.

f' changes sign from positive to negative at $x = \frac{7}{4}$.

Therefore, f has a relative maximum at $x = \frac{7}{4}$.

$$\begin{aligned} (c) \int_5^\infty f(x) dx &= \lim_{b \rightarrow \infty} \int_5^b \frac{3}{2x^2 - 7x + 5} dx = \lim_{b \rightarrow \infty} \int_5^b \left(\frac{2}{2x - 5} - \frac{1}{x - 1} \right) dx \\ &= \lim_{b \rightarrow \infty} \left[\ln(2x - 5) - \ln(x - 1) \right]_5^b = \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2x - 5}{x - 1} \right) \right]_5^b \\ &= \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2b - 5}{b - 1} \right) - \ln \left(\frac{5}{4} \right) \right] = \ln 2 - \ln \left(\frac{5}{4} \right) = \ln \left(\frac{8}{5} \right) \end{aligned}$$

(d) f is continuous, positive, and decreasing on $[5, \infty)$.

The series converges by the integral test since $\int_5^\infty \frac{3}{2x^2 - 7x + 5} dx$ converges.

— OR —

$$\frac{3}{2n^2 - 7n + 5} > 0 \text{ and } \frac{1}{n^2} > 0 \text{ for } n \geq 5.$$

Since $\lim_{n \rightarrow \infty} \frac{\frac{3}{2n^2 - 7n + 5}}{\frac{1}{n^2}} = \frac{3}{2}$ and the series $\sum_{n=5}^\infty \frac{1}{n^2}$ converges,

the series $\sum_{n=5}^\infty \frac{3}{2n^2 - 7n + 5}$ converges by the limit comparison test.

2 : $f'(3)$

2 : $\begin{cases} 1 : x\text{-coordinate} \\ 1 : \text{relative maximum} \\ \text{with justification} \end{cases}$

3 : $\begin{cases} 1 : \text{antiderivative} \\ 1 : \text{limit expression} \\ 1 : \text{answer} \end{cases}$

2 : answer with conditions

2017 SCORING GUIDELINES

Question 6

(a) $f(0) = 0$

$f'(0) = 1$

$f''(0) = -1(1) = -1$

$f'''(0) = -2(-1) = 2$

$f^{(4)}(0) = -3(2) = -6$

The first four nonzero terms are

$$0 + 1x + \frac{-1}{2!}x^2 + \frac{2}{3!}x^3 + \frac{-6}{4!}x^4 = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4}.$$

The general term is $\frac{(-1)^{n+1}x^n}{n}$.

(b) For $x = 1$, the Maclaurin series becomes $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n}$.

The series does not converge absolutely because the harmonic series diverges.

The series alternates with terms that decrease in magnitude to 0, and therefore the series converges conditionally.

$$\begin{aligned} \text{(c)} \quad \int_0^x f(t) dt &= \int_0^x \left(t - \frac{t^2}{2} + \frac{t^3}{3} - \frac{t^4}{4} + \cdots + \frac{(-1)^{n+1}t^n}{n} + \cdots \right) dt \\ &= \left[\frac{t^2}{2} - \frac{t^3}{3 \cdot 2} + \frac{t^4}{4 \cdot 3} - \frac{t^5}{5 \cdot 4} + \cdots + \frac{(-1)^{n+1}t^{n+1}}{(n+1)n} + \cdots \right]_{t=0}^{t=x} \\ &= \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{12} - \frac{x^5}{20} + \cdots + \frac{(-1)^{n+1}x^{n+1}}{(n+1)n} + \cdots \end{aligned}$$

(d) The terms alternate in sign and decrease in magnitude to 0. By the alternating series error bound, the error $\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right|$ is bounded

by the magnitude of the first unused term, $\left| -\frac{(1/2)^5}{20} \right|$.

$$\text{Thus, } \left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right| \leq \left| -\frac{(1/2)^5}{20} \right| = \frac{1}{32 \cdot 20} < \frac{1}{500}.$$

$$3 : \begin{cases} 1 : f''(0), f'''(0), \text{ and } f^{(4)}(0) \\ 1 : \text{verify terms} \\ 1 : \text{general term} \end{cases}$$

2 : converges conditionally
with reason

$$3 : \begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \\ 1 : \text{general term} \end{cases}$$

1 : error bound