

Part A (AB or BC): Graphing calculator required**Question 1****9 points****General Scoring Notes**

- The model solution is presented using standard mathematical notation.
- Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be accurate to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

An invasive species of plant appears in a fruit grove at time $t = 0$ and begins to spread. The function C defined by $C(t) = 7.6 \arctan(0.2t)$ models the number of acres in the fruit grove affected by the species t weeks after the species appears. It can be shown that $C'(t) = \frac{38}{25 + t^2}$.

(Note: Your calculator should be in radian mode.)

	Model Solution	Scoring
A	Find the average number of acres affected by the invasive species from time $t = 0$ to time $t = 4$ weeks. Show the setup for your calculations.	
	$\frac{1}{4 - 0} \int_0^4 C(t) dt$	Average value formula Point 1 (P1)
	$= \frac{1}{4}(11.112896) = 2.778224$	Answer Point 2 (P2)
	From time $t = 0$ to $t = 4$ weeks, the average number of acres affected by the invasive species was 2.778 acres.	

Scoring Notes for Part A

- P1** is earned for the correct integral, with or without the differential, along with evidence of division by 4. In the presence of the correct integral, the correct answer will suffice as evidence of division by 4. These may appear all in one step, as in the model solution, or in multiple steps.
- P2** is earned for the correct answer, with or without supporting work. A reported answer should be accurate to three places after the decimal point, rounded or truncated. An inappropriately rounded answer does not earn the point.
- Incorrect or unclear communication between the correct integral and the correct answer is treated as scratch work and is not considered in scoring. For example:
 - $\int_0^4 C(t) dt = 11.112896$ so the average velocity is 2.778224.
Note: This response earns **P1** for the correct integral with the correct answer as evidence of division by 4. It also earns **P2** for the correct answer.
 - $\int_0^4 C(t) dt = \frac{11.112896}{4} = 2.778224$
Note: This response earns **P1** for the correct integral with the correct answer as evidence of division by 4. It also earns **P2** for the correct answer. (In this instance, incorrect linkage is not considered in scoring.)
 - $\int_0^4 C(t) dt = 2.778224$
Note: This response earns **P1** for the correct integral with the correct answer as evidence of division by 4. It also earns **P2** for the correct answer. (In this instance, incorrect linkage is not considered in scoring.)
- Note that the values $\frac{1}{4}(11.112)$ and $\frac{1}{4}(11.113)$ are accurate to three digits after the decimal and therefore earn **P2**.

- B** Find the time t when the instantaneous rate of change of C equals the average rate of change of C over the time interval $0 \leq t \leq 4$. Show the setup for your calculations.

$$\frac{C(4) - C(0)}{4 - 0} = 1.282008$$

Uses average rate of change **Point 3 (P3)**

$$C'(t) = \frac{38}{25 + t^2} = 1.282008 \Rightarrow t = 2.154298$$

Answer with supporting work **Point 4 (P4)**

The instantaneous rate of change of C equals the average rate of change of C over the interval $0 \leq t \leq 4$ at time $t = 2.154$.

Scoring Notes for Part B

- P3** may be earned by presenting the expression or value for the average rate of change. Note that because $C(0) = 0$ and the interval is $0 \leq t \leq 4$, any of the following will earn **P3**: $\frac{\int_0^4 C'(t) dt}{4}$, $\frac{C(4) - C(0)}{4 - 0}$, $\frac{C(4)}{4}$, $\frac{5.128 - 0}{4 - 0}$, $\frac{5.128}{4}$, or 1.282. However, neither **P3** nor **P4** is earned by just presenting $t = 1.282$.
 - P4** is earned for the correct answer supported by the appropriate equation. A reported answer should be accurate to three places after the decimal point, rounded or truncated. An inappropriately rounded answer does not earn the point, unless an earlier point was not earned due to inappropriate rounding.
- The following response, for example, earns both **P3** and **P4**: $C'(t) = \frac{C(4) - C(0)}{4}$ when $t = 2.154$.

- C** Assume that the invasive species continues to spread according to the given model for all times $t > 0$. Write a limit expression that describes the end behavior of the rate of change in the number of acres affected by the species. Evaluate this limit expression.

$$\lim_{t \rightarrow \infty} C'(t) = \lim_{t \rightarrow \infty} \frac{38}{25 + t^2}$$

Limit expression **Point 5 (P5)**

$$= 0$$

Value **Point 6 (P6)**

Scoring Notes for Part C

- P5** can be earned for either $\lim_{t \rightarrow \infty} C'(t)$ or $\lim_{t \rightarrow \infty} C(t)$.
- A response that includes $\lim_{t \rightarrow \infty} C(t)$ is not eligible to earn **P6**.
- For **P6**, arithmetic with infinity, e.g., $\frac{38}{25 + \infty^2} = 0$, will be considered as scratch work and will not be considered in scoring.

- D** At time $t = 4$ weeks after the invasive species appears in the fruit grove, measures are taken to counter the spread of the species. The function A , defined by $A(t) = C(t) - \int_4^t 0.1 \cdot \ln(x) dx$, models the number of acres affected by the species over the time interval $4 \leq t \leq 36$. At what time t , for $4 \leq t \leq 36$, does A attain its maximum value? Justify your answer.

$$A'(t) = C'(t) - 0.1 \cdot \ln t$$

Considers $A'(t) = 0$ **Point 7 (P7)**

For $4 \leq t \leq 36$, the maximum value of $A(t)$ occurs when $A'(t) = 0$ or at an endpoint.

$$A'(t) = C'(t) - 0.1 \cdot \ln t = 0 \Rightarrow C'(t) = 0.1 \cdot \ln t$$

$$\Rightarrow t = 11.441700$$

Justification **Point 8 (P8)**

t	$A(t)$
4	5.128031
11.441700	7.316978
36	1.743056

Therefore, the number of acres affected by the species is a maximum at time $t = 11.442$ (or 11.441) weeks.

Answer with supporting work **Point 9 (P9)**

Scoring Notes for Part D

- P7** is earned for considering $A'(t) = 0$, $C'(t) - 0.1 \cdot \ln t = 0$, or $C'(t) = 0.1 \cdot \ln t$. **P7** is not earned by just presenting $t = 11.441700$. A response that discusses the sign of $A'(t)$ changing or uses the phrase “critical points of A ” also earns **P7**.
- To earn **P8** using a candidates test, a response must make a global argument by correctly evaluating $A(t)$ at $t = 4$, $t = 11.441700$, and $t = 36$. The evaluations must be correct to the first digit after the decimal, rounded or truncated.
- Alternate justifications:
 - $A'(t) > 0$ for $4 < t < 11.442$, and $A'(t) < 0$ for $11.442 < t < 36$. Therefore, $t = 11.442$ is the location of the absolute maximum for A on the interval $4 \leq t \leq 36$.
 - Because $A'(t)$ changes sign from positive to negative at $t = 11.442$ (this might be presented as “ $A'(t) > 0$ for $t < 11.442$, and $A'(t) < 0$ for $t > 11.442$ ”), it is the location of a relative maximum for A . And because $t = 11.442$ is the only critical point of A in the interval $4 \leq t \leq 36$, it is the location of the absolute maximum for A on the interval.
- A response that presents a local argument (such as a First Derivative Test or a Second Derivative Test) or an incorrect global argument does not earn **P8** but is eligible for **P9** with the correct answer. A reported answer should be accurate to three places after the decimal point, rounded or truncated. An inappropriately rounded answer does not earn the point, unless an earlier point was not earned due to inappropriate rounding.

Part A (AB or BC): Graphing calculator required**Question 1****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

t (minutes)	0	3	7	12
$C(t)$ (degrees Celsius)	100	85	69	55

The temperature of coffee in a cup at time t minutes is modeled by a decreasing differentiable function C , where $C(t)$ is measured in degrees Celsius. For $0 \leq t \leq 12$, selected values of $C(t)$ are given in the table shown.

Model Solution**Scoring**

- (a) Approximate $C'(5)$ using the average rate of change of C over the interval $3 \leq t \leq 7$. Show the work that leads to your answer and include units of measure.

$$C'(5) \approx \frac{C(7) - C(3)}{7 - 3} = \frac{69 - 85}{4} = -4 \text{ degrees Celsius per minute}$$

Estimate with supporting work	1 point
Units	1 point

Scoring notes:

- To earn the first point a response must include a difference and a quotient as the supporting work.
- $\frac{-16}{7-3}$, $\frac{69-85}{7-3}$, or $\frac{69-85}{4}$ is sufficient to earn the first point.
- A response that presents only units without a numerical approximation for $C'(5)$ does not earn the second point.
- The second point is also earned for “degrees per minute” attached to a numerical value.

Total for part (a) 2 points

- (b) Use a left Riemann sum with the three subintervals indicated by the data in the table to approximate the value of $\int_0^{12} C(t) dt$. Interpret the meaning of $\frac{1}{12} \int_0^{12} C(t) dt$ in the context of the problem.

$\int_0^{12} C(t) dt \approx (3-0) \cdot C(0) + (7-3) \cdot C(3) + (12-7) \cdot C(7)$ $= 3 \cdot 100 + 4 \cdot 85 + 5 \cdot 69 = 985$	Form of left Riemann sum	1 point
	Estimate	1 point
$\frac{1}{12} \int_0^{12} C(t) dt$ is the average temperature of the coffee (in degrees Celsius) over the interval from $t = 0$ to $t = 12$.	Interpretation	1 point

Scoring notes:

- Read “=” as “ $\approx$ ” for the first point.
- To earn the first point at least five of the six factors in the Riemann sum must be correct. If any of the six factors is incorrect, the response does not earn the second point.
- A response of $(3-0) \cdot C(0) + (7-3) \cdot C(3) + (12-7) \cdot C(7)$ earns the first point. Values must be pulled from the table to earn the second point.
- A response of $3 \cdot 100 + 4 \cdot 85 + 5 \cdot 69$ earns both the first and second points, unless there is a subsequent error in simplification, in which case the response would earn only the first point.
- A completely correct right Riemann sum (e.g., $3 \cdot 85 + 4 \cdot 69 + 5 \cdot 55$) earns 1 of the first 2 points. An unsupported answer of 806 does not earn either of the first 2 points.
- Units will not affect scoring for the second point.
- To earn the third point the interpretation must include both “average temperature” and the time interval. The response need not include a reference to units. However, if incorrect units are given in the interpretation, the response does not earn the third point.

Total for part (b) 3 points

- (c) For $12 \leq t \leq 20$, the rate of change of the temperature of the coffee is modeled by

$$C'(t) = \frac{-24.55e^{0.01t}}{t}, \text{ where } C'(t) \text{ is measured in degrees Celsius per minute. Find the temperature of the coffee at time } t = 20. \text{ Show the setup for your calculations.}$$

$C(20) = C(12) + \int_{12}^{20} C'(t) dt$	Integral	1 point
$= 55 - 14.670812 = 40.329188$	Uses initial condition	1 point
	Answer	1 point
The temperature of the coffee at time $t = 20$ is 40.329 degrees Celsius.		

Scoring notes:

- The first point is earned for a definite integral with integrand $C'(t)$. If the limits of integration are incorrect, the response does not earn the third point.
- A linkage error such as $C(20) = \int_{12}^{20} C'(t) dt = 55 - 14.670812$ or $\int_{12}^{20} C'(t) dt = -14.670812 = 40.329188$ earns the first 2 points but does not earn the third point.
- Missing differential (dt):
 - Unambiguous responses of $C(20) = C(12) + \int_{12}^{20} C'(t)$ or $C(20) = 55 + \int_{12}^{20} C'(t)$ earn the first 2 points and are eligible for the third point.
 - Ambiguous responses of $C(20) = \int_{12}^{20} C'(t) + C(12)$ or $C(20) = \int_{12}^{20} C'(t) + 55$ do not earn the first point, earn the second point, and earn the third point if the given numeric answer is correct. If there is no numeric answer given, these responses do not earn the third point.
- The second point is earned for adding $C(12)$ or 55 to a definite integral with a lower limit of 12, either symbolically or numerically.
- The third point is earned for an answer of $55 - 14.671$ or $-14.671 + 55$ with no additional simplification, provided there is some supporting work for these values.
- An answer of just 40.329 with no supporting work does not earn any points.

Total for part (c) 3 points

- (d) For the model defined in part (c), it can be shown that $C''(t) = \frac{0.2455e^{0.01t}(100-t)}{t^2}$. For

$12 < t < 20$, determine whether the temperature of the coffee is changing at a decreasing rate or at an increasing rate. Give a reason for your answer.

Because $C''(t) > 0$ on the interval $12 < t < 20$, the rate of change in the temperature of the coffee, $C'(t)$, is increasing on this interval.	Answer with reason	1 point
---	--------------------	----------------

That is, on the interval $12 < t < 20$, the temperature of the coffee is changing at an increasing rate.

Scoring notes:

- This point is earned only for a correct answer with a correct reason that references the sign of the second derivative of C .
- A response that provides a reason based on the evaluation of $C''(t)$ at a single point does not earn this point.
- A response that uses ambiguous pronouns (such as “It is positive, so increasing”) does not earn this point.
- A response does not need to reference the interval $12 < t < 20$ to earn the point.

Total for part (d) 1 point**Total for question 1 9 points****Part A (AB or BC): Graphing calculator required****Question 1****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

t (seconds)	0	60	90	120	135	150
$f(t)$ (gallons per second)	0	0.1	0.15	0.1	0.05	0

A customer at a gas station is pumping gasoline into a gas tank. The rate of flow of gasoline is modeled by a differentiable function f , where $f(t)$ is measured in gallons per second and t is measured in seconds since pumping began. Selected values of $f(t)$ are given in the table.

Model Solution	Scoring
(a) Using correct units, interpret the meaning of $\int_{60}^{135} f(t) dt$ in the context of the problem. Use a right Riemann sum with the three subintervals $[60, 90]$, $[90, 120]$, and $[120, 135]$ to approximate the value of $\int_{60}^{135} f(t) dt$.	
$\int_{60}^{135} f(t) dt$ represents the total number of gallons of gasoline pumped into the gas tank from time $t = 60$ seconds to time $t = 135$ seconds.	Interpretation with units 1 point
$\int_{60}^{135} f(t) dt$ $\approx f(90)(90 - 60) + f(120)(120 - 90) + f(135)(135 - 120)$ $= (0.15)(30) + (0.1)(30) + (0.05)(15) = 8.25$	Form of Riemann sum 1 point Answer 1 point

Scoring notes:

- To earn the first point the response must reference gallons of gasoline added/pumped and the time interval $t = 60$ to $t = 135$.
- To earn the second point at least five of the six factors in the Riemann sum must be correct.
- If there is any error in the Riemann sum, the response does not earn the third point.
- A response of $(0.15)(30) + (0.1)(30) + (0.05)(15)$ earns both the second and third points, unless there is a subsequent error in simplification, in which case the response would earn only the second point.

- A response that presents a correct value with accompanying work that shows the three products in the Riemann sum but does not show all six of the factors and/or the sum process, does not earn the second point but does earn the third point. For example, responses of either $4.5 + 3.0 + 0.75$ or $(0.15)(30)$, $0.1(30)$, $0.05(15) \rightarrow 8.25$ earn the third point but not the second.
- A response of $f(90)(90 - 60) + f(120)(120 - 90) + f(135)(135 - 120) = 8.25$ earns both the second and the third points.
- A response that presents an answer of only 8.25 does not earn either the second or third point.
- A response that provides a completely correct left Riemann sum with accompanying work, $f(60)(30) + f(90)(30) + f(120)(15) = 9$, or $(0.1)(30) + (0.15)(30) + (0.1)(15)$ earns 1 of the last 2 points. A response with any errors or missing factors in a left Riemann sum earns neither of the last 2 points.

Total for part (a) 3 points

- (b) Must there exist a value of c , for $60 < c < 120$, such that $f'(c) = 0$? Justify your answer.

f is differentiable. $\Rightarrow f$ is continuous on $[60, 120]$.

$f(120) - f(60) = 0$ **1 point**

$$\frac{f(120) - f(60)}{120 - 60} = \frac{0.1 - 0.1}{60} = 0$$

Answer with justification **1 point**

By the Mean Value Theorem, there must exist a c , for $60 < c < 120$, such that $f'(c) = 0$.

Scoring notes:

- To earn the first point a response must present either $f(120) - f(60) = 0$, $0.1 - 0.1 = 0$ (perhaps as the numerator of a quotient), or $f(60) = f(120)$.
- To earn the second point a response must:
 - have earned the first point,
 - state that f is continuous because f is differentiable (or equivalent), and
 - answer “yes” in some way.
- A response may reference either the Mean Value Theorem or Rolle’s Theorem.
- A response that references the Intermediate Value Theorem cannot earn the second point.

Total for part (b) 2 points

- (c) The rate of flow of gasoline, in gallons per second, can also be modeled by

$$g(t) = \left(\frac{t}{500}\right) \cos\left(\left(\frac{t}{120}\right)^2\right) \text{ for } 0 \leq t \leq 150. \text{ Using this model, find the average rate of flow of}$$

gasoline over the time interval $0 \leq t \leq 150$. Show the setup for your calculations.

$$\frac{1}{150 - 0} \int_0^{150} g(t) dt$$

Average value formula **1 point**

$$= 0.0959967$$

Answer **1 point**

The average rate of flow of gasoline, in gallons per second, is 0.096 (or 0.095).

Scoring notes:

- The exact value of $\frac{1}{150} \int_0^{150} g(t) dt$ is $\frac{12}{125} \sin\left(\frac{25}{16}\right)$.
- A response may present the average value formula in single or multiple steps. For example, the following response earns both points: $\int_0^{150} g(t) dt = 14.399504$ so the average rate is 0.0959967.
- A response that presents the average value formula in multiple steps but provides incorrect or incomplete communication (e.g., $\int_0^{150} g(t) dt = \frac{14.399504}{150} = 0.0959967$) earns 1 out of 2 points.
- A response of $\int_0^{150} g(t) dt = 0.0959967$ does not earn either point.
- Degree mode: A response that presents answers obtained by using a calculator in degree mode does not earn the first point it would have otherwise earned. The response is generally eligible for all subsequent points (unless no answer is possible in degree mode or the question is made simpler by using degree mode). In degree mode, $\frac{1}{150} \int_0^{150} g(t) dt = 0.149981$ or 0.002618.

Total for part (c) 2 points

- (d) Using the model g defined in part (c), find the value of $g'(140)$. Interpret the meaning of your answer in the context of the problem.

$$g'(140) \approx -0.004908$$

$g'(140)$ **1 point**

$$g'(140) = -0.005 \text{ (or } -0.004 \text{)}$$

The rate at which gasoline is flowing into the tank is decreasing at a rate of 0.005 (or 0.004) gallon per second per second at time $t = 140$ seconds.

Interpretation **1 point**

Scoring notes:

- The exact value of $g'(140)$ is $\frac{1}{500} \cos\left(\frac{49}{36}\right) - \frac{49}{9000} \sin\left(\frac{49}{36}\right)$.
- The value of $g'(140)$ may appear only in the interpretation.
- To be eligible for the second point a response must present some numerical value for $g'(140)$.
- To earn the second point the interpretation must include “the rate of flow of gasoline is changing at a rate of [the declared value of $g'(140)$]” and “at $t = 140$ ” (or equivalent).
- An interpretation of “decreasing at a rate of -0.005 ” or “increasing at a rate of 0.005” does not earn the second point.
- Degree mode: In degree mode, $g'(140) = 0.001997$ or 0.00187.

Total for part (d) 2 points**Total for question 1 9 points**

Part A (AB or BC): Graphing calculator required**Question 1****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

From 5 A.M. to 10 A.M., the rate at which vehicles arrive at a certain toll plaza is given by $A(t) = 450\sqrt{\sin(0.62t)}$, where t is the number of hours after 5 A.M. and $A(t)$ is measured in vehicles per hour. Traffic is flowing smoothly at 5 A.M. with no vehicles waiting in line.

Model Solution		Scoring
(a) Write, but do not evaluate, an integral expression that gives the total number of vehicles that arrive at the toll plaza from 6 A.M. ($t = 1$) to 10 A.M. ($t = 5$).		
The total number of vehicles that arrive at the toll plaza from 6 A.M. to 10 A.M. is given by $\int_1^5 A(t) dt$.	Answer	1 point
Scoring notes: - The response must be a definite integral with correct lower and upper limits to earn this point. - Because $A(t) = A(t)$ for $1 \leq t \leq 5$, a response of $\int_1^5 450\sqrt{\sin(0.62t)} dt$ or $\int_1^5 A(t) dt$ earns the point. - A response missing dt or using dx is eligible to earn the point. - A response with a copy error in the expression for $A(t)$ will earn the point only in the presence of $\int_1^5 A(t) dt$.		
Total for part (a)		1 point

- (b) Find the average value of the rate, in vehicles per hour, at which vehicles arrive at the toll plaza from 6 A.M. ($t = 1$) to 10 A.M. ($t = 5$).

Average = $\frac{1}{5-1} \int_1^5 A(t) dt = 375.536966$	Uses average value formula: $\frac{1}{b-a} \int_a^b A(t) dt$	1 point
The average rate at which vehicles arrive at the toll plaza from 6 A.M. to 10 A.M. is 375.537 (or 375.536) vehicles per hour.	Answer	1 point

Scoring notes:

- The use of the average value formula, indicating that $a = 1$ and $b = 5$, can be presented in single or multiple steps to earn the first point. For example, the following response earns both points:
 $\int_1^5 A(t) dt = 1502.147865$, so the average value is 375.536966 .
- A response that presents a correct integral along with the correct average value, but provides incorrect or incomplete communication, earns 1 out of 2 points. For example, the following response earns 1 out of 2 points: $\int_1^5 A(t) dt = 1502.147865 = 375.536966$.
- The answer must be correct to three decimal places. For example,
 $\frac{1}{5-1} \int_1^5 A(t) dt = 375.536966 \approx 376$ earns only the first point.
- Degree mode: A response that presents answers obtained by using a calculator in degree mode does not earn the first point it would have otherwise earned. The response is generally eligible for all subsequent points (unless no answer is possible in degree mode or the question is made simpler by using degree mode). In degree mode, $\frac{1}{4} \int_1^5 A(t) dt = 79.416068$.
- Special case: $\frac{1}{5} \int_1^5 A(t) dt = 300.429573$ earns 1 out of 2 points.

Total for part (b) 2 points

- (c) Is the rate at which vehicles arrive at the toll plaza at 6 A.M. ($t = 1$) increasing or decreasing? Give a reason for your answer.

$A'(1) = 148.947272$	Considers $A'(1)$	1 point
Because $A'(1) > 0$, the rate at which the vehicles arrive at the toll plaza is increasing.	Answer with reason	1 point

Scoring notes:

- The response need not present the value of $A'(1)$. The second line of the model solution earns both points.
- An incorrect value assigned to $A'(1)$ earns the first point (but will not earn the second point).
- Without a reference to $t = 1$, the first point is earned by any of the following:
 - 148.947 accurate to the number of decimals presented, with zero up to three decimal places (i.e., 149, 148, 148.9, 148.95, or 148.94)
 - $A'(t) = 148.947$ by itself
- To be eligible for the second point, the first point must be earned.
- To earn the second point, there must be a reference to $t = 1$.
- Degree mode: $A'(1) = 23.404311$

Total for part (c) 2 points

- (d) A line forms whenever $A(t) \geq 400$. The number of vehicles in line at time t , for $a \leq t \leq 4$, is given by $N(t) = \int_a^t (A(x) - 400) dx$, where a is the time when a line first begins to form. To the nearest whole number, find the greatest number of vehicles in line at the toll plaza in the time interval $a \leq t \leq 4$. Justify your answer.

$N'(t) = A(t) - 400 = 0$ $\Rightarrow A(t) = 400 \Rightarrow t = 1.469372, t = 3.597713$	Considers $N'(t) = 0$	1 point								
	$t = a$ and $t = b$	1 point								
$a = 1.469372$ $b = 3.597713$										
<table><tr><td>t</td><td>$N(t) = \int_a^t (A(x) - 400) dx$</td></tr><tr><td>$a$</td><td>0</td></tr><tr><td>b</td><td>71.254129</td></tr><tr><td>4</td><td>62.338346</td></tr></table>	t	$N(t) = \int_a^t (A(x) - 400) dx$	a	0	b	71.254129	4	62.338346	Answer	1 point
t	$N(t) = \int_a^t (A(x) - 400) dx$									
a	0									
b	71.254129									
4	62.338346									
	Justification	1 point								

The greatest number of vehicles in line is 71.

Scoring notes:

- It is not necessary to indicate that $A(t) = 400$ to earn the first point, although this statement alone would earn the first point.
- A response of “ $A(t) \geq 400$ when $1.469372 \leq t \leq 3.597713$ ” will earn the first 2 points. A response of “ $A(t) \geq 400$ ” along with the presence of exactly one of the two numbers above will earn the first point, but not the second. A response of “ $A(t) \geq 400$ ” by itself will not earn either of the first 2 points.
- To earn the second point the values for a and b must be accurate to the number of decimals presented, with at least one and up to three decimal places. These may appear only in a candidates table, as limits of integration, or on a number line.
- A response with incorrect notation involving t or x is eligible to earn all 4 points.
- A response that does not earn the first point is still eligible for the remaining 3 points.
- To earn the third point, a response must present the greatest number of vehicles. This point is earned for answers of either 71 or 71.254*** only.
- A correct justification earns the fourth point, even if the third point is not earned because of a decimal presentation error.
- When using a Candidates Test, the response must include the values for $N(a)$, $N(b)$, and $N(4)$ to earn the fourth point. These values must be correct to the number of decimals presented, with up to three decimal places. (Correctly rounded integer values are acceptable.)
- Alternate solution for the third and fourth points:
For $a \leq t \leq b$, $A(t) \geq 400$. For $b \leq t \leq 4$, $A(t) \leq 400$.
Thus, $N(t) = \int_a^t (A(x) - 400) dx$ is greatest at $t = b$.
 $N(b) = 71.254129$, and the greatest number of vehicles in line is 71.
- Degree mode: The response is only eligible to earn the first point because in degree mode $A(t) < 400$.

Total for part (d) 4 points**Total for question 1 9 points**

Part A (AB or BC): Graphing calculator required**Question 1****9 points****General Scoring Notes**

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Scoring guidelines and notes contain examples of the most common approaches seen in student responses. These guidelines can be applied to alternate approaches to ensure that these alternate approaches are scored appropriately.

r (centimeters)	0	1	2	2.5	4
$f(r)$ (milligrams per square centimeter)	1	2	6	10	18

The density of a bacteria population in a circular petri dish at a distance r centimeters from the center of the dish is given by an increasing, differentiable function f , where $f(r)$ is measured in milligrams per square centimeter. Values of $f(r)$ for selected values of r are given in the table above.

Model Solution**Scoring**

- (a) Use the data in the table to estimate $f'(2.25)$. Using correct units, interpret the meaning of your answer in the context of this problem.

$$f'(2.25) \approx \frac{f(2.5) - f(2)}{2.5 - 2} = \frac{10 - 6}{0.5} = 8$$

Estimate **1 point**

At a distance of $r = 2.25$ centimeters from the center of the petri dish, the density of the bacteria population is increasing at a rate of 8 milligrams per square centimeter per centimeter.

Interpretation with units **1 point****Scoring notes:**

- To earn the first point the response must provide both a difference and a quotient and must explicitly use values of f from the table.
- Simplification of the numerical value is not required to earn the first point. If the numerical value is simplified, it must be correct.
- The interpretation requires all of the following: distance $r = 2.25$, density of bacteria (population) is increasing or changing, at a rate of 8, and units of milligrams per square centimeter per centimeter.
- The second point (interpretation) cannot be earned without a nonzero presented value for $f'(2.25)$.
- To earn the second point the interpretation must be consistent with the presented nonzero value for $f'(2.25)$. In particular, if the presented value for $f'(2.25)$ is negative, the interpretation must include “decreasing at a rate of $|f'(2.25)|$ ” or “changing at a rate of $f'(2.25)$.” The second point cannot be earned for an incorrect statement such as “the bacteria density is decreasing at a rate of $-8 \dots$ ” even for a presented $f'(2.25) = -8$.
- The units (mg/cm²/cm) may be attached to the estimate of $f'(2.25)$ and, if so, do not need to be repeated in the interpretation.
- If units attached to the estimate do not agree with units in the interpretation, read the units in the interpretation.

Total for part (a) 2 points

- (b) The total mass, in milligrams, of bacteria in the petri dish is given by the integral expression $2\pi \int_0^4 r f(r) dr$. Approximate the value of $2\pi \int_0^4 r f(r) dr$ using a right Riemann sum with the four subintervals indicated by the data in the table.

$2\pi \int_0^4 r f(r) dr \approx 2\pi(1 \cdot f(1) \cdot (1 - 0) + 2 \cdot f(2) \cdot (2 - 1) + 2.5 \cdot f(2.5) \cdot (2.5 - 2) + 4 \cdot f(4) \cdot (4 - 2.5))$	Right Riemann sum setup	1 point
$= 2\pi(1 \cdot 2 \cdot 1 + 2 \cdot 6 \cdot 1 + 2.5 \cdot 10 \cdot 0.5 + 4 \cdot 18 \cdot 1.5)$ $= 269\pi \approx 845.088$	Approximation	1 point

Scoring notes:

- The presence or absence of 2π has no bearing on earning the first point.
- The first point is earned for a sum of four products with at most one error in any single value among the four products. Multiplication by 1 in any term does not need to be shown, but all other products must be explicitly shown.
- A response of $1 \cdot f(1) \cdot (1 - 0) + 2 \cdot f(2) \cdot (2 - 1) + 2.5 \cdot f(2.5) \cdot (2.5 - 2) + 4 \cdot f(4) \cdot (4 - 2.5)$ earns the first point but not the second.
- A response with any error in the Riemann sum is not eligible for the second point.
- A response that provides a completely correct left Riemann sum for $2\pi \int_0^4 r f(r) dr$ and approximation (91π) earns one of the two points. A response that has any error in a left Riemann sum or evaluation for $2\pi \int_0^4 r f(r) dr$ earns no points.
- A response that provides a completely correct right Riemann sum for $2\pi \int_0^4 r f(r) dr$ and approximation (80π) earns one of the two points. A response that has any error in a right Riemann sum or evaluation for $2\pi \int_0^4 r f(r) dr$ earns no points.
- Simplification of the numerical value is not required to earn the second point. If a numerical value is given, it must be correct to three decimal places.

Total for part (b) 2 points

- (c) Is the approximation found in part (b) an overestimate or underestimate of the total mass of bacteria in the petri dish? Explain your reasoning.

$\frac{d}{dr}(rf(r)) = f(r) + rf'(r)$	Product rule expression for $\frac{d}{dr}(rf(r))$	1 point
Because f is nonnegative and increasing, $\frac{d}{dr}(rf(r)) > 0$ on the interval $0 \leq r \leq 4$. Thus, the integrand $rf(r)$ is strictly increasing.	Answer with explanation	1 point
Therefore, the right Riemann sum approximation of $2\pi \int_0^4 r f(r) dr$ is an overestimate.		

Scoring notes:

- To earn the second point a response must explain that $rf(r)$ is increasing and, therefore, the right Riemann sum is an overestimate. The second point can be earned without having earned the first point.
- A response that attempts to explain based on a left Riemann sum for $2\pi \int_0^4 r f(r) dr$ from part (b) earns no points.
- A response that attempts to explain based on a right Riemann sum for $2\pi \int_0^4 r f(r) dr$ from part (b) earns no points.

Total for part (c) 2 points

- (d) The density of bacteria in the petri dish, for $1 \leq r \leq 4$, is modeled by the function g defined by $g(r) = 2 - 16(\cos(1.57\sqrt{r}))^3$. For what value of k , $1 < k < 4$, is $g(k)$ equal to the average value of $g(r)$ on the interval $1 \leq r \leq 4$?

Average value $= g_{\text{avg}} = \frac{1}{4-1} \int_1^4 g(r) dr$	Definite integral	1 point
$\frac{1}{4-1} \int_1^4 g(r) dr = 9.875795$	Average value	1 point
$g(k) = g_{\text{avg}} \Rightarrow k = 2.497$	Answer	1 point

Scoring notes:

- The first point is earned for a definite integral, with or without $\frac{1}{4-1}$ or $\frac{1}{3}$.
- A response that presents a definite integral with incorrect limits but a correct integrand earns the first point.
- Presentation of the numerical value 9.875795 is not required to earn the second point. This point can be earned by the average value setup: $\frac{1}{3} \int_1^4 g(r) dr$.
- Once earned for the average value setup, the second point cannot be lost. Subsequent errors will result in not earning the third point.
- The third point is earned only for the value $k = 2.497$.
- The third point cannot be earned without the second.
- Special case: A response that does not provide the average value setup but presents an average value of -13.955 is using degree mode on their calculator. This response would not earn the second point but could earn the third point for an answer of $k = 2.5$ (or 2.499).

Total for part (d) 3 points**Total for question 1 9 points**

2019

AP[®] CollegeBoard

AP[®] Calculus BC

Scoring Guidelines

2019 SCORING GUIDELINES**Question 1**

(a) $\int_0^5 E(t) dt = 153.457690$

To the nearest whole number, 153 fish enter the lake from midnight to 5 A.M.

(b) $\frac{1}{5-0} \int_0^5 L(t) dt = 6.059038$

The average number of fish that leave the lake per hour from midnight to 5 A.M. is 6.059 fish per hour.

- (c) The rate of change in the number of fish in the lake at time t is given by $E(t) - L(t)$.

$$E(t) - L(t) = 0 \Rightarrow t = 6.20356$$

$E(t) - L(t) > 0$ for $0 \leq t < 6.20356$, and $E(t) - L(t) < 0$ for $6.20356 < t \leq 8$. Therefore the greatest number of fish in the lake is at time $t = 6.204$ (or 6.203).

— OR —

Let $A(t)$ be the change in the number of fish in the lake from midnight to t hours after midnight.

$$A(t) = \int_0^t (E(s) - L(s)) ds$$

$$A'(t) = E(t) - L(t) = 0 \Rightarrow t = C = 6.20356$$

t	$A(t)$
0	0
C	135.01492
8	80.91998

Therefore the greatest number of fish in the lake is at time $t = 6.204$ (or 6.203).

(d) $E'(5) - L'(5) = -10.7228 < 0$

Because $E'(5) - L'(5) < 0$, the rate of change in the number of fish is decreasing at time $t = 5$.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

3 : $\begin{cases} 1 : \text{sets } E(t) - L(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

2 : $\begin{cases} 1 : \text{considers } E'(5) \text{ and } L'(5) \\ 1 : \text{answer with explanation} \end{cases}$

2019 SCORING GUIDELINES

Question 2

(a) $\frac{1}{2} \int_0^{\sqrt{\pi}} (r(\theta))^2 d\theta = 3.534292$

The area of S is 3.534.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) $\frac{1}{\sqrt{\pi}-0} \int_0^{\sqrt{\pi}} r(\theta) d\theta = 1.579933$

The average distance from the origin to a point on the curve $r = r(\theta)$ for $0 \leq \theta \leq \sqrt{\pi}$ is 1.580 (or 1.579).

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(c) $\tan \theta = \frac{y}{x} = m \Rightarrow \theta = \tan^{-1} m$

$$\frac{1}{2} \int_0^{\tan^{-1} m} (r(\theta))^2 d\theta = \frac{1}{2} \left(\frac{1}{2} \int_0^{\sqrt{\pi}} (r(\theta))^2 d\theta \right)$$

3 : $\begin{cases} 1 : \text{equates polar areas} \\ 1 : \text{inverse trigonometric function} \\ \text{applied to } m \text{ as limit of} \\ \text{integration} \\ 1 : \text{equation} \end{cases}$

(d) As $k \rightarrow \infty$, the circle $r = k \cos \theta$ grows to enclose all points to the right of the y -axis.

$$\begin{aligned} \lim_{k \rightarrow \infty} A(k) &= \frac{1}{2} \int_0^{\pi/2} (r(\theta))^2 d\theta \\ &= \frac{1}{2} \int_0^{\pi/2} (3\sqrt{\theta} \sin(\theta^2))^2 d\theta = 3.324 \end{aligned}$$

2 : $\begin{cases} 1 : \text{limits of integration} \\ 1 : \text{answer with integral} \end{cases}$

2019 SCORING GUIDELINES

Question 3

(a) $\int_{-6}^5 f(x) dx = \int_{-6}^{-2} f(x) dx + \int_{-2}^5 f(x) dx$

$$\Rightarrow 7 = \int_{-6}^{-2} f(x) dx + 2 + \left(9 - \frac{9\pi}{4}\right)$$

$$\Rightarrow \int_{-6}^{-2} f(x) dx = 7 - \left(11 - \frac{9\pi}{4}\right) = \frac{9\pi}{4} - 4$$

3 : $\begin{cases} 1 : \int_{-6}^5 f(x) dx = \int_{-6}^{-2} f(x) dx + \int_{-2}^5 f(x) dx \\ 1 : \int_{-2}^5 f(x) dx \\ 1 : \text{answer} \end{cases}$

(b) $\begin{aligned} \int_3^5 (2f'(x) + 4) dx &= 2 \int_3^5 f'(x) dx + \int_3^5 4 dx \\ &= 2(f(5) - f(3)) + 4(5 - 3) \\ &= 2(0 - (3 - \sqrt{5})) + 8 \\ &= 2(-3 + \sqrt{5}) + 8 = 2 + 2\sqrt{5} \end{aligned}$

2 : $\begin{cases} 1 : \text{Fundamental Theorem of Calculus} \\ 1 : \text{answer} \end{cases}$

— OR —

$$\begin{aligned} \int_3^5 (2f'(x) + 4) dx &= [2f(x) + 4x]_{x=3}^{x=5} \\ &= (2f(5) + 20) - (2f(3) + 12) \\ &= (2 \cdot 0 + 20) - (2(3 - \sqrt{5}) + 12) \\ &= 2 + 2\sqrt{5} \end{aligned}$$

(c) $g'(x) = f(x) = 0 \Rightarrow x = -1, x = \frac{1}{2}, x = 5$

x	$g(x)$
-2	0
-1	$\frac{1}{2}$
$\frac{1}{2}$	$-\frac{1}{4}$
5	$11 - \frac{9\pi}{4}$

3 : $\begin{cases} 1 : g'(x) = f(x) \\ 1 : \text{identifies } x = -1 \text{ as a candidate} \\ 1 : \text{answer with justification} \end{cases}$

On the interval $-2 \leq x \leq 5$, the absolute maximum value of g is $g(5) = 11 - \frac{9\pi}{4}$.

(d) $\begin{aligned} \lim_{x \rightarrow 1} \frac{10^x - 3f'(x)}{f(x) - \arctan x} &= \frac{10^1 - 3f'(1)}{f(1) - \arctan 1} \\ &= \frac{10 - 3 \cdot 2}{1 - \arctan 1} = \frac{4}{1 - \frac{\pi}{4}} \end{aligned}$

1 : answer

2019 SCORING GUIDELINES

Question 4

(a) $V = \pi r^2 h = \pi(1)^2 h = \pi h$

$$\left. \frac{dV}{dt} \right|_{h=4} = \pi \left. \frac{dh}{dt} \right|_{h=4} = \pi \left(-\frac{1}{10} \sqrt{4} \right) = -\frac{\pi}{5} \text{ cubic feet per second}$$

2 : $\begin{cases} 1 : \frac{dV}{dt} = \pi \frac{dh}{dt} \\ 1 : \text{answer with units} \end{cases}$

(b) $\frac{d^2 h}{dt^2} = -\frac{1}{20\sqrt{h}} \cdot \frac{dh}{dt} = -\frac{1}{20\sqrt{h}} \cdot \left(-\frac{1}{10} \sqrt{h} \right) = \frac{1}{200}$

Because $\frac{d^2 h}{dt^2} = \frac{1}{200} > 0$ for $h > 0$, the rate of change of the height is increasing when the height of the water is 3 feet.

3 : $\begin{cases} 1 : \frac{d}{dh} \left(-\frac{1}{10} \sqrt{h} \right) = -\frac{1}{20\sqrt{h}} \\ 1 : \frac{d^2 h}{dt^2} = -\frac{1}{20\sqrt{h}} \cdot \frac{dh}{dt} \\ 1 : \text{answer with explanation} \end{cases}$

(c) $\frac{dh}{\sqrt{h}} = -\frac{1}{10} dt$

$$\int \frac{dh}{\sqrt{h}} = \int -\frac{1}{10} dt$$

$$2\sqrt{h} = -\frac{1}{10}t + C$$

$$2\sqrt{5} = -\frac{1}{10} \cdot 0 + C \Rightarrow C = 2\sqrt{5}$$

$$2\sqrt{h} = -\frac{1}{10}t + 2\sqrt{5}$$

$$h(t) = \left(-\frac{1}{20}t + \sqrt{5} \right)^2$$

4 : $\begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ \text{and uses initial condition} \\ 1 : h(t) \end{cases}$

Note: 0/4 if no separation of variables

Note: max 2/4 [1-1-0-0] if no constant of integration

2019 SCORING GUIDELINES

Question 5

(a) $f'(x) = \frac{-(2x-2)}{(x^2-2x+k)^2}$

$$f'(0) = \frac{2}{k^2} = 6 \Rightarrow k^2 = \frac{1}{3} \Rightarrow k = \frac{1}{\sqrt{3}}$$

3 : $\begin{cases} 1 : \text{denominator of } f'(x) \\ 1 : f'(x) \\ 1 : \text{answer} \end{cases}$

(b) $\frac{1}{x^2-2x-8} = \frac{1}{(x-4)(x+2)} = \frac{A}{x-4} + \frac{B}{x+2}$

$$\Rightarrow 1 = A(x+2) + B(x-4)$$

$$\Rightarrow A = \frac{1}{6}, B = -\frac{1}{6}$$

3 : $\begin{cases} 1 : \text{partial fraction} \\ \text{decomposition} \\ 1 : \text{antiderivatives} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \int_0^1 f(x) dx &= \int_0^1 \left(\frac{1}{6} \frac{1}{x-4} - \frac{1}{6} \frac{1}{x+2} \right) dx \\ &= \left[\frac{1}{6} \ln|x-4| - \frac{1}{6} \ln|x+2| \right]_{x=0}^{x=1} \\ &= \left(\frac{1}{6} \ln 3 - \frac{1}{6} \ln 3 \right) - \left(\frac{1}{6} \ln 4 - \frac{1}{6} \ln 2 \right) = -\frac{1}{6} \ln 2 \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad \int_0^2 \frac{1}{x^2-2x+1} dx &= \int_0^2 \frac{1}{(x-1)^2} dx = \int_0^1 \frac{1}{(x-1)^2} dx + \int_1^2 \frac{1}{(x-1)^2} dx \\ &= \lim_{b \rightarrow 1^-} \int_0^b \frac{1}{(x-1)^2} dx + \lim_{b \rightarrow 1^+} \int_b^2 \frac{1}{(x-1)^2} dx \\ &= \lim_{b \rightarrow 1^-} \left(-\frac{1}{x-1} \Big|_{x=0}^{x=b} \right) + \lim_{b \rightarrow 1^+} \left(-\frac{1}{x-1} \Big|_{x=b}^{x=2} \right) \\ &= \lim_{b \rightarrow 1^-} \left(-\frac{1}{b-1} - 1 \right) + \lim_{b \rightarrow 1^+} \left(-1 + \frac{1}{b-1} \right) \end{aligned}$$

Because $\lim_{b \rightarrow 1^-} \left(-\frac{1}{b-1} \right)$ does not exist, the integral diverges.

3 : $\begin{cases} 1 : \text{improper integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer with reason} \end{cases}$

2019 SCORING GUIDELINES

Question 6

- (a)
- $f(0) = 3$
- and
- $f'(0) = -2$

The third-degree Taylor polynomial for f about $x = 0$ is

$$3 - 2x + \frac{3}{2!}x^2 + \frac{-2}{3!}x^3 = 3 - 2x + \frac{3}{2}x^2 - \frac{23}{12}x^3.$$

- (b) The first three nonzero terms of the Maclaurin series for
- e^x
- are

$$1 + x + \frac{1}{2!}x^2.$$

The second-degree Taylor polynomial for $e^x f(x)$ about $x = 0$ is

$$\begin{aligned} 3\left(1 + x + \frac{1}{2!}x^2\right) - 2x(1 + x) + \frac{3}{2}x^2(1) \\ = 3 + (3 - 2)x + \left(\frac{3}{2} - 2 + \frac{3}{2}\right)x^2 \\ = 3 + x + x^2. \end{aligned}$$

- (c)
- $h(1) = \int_0^1 f(t) dt$

$$\begin{aligned} &\approx \int_0^1 \left(3 - 2t + \frac{3}{2}t^2 - \frac{23}{12}t^3\right) dt \\ &= \left[3t - t^2 + \frac{1}{2}t^3 - \frac{23}{48}t^4\right]_{t=0}^{t=1} \\ &= 3 - 1 + \frac{1}{2} - \frac{23}{48} = \frac{97}{48} \end{aligned}$$

- (d) The alternating series error bound is the absolute value of the first omitted term of the series for
- $h(1)$
- .

$$\int_0^1 \left(\frac{54}{4!}t^4\right) dt = \left[\frac{9}{20}t^5\right]_{t=0}^{t=1} = \frac{9}{20}$$

$$\text{Error} \leq \left|\frac{9}{20}\right| = 0.45$$

$$2 : \begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \end{cases}$$

$$2 : \begin{cases} 1 : \text{three terms for } e^x \\ 1 : \text{three terms for } e^x f(x) \end{cases}$$

$$2 : \begin{cases} 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : \text{uses fourth-degree term} \\ \quad \text{of Maclaurin series for } f \\ 1 : \text{uses first omitted term} \\ \quad \text{of series for } h(1) \\ 1 : \text{error bound} \end{cases}$$

2018

AP[®]

CollegeBoard

AP Calculus BC

Scoring Guidelines

2018 SCORING GUIDELINES

Question 1

(a) $\int_0^{300} r(t) dt = 270$

According to the model, 270 people enter the line for the escalator during the time interval $0 \leq t \leq 300$.

(b) $20 + \int_0^{300} (r(t) - 0.7) dt = 20 + \int_0^{300} r(t) dt - 0.7 \cdot 300 = 80$

According to the model, 80 people are in line at time $t = 300$.

(c) Based on part (b), the number of people in line at time $t = 300$ is 80.

The first time t that there are no people in line is

$$300 + \frac{80}{0.7} = 414.286 \text{ (or } 414.285 \text{) seconds.}$$

(d) The total number of people in line at time t , $0 \leq t \leq 300$, is modeled by

$$20 + \int_0^t r(x) dx - 0.7t.$$

$$r(t) - 0.7 = 0 \Rightarrow t_1 = 33.013298, t_2 = 166.574719$$

t	People in line for escalator
0	20
t_1	3.803
t_2	158.070
300	80

The number of people in line is a minimum at time $t = 33.013$ seconds, when there are 4 people in line.

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

$$2 : \begin{cases} 1 : \text{considers rate out} \\ 1 : \text{answer} \end{cases}$$

$$1 : \text{answer}$$

$$4 : \begin{cases} 1 : \text{considers } r(t) - 0.7 = 0 \\ 1 : \text{identifies } t = 33.013 \\ 1 : \text{answers} \\ 1 : \text{justification} \end{cases}$$

2018 SCORING GUIDELINES

Question 2

(a) $p'(25) = -1.179$

At a depth of 25 meters, the density of plankton cells is changing at a rate of -1.179 million cells per cubic meter per meter.

(b) $\int_0^{30} 3p(h) dh = 1675.414936$

There are 1675 million plankton cells in the column of water between $h = 0$ and $h = 30$ meters.

(c) $\int_{30}^K 3f(h) dh$ represents the number of plankton cells, in millions, in the column of water from a depth of 30 meters to a depth of K meters.

The number of plankton cells, in millions, in the entire column of water is given by $\int_0^{30} 3p(h) dh + \int_{30}^K 3f(h) dh$.

Because $0 \leq f(h) \leq u(h)$ for all $h \geq 30$,

$$3 \int_{30}^K f(h) dh \leq 3 \int_{30}^K u(h) dh \leq 3 \int_{30}^{\infty} u(h) dh = 3 \cdot 105 = 315.$$

The total number of plankton cells in the column of water is bounded by $1675.415 + 315 = 1990.415 \leq 2000$ million.

(d) $\int_0^1 \sqrt{(x'(t))^2 + (y'(t))^2} dt = 757.455862$

The total distance traveled by the boat over the time interval $0 \leq t \leq 1$ is 757.456 (or 757.455) meters.

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{meaning with units} \end{cases}$$

$$2 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 1 : \text{integral expression} \\ 1 : \text{compares improper integral} \\ 1 : \text{explanation} \end{cases}$$

$$2 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{total distance} \end{cases}$$

2018 SCORING GUIDELINES

Question 3

$$\begin{aligned} \text{(a)} \quad f(-5) &= f(1) + \int_1^{-5} g(x) \, dx = f(1) - \int_{-5}^1 g(x) \, dx \\ &= 3 - \left(-9 - \frac{3}{2} + 1\right) = 3 - \left(-\frac{19}{2}\right) = \frac{25}{2} \end{aligned}$$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b)} \quad \int_1^6 g(x) \, dx &= \int_1^3 g(x) \, dx + \int_3^6 g(x) \, dx \\ &= \int_1^3 2 \, dx + \int_3^6 2(x-4)^2 \, dx \\ &= 4 + \left[\frac{2}{3}(x-4)^3\right]_{x=3}^{x=6} = 4 + \frac{16}{3} - \left(-\frac{2}{3}\right) = 10 \end{aligned}$$

3 : $\begin{cases} 1 : \text{split at } x = 3 \\ 1 : \text{antiderivative of } 2(x-4)^2 \\ 1 : \text{answer} \end{cases}$

(c) The graph of f is increasing and concave up on $0 < x < 1$ and $4 < x < 6$ because $f'(x) = g(x) > 0$ and $f'(x) = g(x)$ is increasing on those intervals.

2 : $\begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$

(d) The graph of f has a point of inflection at $x = 4$ because $f'(x) = g(x)$ changes from decreasing to increasing at $x = 4$.

2 : $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

2018 SCORING GUIDELINES

Question 4

$$\text{(a)} \quad H'(6) \approx \frac{H(7) - H(5)}{7 - 5} = \frac{11 - 6}{2} = \frac{5}{2}$$

$H'(6)$ is the rate at which the height of the tree is changing, in meters per year, at time $t = 6$ years.

$$\text{(b)} \quad \frac{H(5) - H(3)}{5 - 3} = \frac{6 - 2}{2} = 2$$

Because H is differentiable on $3 \leq t \leq 5$, H is continuous on $3 \leq t \leq 5$.

By the Mean Value Theorem, there exists a value c , $3 < c < 5$, such that $H'(c) = 2$.

(c) The average height of the tree over the time interval $2 \leq t \leq 10$ is given by $\frac{1}{10-2} \int_2^{10} H(t) \, dt$.

$$\begin{aligned} \frac{1}{8} \int_2^{10} H(t) \, dt &\approx \frac{1}{8} \left(\frac{1.5+2}{2} \cdot 1 + \frac{2+6}{2} \cdot 2 + \frac{6+11}{2} \cdot 2 + \frac{11+15}{2} \cdot 3 \right) \\ &= \frac{1}{8} (65.75) = \frac{263}{32} \end{aligned}$$

The average height of the tree over the time interval $2 \leq t \leq 10$ is $\frac{263}{32}$ meters.

$$\text{(d)} \quad G(x) = 50 \Rightarrow x = 1$$

$$\frac{d}{dx}(G(x)) = \frac{d}{dx}(G(x)) \cdot \frac{dx}{dt} = \frac{(1+x)100 - 100x \cdot 1}{(1+x)^2} \cdot \frac{dx}{dt} = \frac{100}{(1+x)^2} \cdot \frac{dx}{dt}$$

$$\left. \frac{d}{dx}(G(x)) \right|_{x=1} = \frac{100}{(1+1)^2} \cdot 0.03 = \frac{3}{4}$$

According to the model, the rate of change of the height of the tree with respect to time when the tree is 50 meters tall is $\frac{3}{4}$ meter per year.

2 : $\begin{cases} 1 : \text{estimate} \\ 1 : \text{interpretation with units} \end{cases}$

2 : $\begin{cases} 1 : \frac{H(5) - H(3)}{5 - 3} \\ 1 : \text{conclusion using Mean Value Theorem} \end{cases}$

2 : $\begin{cases} 1 : \text{trapezoidal sum} \\ 1 : \text{approximation} \end{cases}$

3 : $\begin{cases} 2 : \frac{d}{dt}(G(x)) \\ 1 : \text{answer} \end{cases}$

Note: max 1/3 [1-0] if no chain rule

2018 SCORING GUIDELINES

Question 5

$$(a) \text{ Area} = \frac{1}{2} \int_{\pi/3}^{5\pi/3} (4^2 - (3 + 2\cos\theta)^2) d\theta$$

$$(b) \frac{dr}{d\theta} = -2\sin\theta \Rightarrow \left. \frac{dr}{d\theta} \right|_{\theta=\pi/2} = -2$$

$$r\left(\frac{\pi}{2}\right) = 3 + 2\cos\left(\frac{\pi}{2}\right) = 3$$

$$y = r\sin\theta \Rightarrow \frac{dy}{d\theta} = \frac{dr}{d\theta}\sin\theta + r\cos\theta$$

$$x = r\cos\theta \Rightarrow \frac{dx}{d\theta} = \frac{dr}{d\theta}\cos\theta - r\sin\theta$$

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/2} = \left. \frac{dy/d\theta}{dx/d\theta} \right|_{\theta=\pi/2} = \frac{-2\sin\left(\frac{\pi}{2}\right) + 3\cos\left(\frac{\pi}{2}\right)}{-2\cos\left(\frac{\pi}{2}\right) - 3\sin\left(\frac{\pi}{2}\right)} = \frac{2}{3}$$

The slope of the line tangent to the graph of $r = 3 + 2\cos\theta$

at $\theta = \frac{\pi}{2}$ is $\frac{2}{3}$.

— OR —

$$y = r\sin\theta = (3 + 2\cos\theta)\sin\theta \Rightarrow \frac{dy}{d\theta} = 3\cos\theta + 2\cos^2\theta - 2\sin^2\theta$$

$$x = r\cos\theta = (3 + 2\cos\theta)\cos\theta \Rightarrow \frac{dx}{d\theta} = -3\sin\theta - 4\sin\theta\cos\theta$$

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/2} = \left. \frac{dy/d\theta}{dx/d\theta} \right|_{\theta=\pi/2} = \frac{3\cos\left(\frac{\pi}{2}\right) + 2\cos^2\left(\frac{\pi}{2}\right) - 2\sin^2\left(\frac{\pi}{2}\right)}{-3\sin\left(\frac{\pi}{2}\right) - 4\sin\left(\frac{\pi}{2}\right)\cos\left(\frac{\pi}{2}\right)} = \frac{2}{3}$$

The slope of the line tangent to the graph of $r = 3 + 2\cos\theta$

at $\theta = \frac{\pi}{2}$ is $\frac{2}{3}$.

$$(c) \frac{dr}{dt} = \frac{dr}{d\theta} \cdot \frac{d\theta}{dt} = -2\sin\theta \cdot \frac{d\theta}{dt} \Rightarrow \frac{d\theta}{dt} = \frac{dr}{dt} \cdot \frac{1}{-2\sin\theta}$$

$$\left. \frac{d\theta}{dt} \right|_{\theta=\pi/3} = 3 \cdot \frac{1}{-2\sin\left(\frac{\pi}{3}\right)} = \frac{3}{-\sqrt{3}} = -\sqrt{3} \text{ radians per second}$$

3 : $\begin{cases} 1 : \text{constant and limits} \\ 2 : \text{integrand} \end{cases}$

3 : $\begin{cases} 1 : \frac{dy}{d\theta} = \frac{dr}{d\theta}\sin\theta + r\cos\theta \\ \text{or } \frac{dx}{d\theta} = \frac{dr}{d\theta}\cos\theta - r\sin\theta \\ 1 : \frac{dy}{dx} = \frac{\frac{dr}{d\theta}\sin\theta + r\cos\theta}{\frac{dr}{d\theta}\cos\theta - r\sin\theta} \\ 1 : \text{answer} \end{cases}$

2018 SCORING GUIDELINES

Question 6

$$(a) \text{ The first four nonzero terms are } \frac{x^2}{3} - \frac{x^3}{2 \cdot 3^2} + \frac{x^4}{3 \cdot 3^3} - \frac{x^5}{4 \cdot 3^4}.$$

$$\text{The general term is } (-1)^{n+1} \frac{x^{n+1}}{n \cdot 3^n}.$$

$$(b) \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+2} x^{n+2}}{(n+1)(3^{n+1})} \cdot \frac{(n \cdot 3^n)}{(-1)^{n+1} x^{n+1}} \right| = \lim_{n \rightarrow \infty} \left| \frac{-x}{3} \cdot \frac{n}{(n+1)} \right| = \left| \frac{x}{3} \right|$$

$$\left| \frac{x}{3} \right| < 1 \text{ for } |x| < 3$$

Therefore, the radius of convergence of the Maclaurin series for f is 3.

— OR —

The radius of convergence of the Maclaurin series for $\ln(1+x)$ is 1,

so the series for $f(x) = x \ln\left(1 + \frac{x}{3}\right)$ converges absolutely for $\left|\frac{x}{3}\right| < 1$.

$$\left| \frac{x}{3} \right| < 1 \Rightarrow |x| < 3$$

Therefore, the radius of convergence of the Maclaurin series for f is 3.

When $x = -3$, the series is $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{(-3)^{n+1}}{n \cdot 3^n} = \sum_{n=1}^{\infty} \frac{3}{n}$, which diverges by comparison to the harmonic series.

When $x = 3$, the series is $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{3^{n+1}}{n \cdot 3^n} = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{3}{n}$, which converges by the alternating series test.

The interval of convergence of the Maclaurin series for f is $-3 < x \leq 3$.

(c) By the alternating series error bound, an upper bound for $|P_4(2) - f(2)|$ is the magnitude of the next term of the alternating series.

$$|P_4(2) - f(2)| < \left| -\frac{2^5}{4 \cdot 3^4} \right| = \frac{8}{81}$$

2 : $\begin{cases} 1 : \text{first four terms} \\ 1 : \text{general term} \end{cases}$

5 : $\begin{cases} 1 : \text{sets up ratio} \\ 1 : \text{computes limit of ratio} \\ 1 : \text{radius of convergence} \\ 1 : \text{considers both endpoints} \\ 1 : \text{analysis and interval of convergence} \end{cases}$

— OR —

5 : $\begin{cases} 1 : \text{radius for } \ln(1+x) \text{ series} \\ 1 : \text{substitutes } \frac{x}{3} \\ 1 : \text{radius of convergence} \\ 1 : \text{considers both endpoints} \\ 1 : \text{analysis and interval of convergence} \end{cases}$

2 : $\begin{cases} 1 : \text{uses fifth-degree term as error bound} \\ 1 : \text{answer} \end{cases}$

2017

AP Calculus BC

Scoring Guidelines

2017 SCORING GUIDELINES

Question 1

	1 : units in parts (a), (c), and (d)
(a) $\text{Volume} = \int_0^{10} A(h) \, dh$ $\approx (2 - 0) \cdot A(0) + (5 - 2) \cdot A(2) + (10 - 5) \cdot A(5)$ $= 2 \cdot 50.3 + 3 \cdot 14.4 + 5 \cdot 6.5$ $= 176.3 \text{ cubic feet}$	2 : $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{approximation} \end{cases}$
(b) The approximation in part (a) is an overestimate because a left Riemann sum is used and A is decreasing.	1 : overestimate with reason
(c) $\int_0^{10} f(h) \, dh = 101.325338$ The volume is 101.325 cubic feet.	2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$
(d) Using the model, $V(h) = \int_0^h f(x) \, dx$. $\left. \frac{dV}{dt} \right _{h=5} = \left[\frac{dV}{dh} \cdot \frac{dh}{dt} \right]_{h=5}$ $= \left[f(h) \cdot \frac{dh}{dt} \right]_{h=5}$ $= f(5) \cdot 0.26 = 1.694419$ When $h = 5$, the volume of water is changing at a rate of 1.694 cubic feet per minute.	3 : $\begin{cases} 2 : \frac{dV}{dt} \\ 1 : \text{answer} \end{cases}$

2017 SCORING GUIDELINES

Question 2

$$(a) \frac{1}{2} \int_0^{\pi/2} (f(\theta))^2 d\theta = 0.648414$$

The area of R is 0.648.

$$(b) \int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \frac{1}{2} \int_0^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

— OR —

$$\int_0^k ((g(\theta))^2 - (f(\theta))^2) d\theta = \int_k^{\pi/2} ((g(\theta))^2 - (f(\theta))^2) d\theta$$

$$(c) w(\theta) = g(\theta) - f(\theta)$$

$$w_A = \frac{\int_0^{\pi/2} w(\theta) d\theta}{\frac{\pi}{2} - 0} = 0.485446$$

The average value of $w(\theta)$ on the interval $\left[0, \frac{\pi}{2}\right]$ is 0.485.

$$(d) w(\theta) = w_A \text{ for } 0 \leq \theta \leq \frac{\pi}{2} \Rightarrow \theta = 0.517688$$

$$w(\theta) = w_A \text{ at } \theta = 0.518 \text{ (or } 0.517).$$

$$w'(0.518) < 0 \Rightarrow w(\theta) \text{ is decreasing at } \theta = 0.518.$$

$$2: \begin{cases} 1: \text{integral} \\ 1: \text{answer} \end{cases}$$

$$2: \begin{cases} 1: \text{integral expression} \\ \quad \text{for one region} \\ 1: \text{equation} \end{cases}$$

$$3: \begin{cases} 1: w(\theta) \\ 1: \text{integral} \\ 1: \text{average value} \end{cases}$$

$$2: \begin{cases} 1: \text{solves } w(\theta) = w_A \\ 1: \text{answer with reason} \end{cases}$$

2017 SCORING GUIDELINES

Question 3

$$(a) f(-6) = f(-2) + \int_{-2}^{-6} f'(x) dx = 7 - \int_{-6}^{-2} f'(x) dx = 7 - 4 = 3$$

$$f(5) = f(-2) + \int_{-2}^5 f'(x) dx = 7 - 2\pi + 3 = 10 - 2\pi$$

$$(b) f'(x) > 0 \text{ on the intervals } [-6, -2) \text{ and } (2, 5).$$

Therefore, f is increasing on the intervals $[-6, -2]$ and $[2, 5]$.

$$(c) \text{The absolute minimum will occur at a critical point where } f'(x) = 0 \text{ or at an endpoint.}$$

$$f'(x) = 0 \Rightarrow x = -2, x = 2$$

x	$f(x)$
-6	3
-2	7
2	$7 - 2\pi$
5	$10 - 2\pi$

The absolute minimum value is $f(2) = 7 - 2\pi$.

$$(d) f''(-5) = \frac{2 - 0}{-6 - (-2)} = -\frac{1}{2}$$

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} = 2 \text{ and } \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3} = -1$$

$f''(3)$ does not exist because

$$\lim_{x \rightarrow 3^-} \frac{f'(x) - f'(3)}{x - 3} \neq \lim_{x \rightarrow 3^+} \frac{f'(x) - f'(3)}{x - 3}.$$

$$3: \begin{cases} 1: \text{uses initial condition} \\ 1: f(-6) \\ 1: f(5) \end{cases}$$

$$2: \text{answer with justification}$$

$$2: \begin{cases} 1: \text{considers } x = 2 \\ 1: \text{answer with justification} \end{cases}$$

$$2: \begin{cases} 1: f''(-5) \\ 1: f''(3) \text{ does not exist,} \\ \quad \text{with explanation} \end{cases}$$

2017 SCORING GUIDELINES

Question 4

(a) $H'(0) = -\frac{1}{4}(91 - 27) = -16$
 $H(0) = 91$

An equation for the tangent line is $y = 91 - 16t$.

The internal temperature of the potato at time $t = 3$ minutes is approximately $91 - 16 \cdot 3 = 43$ degrees Celsius.

(b) $\frac{d^2H}{dt^2} = -\frac{1}{4} \frac{dH}{dt} = \left(-\frac{1}{4}\right)\left(-\frac{1}{4}\right)(H - 27) = \frac{1}{16}(H - 27)$

$$H > 27 \text{ for } t > 0 \Rightarrow \frac{d^2H}{dt^2} = \frac{1}{16}(H - 27) > 0 \text{ for } t > 0$$

Therefore, the graph of H is concave up for $t > 0$. Thus, the answer in part (a) is an underestimate.

(c) $\frac{dG}{(G - 27)^{2/3}} = -dt$
 $\int \frac{dG}{(G - 27)^{2/3}} = \int (-1) dt$
 $3(G - 27)^{1/3} = -t + C$
 $3(91 - 27)^{1/3} = 0 + C \Rightarrow C = 12$
 $3(G - 27)^{1/3} = 12 - t$
 $G(t) = 27 + \left(\frac{12 - t}{3}\right)^3 \text{ for } 0 \leq t < 10$

The internal temperature of the potato at time $t = 3$ minutes is $27 + \left(\frac{12 - 3}{3}\right)^3 = 54$ degrees Celsius.

3 : $\begin{cases} 1 : \text{slope} \\ 1 : \text{tangent line} \\ 1 : \text{approximation} \end{cases}$

1 : underestimate with reason

5 : $\begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration and uses initial condition} \\ 1 : \text{equation involving } G \text{ and } t \\ 1 : G(t) \text{ and } G(3) \end{cases}$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

2017 SCORING GUIDELINES

Question 5

(a) $f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2}$

$$f'(3) = \frac{(-3)(5)}{(18 - 21 + 5)^2} = -\frac{15}{4}$$

(b) $f'(x) = \frac{-3(4x - 7)}{(2x^2 - 7x + 5)^2} = 0 \Rightarrow x = \frac{7}{4}$

The only critical point in the interval $1 < x < 2.5$ has x -coordinate $\frac{7}{4}$.

f' changes sign from positive to negative at $x = \frac{7}{4}$.

Therefore, f has a relative maximum at $x = \frac{7}{4}$.

(c) $\int_5^\infty f(x) dx = \lim_{b \rightarrow \infty} \int_5^b \frac{3}{2x^2 - 7x + 5} dx = \lim_{b \rightarrow \infty} \int_5^b \left(\frac{2}{2x - 5} - \frac{1}{x - 1} \right) dx$
 $= \lim_{b \rightarrow \infty} \left[\ln(2x - 5) - \ln(x - 1) \right]_5^b = \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2x - 5}{x - 1} \right) \right]_5^b$
 $= \lim_{b \rightarrow \infty} \left[\ln \left(\frac{2b - 5}{b - 1} \right) - \ln \left(\frac{5}{4} \right) \right] = \ln 2 - \ln \left(\frac{5}{4} \right) = \ln \left(\frac{8}{5} \right)$

(d) f is continuous, positive, and decreasing on $[5, \infty)$.

The series converges by the integral test since $\int_5^\infty \frac{3}{2x^2 - 7x + 5} dx$ converges.

— OR —

$$\frac{3}{2n^2 - 7n + 5} > 0 \text{ and } \frac{1}{n^2} > 0 \text{ for } n \geq 5.$$

Since $\lim_{n \rightarrow \infty} \frac{\frac{3}{2n^2 - 7n + 5}}{\frac{1}{n^2}} = \frac{3}{2}$ and the series $\sum_{n=5}^\infty \frac{1}{n^2}$ converges,

the series $\sum_{n=5}^\infty \frac{3}{2n^2 - 7n + 5}$ converges by the limit comparison test.

2 : $f'(3)$

2 : $\begin{cases} 1 : x\text{-coordinate} \\ 1 : \text{relative maximum with justification} \end{cases}$

3 : $\begin{cases} 1 : \text{antiderivative} \\ 1 : \text{limit expression} \\ 1 : \text{answer} \end{cases}$

2 : answer with conditions

2017 SCORING GUIDELINES

Question 6

(a) $f(0) = 0$

$f'(0) = 1$

$f''(0) = -1(1) = -1$

$f'''(0) = -2(-1) = 2$

$f^{(4)}(0) = -3(2) = -6$

The first four nonzero terms are

$$0 + 1x + \frac{-1}{2!}x^2 + \frac{2}{3!}x^3 + \frac{-6}{4!}x^4 = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4}.$$

The general term is $\frac{(-1)^{n+1}x^n}{n}$.

(b) For $x = 1$, the Maclaurin series becomes $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n}$.

The series does not converge absolutely because the harmonic series diverges.

The series alternates with terms that decrease in magnitude to 0, and therefore the series converges conditionally.

(c)
$$\begin{aligned} \int_0^x f(t) dt &= \int_0^x \left(t - \frac{t^2}{2} + \frac{t^3}{3} - \frac{t^4}{4} + \cdots + \frac{(-1)^{n+1}t^n}{n} + \cdots \right) dt \\ &= \left[\frac{t^2}{2} - \frac{t^3}{3 \cdot 2} + \frac{t^4}{4 \cdot 3} - \frac{t^5}{5 \cdot 4} + \cdots + \frac{(-1)^{n+1}t^{n+1}}{(n+1)n} + \cdots \right]_{t=0}^{t=x} \\ &= \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{12} - \frac{x^5}{20} + \cdots + \frac{(-1)^{n+1}x^{n+1}}{(n+1)n} + \cdots \end{aligned}$$

(d) The terms alternate in sign and decrease in magnitude to 0. By the alternating series error bound, the error $\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right|$ is boundedby the magnitude of the first unused term, $\left| -\frac{(1/2)^5}{20} \right|$.

Thus, $\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right| \leq \left| -\frac{(1/2)^5}{20} \right| = \frac{1}{32 \cdot 20} < \frac{1}{500}.$

3 : $\begin{cases} 1 : f''(0), f'''(0), \text{ and } f^{(4)}(0) \\ 1 : \text{verify terms} \\ 1 : \text{general term} \end{cases}$

2 : converges conditionally
with reason

3 : $\begin{cases} 1 : \text{two terms} \\ 1 : \text{remaining terms} \\ 1 : \text{general term} \end{cases}$

1 : error bound

AP[®] Calculus BC
2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2016 SCORING GUIDELINES

Question 1

t (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.

- (a) Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.
- (b) Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
- (c) Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
- (d) For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

(a) $R'(2) \approx \frac{R(3) - R(1)}{3 - 1} = \frac{950 - 1190}{3 - 1} = -120$ liters/hr²

2 : $\begin{cases} 1 : \text{estimate} \\ 1 : \text{units} \end{cases}$

(b) The total amount of water removed is given by $\int_0^8 R(t) dt$.

$$\begin{aligned} \int_0^8 R(t) dt &\approx 1 \cdot R(0) + 2 \cdot R(1) + 3 \cdot R(3) + 2 \cdot R(6) \\ &= 1(1340) + 2(1190) + 3(950) + 2(740) \\ &= 8050 \text{ liters} \end{aligned}$$

3 : $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{estimate} \\ 1 : \text{overestimate with reason} \end{cases}$

This is an overestimate since R is a decreasing function.

(c) Total $\approx 50000 + \int_0^8 W(t) dt - 8050$
 $= 50000 + 7836.195325 - 8050 \approx 49786$ liters

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{estimate} \end{cases}$

(d) $W(0) - R(0) > 0$, $W(8) - R(8) < 0$, and $W(t) - R(t)$ is continuous.


2 : $\begin{cases} 1 : \text{considers } W(t) - R(t) \\ 1 : \text{answer with explanation} \end{cases}$

Therefore, the Intermediate Value Theorem guarantees at least one time t , $0 < t < 8$, for which $W(t) - R(t) = 0$, or $W(t) = R(t)$.

For this value of t , the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank.

2016 SCORING GUIDELINES

Question 2



At time t , the position of a particle moving in the xy -plane is given by the parametric functions $(x(t), y(t))$, where $\frac{dx}{dt} = t^2 + \sin(3t^2)$. The graph of y , consisting of three line segments, is shown in the figure above. At $t = 0$, the particle is at position $(5, 1)$.

- (a) Find the position of the particle at $t = 3$.
- (b) Find the slope of the line tangent to the path of the particle at $t = 3$.
- (c) Find the speed of the particle at $t = 3$.
- (d) Find the total distance traveled by the particle from $t = 0$ to $t = 2$.

(a) $x(3) = x(0) + \int_0^3 x'(t) dt = 5 + 9.377035 = 14.377$

$$y(3) = -\frac{1}{2}$$

The position of the particle at $t = 3$ is $(14.377, -0.5)$.

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

(b) Slope $= \frac{y'(3)}{x'(3)} = \frac{0.5}{9.956376} = 0.05$

1 : slope

(c) Speed $= \sqrt{(x'(3))^2 + (y'(3))^2} = 9.969$ (or 9.968)

2 : $\begin{cases} 1 : \text{expression for speed} \\ 1 : \text{answer} \end{cases}$

(d) Distance $= \int_0^2 \sqrt{(x'(t))^2 + (y'(t))^2} dt$
 $= \int_0^1 \sqrt{(x'(t))^2 + (-2)^2} dt + \int_1^2 \sqrt{(x'(t))^2 + 0^2} dt$
 $= 2.237871 + 2.112003 = 4.350$ (or 4.349)

3 : $\begin{cases} 1 : \text{expression for distance} \\ 1 : \text{integrals} \\ 1 : \text{answer} \end{cases}$


2016 SCORING GUIDELINES

Question 3

The figure above shows the graph of the piecewise-linear function f . For $-4 \leq x \leq 12$, the function g is defined by

$$g(x) = \int_2^x f(t) dt.$$

- (a) Does g have a relative minimum, a relative maximum, or neither at $x = 10$? Justify your answer.
- (b) Does the graph of g have a point of inflection at $x = 4$? Justify your answer.
- (c) Find the absolute minimum value and the absolute maximum value of g on the interval $-4 \leq x \leq 12$. Justify your answers.
- (d) For $-4 \leq x \leq 12$, find all intervals for which $g(x) \leq 0$.

Graph of f

- (a) The function g has neither a relative minimum nor a relative maximum at $x = 10$ since $g'(x) = f(x)$ and $f(x) \leq 0$ for $8 \leq x \leq 12$.
- (b) The graph of g has a point of inflection at $x = 4$ since $g'(x) = f(x)$ is increasing for $2 \leq x \leq 4$ and decreasing for $4 \leq x \leq 8$.
- (c) $g'(x) = f(x)$ changes sign only at $x = -2$ and $x = 6$.

x	$g(x)$
-4	-4
-2	-8
6	8
12	-4

On the interval $-4 \leq x \leq 12$, the absolute minimum value is $g(-2) = -8$ and the absolute maximum value is $g(6) = 8$.

- (d) $g(x) \leq 0$ for $-4 \leq x \leq 2$ and $10 \leq x \leq 12$.

1 : $g'(x) = f(x)$ in (a), (b), (c), or (d)

1 : answer with justification

1 : answer with justification

4 : $\begin{cases} 1 : \text{considers } x = -2 \text{ and } x = 6 \\ \text{as candidates} \\ 1 : \text{considers } x = -4 \text{ and } x = 12 \\ 2 : \text{answers with justification} \end{cases}$

2 : intervals

2016 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = x^2 - \frac{1}{2}y$.

- (a) Find $\frac{d^2y}{dx^2}$ in terms of x and y .
- (b) Let $y = f(x)$ be the particular solution to the given differential equation whose graph passes through the point $(-2, 8)$. Does the graph of f have a relative minimum, a relative maximum, or neither at the point $(-2, 8)$? Justify your answer.
- (c) Let $y = g(x)$ be the particular solution to the given differential equation with $g(-1) = 2$. Find $\lim_{x \rightarrow -1} \left(\frac{g(x) - 2}{3(x+1)^2} \right)$. Show the work that leads to your answer.
- (d) Let $y = h(x)$ be the particular solution to the given differential equation with $h(0) = 2$. Use Euler's method, starting at $x = 0$ with two steps of equal size, to approximate $h(1)$.

(a) $\frac{d^2y}{dx^2} = 2x - \frac{1}{2} \frac{dy}{dx} = 2x - \frac{1}{2} \left(x^2 - \frac{1}{2}y \right)$

2 : $\frac{d^2y}{dx^2}$ in terms of x and y

(b) $\left. \frac{dy}{dx} \right|_{(x,y)=(-2,8)} = (-2)^2 - \frac{1}{2} \cdot 8 = 0$
 $\left. \frac{d^2y}{dx^2} \right|_{(x,y)=(-2,8)} = 2(-2) - \frac{1}{2} \left((-2)^2 - \frac{1}{2} \cdot 8 \right) = -4 < 0$

Thus, the graph of f has a relative maximum at the point $(-2, 8)$.

2 : conclusion with justification

(c) $\lim_{x \rightarrow -1} (g(x) - 2) = 0$ and $\lim_{x \rightarrow -1} 3(x+1)^2 = 0$

Using L'Hospital's Rule,

$$\lim_{x \rightarrow -1} \left(\frac{g(x) - 2}{3(x+1)^2} \right) = \lim_{x \rightarrow -1} \left(\frac{g'(x)}{6(x+1)} \right)$$

$$\lim_{x \rightarrow -1} g'(x) = 0 \text{ and } \lim_{x \rightarrow -1} 6(x+1) = 0$$

Using L'Hospital's Rule,

$$\lim_{x \rightarrow -1} \left(\frac{g'(x)}{6(x+1)} \right) = \lim_{x \rightarrow -1} \left(\frac{g''(x)}{6} \right) = \frac{-2}{6} = -\frac{1}{3}$$


3 : $\begin{cases} 2 : \text{L'Hospital's Rule} \\ 1 : \text{answer} \end{cases}$

(d) $h\left(\frac{1}{2}\right) \approx h(0) + h'(0) \cdot \frac{1}{2} = 2 + (-1) \cdot \frac{1}{2} = \frac{3}{2}$
 $h(1) \approx h\left(\frac{1}{2}\right) + h'\left(\frac{1}{2}\right) \cdot \frac{1}{2} \approx \frac{3}{2} + \left(-\frac{1}{2}\right) \cdot \frac{1}{2} = \frac{5}{4}$

2 : $\begin{cases} 1 : \text{Euler's method} \\ 1 : \text{approximation} \end{cases}$

2016 SCORING GUIDELINES

Question 5



The inside of a funnel of height 10 inches has circular cross sections, as shown in the figure above. At height h , the radius of the funnel is given by $r = \frac{1}{20}(3 + h^2)$, where $0 \leq h \leq 10$. The units of r and h are inches.

- Find the average value of the radius of the funnel.
- Find the volume of the funnel.
- The funnel contains liquid that is draining from the bottom. At the instant when the height of the liquid is $h = 3$ inches, the radius of the surface of the liquid is decreasing at a rate of $\frac{1}{5}$ inch per second. At this instant, what is the rate of change of the height of the liquid with respect to time?

$$\begin{aligned} \text{(a) Average radius} &= \frac{1}{10} \int_0^{10} \frac{1}{20}(3 + h^2) dh = \frac{1}{200} \left[3h + \frac{h^3}{3} \right]_0^{10} \\ &= \frac{1}{200} \left(\left(30 + \frac{1000}{3} \right) - 0 \right) = \frac{109}{60} \text{ in} \end{aligned}$$

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b) Volume} &= \pi \int_0^{10} \left(\frac{1}{20}(3 + h^2) \right)^2 dh = \frac{\pi}{400} \int_0^{10} (9 + 6h^2 + h^4) dh \\ &= \frac{\pi}{400} \left[9h + 2h^3 + \frac{h^5}{5} \right]_0^{10} \\ &= \frac{\pi}{400} \left(\left(90 + 2000 + \frac{100000}{5} \right) - 0 \right) = \frac{2209\pi}{40} \text{ in}^3 \end{aligned}$$

3 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(c) } \frac{dr}{dt} &= \frac{1}{20}(2h) \frac{dh}{dt} \\ -\frac{1}{5} &= \frac{3}{10} \frac{dh}{dt} \\ \frac{dh}{dt} &= -\frac{1}{5} \cdot \frac{10}{3} = -\frac{2}{3} \text{ in/sec} \end{aligned}$$

3 : $\begin{cases} 2 : \text{chain rule} \\ 1 : \text{answer} \end{cases}$

2016 SCORING GUIDELINES

Question 6

The function f has a Taylor series about $x = 1$ that converges to $f(x)$ for all x in the interval of convergence.

It is known that $f(1) = 1$, $f'(1) = -\frac{1}{2}$, and the n th derivative of f at $x = 1$ is given by

$$f^{(n)}(1) = (-1)^n \frac{(n-1)!}{2^n} \text{ for } n \geq 2.$$

- Write the first four nonzero terms and the general term of the Taylor series for f about $x = 1$.
- The Taylor series for f about $x = 1$ has a radius of convergence of 2. Find the interval of convergence. Show the work that leads to your answer.
- The Taylor series for f about $x = 1$ can be used to represent $f(1.2)$ as an alternating series. Use the first three nonzero terms of the alternating series to approximate $f(1.2)$.
- Show that the approximation found in part (c) is within 0.001 of the exact value of $f(1.2)$.

$$\begin{aligned} \text{(a) } f(1) &= 1, \quad f'(1) = -\frac{1}{2}, \quad f''(1) = \frac{1}{2^2}, \quad f'''(1) = -\frac{2}{2^3} \\ f(x) &= 1 - \frac{1}{2}(x-1) + \frac{1}{2^2} \cdot \frac{1}{2}(x-1)^2 - \frac{1}{2^3} \cdot \frac{1}{3}(x-1)^3 + \cdots \\ &\quad + \frac{(-1)^n}{2^n \cdot n}(x-1)^n + \cdots \end{aligned}$$

4 : $\begin{cases} 1 : \text{first two terms} \\ 1 : \text{third term} \\ 1 : \text{fourth term} \\ 1 : \text{general term} \end{cases}$

- $R = 2$. The series converges on the interval $(-1, 3)$.

When $x = -1$, the series is $1 + 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots$.

Since the harmonic series diverges, this series diverges.

When $x = 3$, the series is $1 - 1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \cdots$.

Since the alternating harmonic series converges, this series converges.

Therefore, the interval of convergence is $-1 < x \leq 3$.

$$\text{(c) } f(1.2) \approx 1 - \frac{1}{2}(0.2) + \frac{1}{8}(0.2)^2 = 1 - 0.1 + 0.005 = 0.905$$

1 : approximation

- The series for $f(1.2)$ alternates with terms that decrease in magnitude to 0.

2 : $\begin{cases} 1 : \text{error form} \\ 1 : \text{analysis} \end{cases}$

$$|f(1.2) - T_2(1.2)| \leq \left| \frac{-1}{2^3 \cdot 3}(0.2)^3 \right| = \frac{1}{3000} \leq 0.001$$

AP[®] Calculus BC

2015 Scoring Guidelines

© 2015 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.
Visit the College Board on the Web: www.collegeboard.org.
AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2015 SCORING GUIDELINES

Question 1

The rate at which rainwater flows into a drainpipe is modeled by the function R , where $R(t) = 20\sin\left(\frac{t^2}{35}\right)$ cubic feet per hour, t is measured in hours, and $0 \leq t \leq 8$. The pipe is partially blocked, allowing water to drain out the other end of the pipe at a rate modeled by $D(t) = -0.04t^3 + 0.4t^2 + 0.96t$ cubic feet per hour, for $0 \leq t \leq 8$. There are 30 cubic feet of water in the pipe at time $t = 0$.

- How many cubic feet of rainwater flow into the pipe during the 8-hour time interval $0 \leq t \leq 8$?
- Is the amount of water in the pipe increasing or decreasing at time $t = 3$ hours? Give a reason for your answer.
- At what time t , $0 \leq t \leq 8$, is the amount of water in the pipe at a minimum? Justify your answer.
- The pipe can hold 50 cubic feet of water before overflowing. For $t > 8$, water continues to flow into and out of the pipe at the given rates until the pipe begins to overflow. Write, but do not solve, an equation involving one or more integrals that gives the time w when the pipe will begin to overflow.

(a) $\int_0^8 R(t) dt = 76.570$

2 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b) $R(3) - D(3) = -0.313632 < 0$

Since $R(3) < D(3)$, the amount of water in the pipe is decreasing at time $t = 3$ hours.

2 : $\begin{cases} 1 : \text{considers } R(3) \text{ and } D(3) \\ 1 : \text{answer and reason} \end{cases}$

(c) The amount of water in the pipe at time t , $0 \leq t \leq 8$, is $30 + \int_0^t [R(x) - D(x)] dx$.

3 : $\begin{cases} 1 : \text{considers } R(t) - D(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

$$R(t) - D(t) = 0 \Rightarrow t = 0, 3.271658$$

t	Amount of water in the pipe
0	30
3.271658	27.964561
8	48.543686

The amount of water in the pipe is a minimum at time $t = 3.272$ (or 3.271) hours.

(d) $30 + \int_0^w [R(t) - D(t)] dt = 50$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{equation} \end{cases}$

2015 SCORING GUIDELINES

Question 2

At time $t \geq 0$, a particle moving along a curve in the xy -plane has position $(x(t), y(t))$ with velocity vector $v(t) = (\cos(t^2), e^{0.5t})$. At $t = 1$, the particle is at the point $(3, 5)$.

- (a) Find the x -coordinate of the position of the particle at time $t = 2$.
 (b) For $0 < t < 1$, there is a point on the curve at which the line tangent to the curve has a slope of 2. At what time is the object at that point?
 (c) Find the time at which the speed of the particle is 3.
 (d) Find the total distance traveled by the particle from time $t = 0$ to time $t = 1$.

(a) $x(2) = 3 + \int_1^2 \cos(t^2) dt = 2.557$ (or 2.556)

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

(b) $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{e^{0.5t}}{\cos(t^2)}$

$$\frac{e^{0.5t}}{\cos(t^2)} = 2$$

$$t = 0.840$$

2 : $\begin{cases} 1 : \text{slope in terms of } t \\ 1 : \text{answer} \end{cases}$

(c) Speed = $\sqrt{\cos^2(t^2) + e^t}$

$$\sqrt{\cos^2(t^2) + e^t} = 3$$

$$t = 2.196$$
 (or 2.195)

2 : $\begin{cases} 1 : \text{speed in terms of } t \\ 1 : \text{answer} \end{cases}$

(d) Distance = $\int_0^1 \sqrt{\cos^2(t^2) + e^t} dt = 1.595$ (or 1.594)

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 3

t (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	-220	150

Johanna jogs along a straight path. For $0 \leq t \leq 40$, Johanna's velocity is given by a differentiable function v . Selected values of $v(t)$, where t is measured in minutes and $v(t)$ is measured in meters per minute, are given in the table above.

- (a) Use the data in the table to estimate the value of $v'(16)$.

- (b) Using correct units, explain the meaning of the definite integral $\int_0^{40} |v(t)| dt$ in the context of the problem.

Approximate the value of $\int_0^{40} |v(t)| dt$ using a right Riemann sum with the four subintervals indicated in the table.

- (c) Bob is riding his bicycle along the same path. For $0 \leq t \leq 10$, Bob's velocity is modeled by $B(t) = t^3 - 6t^2 + 300$, where t is measured in minutes and $B(t)$ is measured in meters per minute.

Find Bob's acceleration at time $t = 5$.

- (d) Based on the model B from part (c), find Bob's average velocity during the interval $0 \leq t \leq 10$.

(a) $v'(16) \approx \frac{240 - 200}{20 - 12} = 5$ meters/min²

1 : approximation

- (b) $\int_0^{40} |v(t)| dt$ is the total distance Johanna jogs, in meters, over the time interval $0 \leq t \leq 40$ minutes.

3 : $\begin{cases} 1 : \text{explanation} \\ 1 : \text{right Riemann sum} \\ 1 : \text{approximation} \end{cases}$

$$\begin{aligned} \int_0^{40} |v(t)| dt &\approx 12 \cdot |v(12)| + 8 \cdot |v(20)| + 4 \cdot |v(24)| + 16 \cdot |v(40)| \\ &= 12 \cdot 200 + 8 \cdot 240 + 4 \cdot 220 + 16 \cdot 150 \\ &= 2400 + 1920 + 880 + 2400 \\ &= 7600 \text{ meters} \end{aligned}$$

- (c) Bob's acceleration is $B'(t) = 3t^2 - 12t$.

$$B'(5) = 3(25) - 12(5) = 15 \text{ meters/min}^2$$

2 : $\begin{cases} 1 : \text{uses } B'(t) \\ 1 : \text{answer} \end{cases}$

(d) Avg vel = $\frac{1}{10} \int_0^{10} (t^3 - 6t^2 + 300) dt$

$$= \frac{1}{10} \left[\frac{t^4}{4} - 2t^3 + 300t \right]_0^{10}$$

$$= \frac{1}{10} \left[\frac{10000}{4} - 2000 + 3000 \right] = 350 \text{ meters/min}$$


3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 4

Consider the differential equation $\frac{dy}{dx} = 2x - y$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.
- (b) Find $\frac{d^2y}{dx^2}$ in terms of x and y . Determine the concavity of all solution curves for the given differential equation in Quadrant II. Give a reason for your answer.
- (c) Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(2) = 3$. Does f have a relative minimum, a relative maximum, or neither at $x = 2$? Justify your answer.
- (d) Find the values of the constants m and b for which $y = mx + b$ is a solution to the differential equation.



- 2 : $\begin{cases} 1 : \text{slopes where } x = 0 \\ 1 : \text{slopes where } x = 1 \end{cases}$

(b) $\frac{d^2y}{dx^2} = 2 - \frac{dy}{dx} = 2 - (2x - y) = 2 - 2x + y$

In Quadrant II, $x < 0$ and $y > 0$, so $2 - 2x + y > 0$.
Therefore, all solution curves are concave up in Quadrant II.

- 2 : $\begin{cases} 1 : \frac{d^2y}{dx^2} \\ 1 : \text{concave up with reason} \end{cases}$

(c) $\left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} = 2(2) - 3 = 1 \neq 0$

Therefore, f has neither a relative minimum nor a relative maximum at $x = 2$.

- 2 : $\begin{cases} 1 : \text{considers } \left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} \\ 1 : \text{conclusion with justification} \end{cases}$

(d) $y = mx + b \Rightarrow \frac{dy}{dx} = \frac{d}{dx}(mx + b) = m$

$$2x - y = m$$

$$2x - (mx + b) = m$$

$$(2 - m)x - (m + b) = 0$$

$$2 - m = 0 \Rightarrow m = 2$$

$$b = -m \Rightarrow b = -2$$

- 3 : $\begin{cases} 1 : \frac{d}{dx}(mx + b) = m \\ 1 : 2x - y = m \\ 1 : \text{answer} \end{cases}$

2015 SCORING GUIDELINES

Question 5

Consider the function $f(x) = \frac{1}{x^2 - kx}$, where k is a nonzero constant. The derivative of f is given by

$$f'(x) = \frac{k - 2x}{(x^2 - kx)^2}.$$

- (a) Let $k = 3$, so that $f(x) = \frac{1}{x^2 - 3x}$. Write an equation for the line tangent to the graph of f at the point whose x -coordinate is 4.
- (b) Let $k = 4$, so that $f(x) = \frac{1}{x^2 - 4x}$. Determine whether f has a relative minimum, a relative maximum, or neither at $x = 2$. Justify your answer.
- (c) Find the value of k for which f has a critical point at $x = -5$.
- (d) Let $k = 6$, so that $f(x) = \frac{1}{x^2 - 6x}$. Find the partial fraction decomposition for the function f .

Find $\int f(x) dx$.

(a) $f(4) = \frac{1}{4^2 - 3 \cdot 4} = \frac{1}{4}$ $f'(4) = \frac{3 - 2 \cdot 4}{(4^2 - 3 \cdot 4)^2} = -\frac{5}{16}$

An equation for the line tangent to the graph of f at the point whose x -coordinate is 4 is $y = -\frac{5}{16}(x - 4) + \frac{1}{4}$.

- 2 : $\begin{cases} 1 : \text{slope} \\ 1 : \text{tangent line equation} \end{cases}$

(b) $f'(x) = \frac{4 - 2x}{(x^2 - 4x)^2}$ $f'(2) = \frac{4 - 2 \cdot 2}{(2^2 - 4 \cdot 2)^2} = 0$

$f'(x)$ changes sign from positive to negative at $x = 2$.
Therefore, f has a relative maximum at $x = 2$.

- 2 : $\begin{cases} 1 : \text{considers } f'(2) \\ 1 : \text{answer with justification} \end{cases}$

(c) $f'(-5) = \frac{k - 2 \cdot (-5)}{((-5)^2 - k \cdot (-5))^2} = 0 \Rightarrow k = -10$

- 1 : answer

(d) $\frac{1}{x^2 - 6x} = \frac{1}{x(x - 6)} = \frac{A}{x} + \frac{B}{x - 6} \Rightarrow 1 = A(x - 6) + Bx$

$$x = 0 \Rightarrow 1 = A \cdot (-6) \Rightarrow A = -\frac{1}{6}$$

$$x = 6 \Rightarrow 1 = B \cdot (6) \Rightarrow B = \frac{1}{6}$$

$$\frac{1}{x(x - 6)} = \frac{-1/6}{x} + \frac{1/6}{x - 6}$$

$$\int f(x) dx = \int \left(\frac{-1/6}{x} + \frac{1/6}{x - 6} \right) dx$$

$$= -\frac{1}{6} \ln|x| + \frac{1}{6} \ln|x - 6| + C = \frac{1}{6} \ln \left| \frac{x - 6}{x} \right| + C$$

- 4 : $\begin{cases} 2 : \text{partial fraction decomposition} \\ 2 : \text{general antiderivative} \end{cases}$

2015 SCORING GUIDELINES

Question 6

The Maclaurin series for a function f is given by $\sum_{n=1}^{\infty} \frac{(-3)^{n-1}}{n} x^n = x - \frac{3}{2}x^2 + 3x^3 - \dots + \frac{(-3)^{n-1}}{n} x^n + \dots$ and converges to $f(x)$ for $|x| < R$, where R is the radius of convergence of the Maclaurin series.

- (a) Use the ratio test to find R .
- (b) Write the first four nonzero terms of the Maclaurin series for f' , the derivative of f . Express f' as a rational function for $|x| < R$.
- (c) Write the first four nonzero terms of the Maclaurin series for e^x . Use the Maclaurin series for e^x to write the third-degree Taylor polynomial for $g(x) = e^x f(x)$ about $x = 0$.

- (a) Let a_n be the n th term of the Maclaurin series.

$$\frac{a_{n+1}}{a_n} = \frac{(-3)^n x^{n+1}}{n+1} \cdot \frac{n}{(-3)^{n-1} x^n} = \frac{-3n}{n+1} \cdot x$$

$$\lim_{n \rightarrow \infty} \left| \frac{-3n}{n+1} \cdot x \right| = 3|x|$$

$$3|x| < 1 \Rightarrow |x| < \frac{1}{3}$$

The radius of convergence is $R = \frac{1}{3}$.

- (b) The first four nonzero terms of the Maclaurin series for f' are

$$1 - 3x + 9x^2 - 27x^3.$$

$$f'(x) = \frac{1}{1 - (-3x)} = \frac{1}{1 + 3x}$$

- (c) The first four nonzero terms of the Maclaurin series for e^x are

$$1 + x + \frac{x^2}{2!} + \frac{x^3}{3!}.$$

The product of the Maclaurin series for e^x and the Maclaurin series for f is

$$\left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots\right) \left(x - \frac{3}{2}x^2 + 3x^3 - \dots\right) \\ = x - \frac{1}{2}x^2 + 2x^3 + \dots$$

The third-degree Taylor polynomial for $g(x) = e^x f(x)$

about $x = 0$ is $T_3(x) = x - \frac{1}{2}x^2 + 2x^3$.

1 : sets up ratio
3 : $\begin{cases} 1 : \text{computes limit of ratio} \\ 1 : \text{determines radius of convergence} \end{cases}$

3 : $\begin{cases} 2 : \text{first four nonzero terms} \\ 1 : \text{rational function} \end{cases}$

3 : $\begin{cases} 1 : \text{first four nonzero terms} \\ \text{of the Maclaurin series for } e^x \\ 2 : \text{Taylor polynomial} \end{cases}$