


# Differentiation: Definition and Fundamental Properties

## AP Calculus BC

### Average and Instantaneous Rates of Change at a Point

Unit 1 built the limit. Unit 2 uses it to define the **derivative** 导数—the exact rate of change at a point.



*The instantaneous rate of change is the gradient of the tangent at a point*

Over an interval, the average rate of change is a **difference quotient** 差商. Two equivalent forms appear:

$$\frac{f(a+h) - f(a)}{h} \quad \text{and} \quad \frac{f(x) - f(a)}{x - a}.$$

The first uses a step of size  $h$  from  $a$ ; the second uses two points  $x$  and  $a$ . Both compute  $\frac{\text{change in output}}{\text{change in input}}$  over the interval.

The **instantaneous** 瞬时 rate of change at  $x = a$  is what the difference quotient approaches as the interval shrinks to zero. This limit is the derivative at  $a$ , written  $f'(a)$ :

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a},$$

provided the limit exists.

# Defining the Derivative and Reading Its Notation

Let the point  $a$  vary and the derivative becomes a **new function**:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}.$$

This is the **definition of the derivative** (sometimes called differentiating "by first principles" 用定义求导). Its value at each  $x$  is the instantaneous rate of change there.

Common **notations** 记号 for the derivative of  $y = f(x)$  are:


$$\frac{dy}{dx}, \quad f'(x), \quad y'.$$

The derivative can be represented graphically, numerically, analytically, and verbally – be ready to move between them.

**Geometric meaning.** The derivative at a point is the **slope** 斜率 of the **tangent line** 切线 to the graph there. So the tangent line at  $x = a$  passes through  $(a, f(a))$  with slope  $f'(a)$ :

$$y - f(a) = f'(a)(x - a).$$

Writing this line is a routine exam task, so keep the point-slope form ready.



*Secant slopes approach the tangent slope: the derivative is the limit of average rates*

## Estimating a Derivative at a Point

You do not always have a formula. When a function is given by a **table** 表格 or a graph, estimate the derivative  $f'(a)$  with a difference quotient over a **small interval around**  $a$ . A table with values on both sides of  $a$  gives the best estimate:

$$f'(a) \approx \frac{f(b) - f(c)}{b - c}, \quad \text{where } c < a < b \text{ are the closest table inputs.}$$

Technology (a calculator) can also estimate a derivative at a point.

**Exam skill (appears almost every year).** Questions such as "Approximate  $M'(7.5)$  using the average rate of change of  $M$  over the interval  $5 \leq t \leq 10$ " ask for exactly this difference quotient. Show the setup:

$$M'(7.5) \approx \frac{M(10) - M(5)}{10 - 5}.$$

Full credit needs the numbers plugged in **and** the correct **units** 单位 (output units per input unit), since these come from real-world models.


## Differentiability and Continuity: When a Derivative Exists

Differentiability is stronger than continuity. The key relationship:

**If  $f$  is differentiable 可导 at a point, then  $f$  is continuous 连续 there.**

So differentiability *implies* continuity. The reverse is false: a continuous function can fail to be differentiable. Two ways this happens:

- A **corner** 尖点: the left and right difference-quotient limits disagree, as with  $f(x) = |x|$  at  $x = 0$ .
- A **vertical tangent** 垂直切线: the slope is infinite (no real number), as with  $f(x) = \sqrt[3]{x}$  at  $x = 0$ .



*Two ways a continuous function is not differentiable: a corner and a vertical tangent*

Also, a point outside the domain of  $f$  cannot be in the domain of  $f'$ . Use the contrapositive on the exam: **if  $f$  is not continuous at  $a$ , then  $f$  is not differentiable at  $a$ .**

## The Power Rule

From here we use **rules** instead of the limit definition each time. The **power rule** 幂法则 handles any power of  $x$ :

$$\frac{d}{dx} x^r = r x^{r-1} \quad \text{for any real } r.$$

It works for whole-number powers, negative powers ( $\frac{1}{x} = x^{-1}$ ), and roots ( $\sqrt{x} = x^{1/2}$ ) – rewrite as a power first, then apply the rule.

## Constant, Sum, Difference, and Constant Multiple Rules

These rules let you differentiate term by term:

- **Constant:**  $\frac{d}{dx} k = 0$  (a constant does not change).
- **Constant multiple** 常数倍:  $\frac{d}{dx} [k f(x)] = k f'(x)$ .
- **Sum / difference:**  $\frac{d}{dx} [f(x) \pm g(x)] = f'(x) \pm g'(x)$ .

Combined with the power rule, they differentiate any **polynomial** 多项式 term by term. Example:

$$\frac{d}{dx} (4x^3 - 5x + 7) = 12x^2 - 5.$$

## Derivatives of $\cos x$ , $\sin x$ , $e^x$ , and $\ln x$

Learn these four building-block derivatives by heart:

$$\frac{d}{dx} \sin x = \cos x, \quad \frac{d}{dx} \cos x = -\sin x,$$

$$\frac{d}{dx} e^x = e^x, \quad \frac{d}{dx} \ln x = \frac{1}{x} \quad (x > 0).$$

Note the minus sign on the derivative of cosine, and that  $e^x$  is its own derivative.

**A limit that is really a derivative (LIM-3.A.1).** Sometimes a limit is secretly the definition of a known derivative. If you recognize

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

for a function  $f$  whose derivative you know, just evaluate  $f'(a)$ . For example,

$$\lim_{h \rightarrow 0} \frac{\sin\left(\frac{\pi}{2} + h\right) - 1}{h} = \left. \frac{d}{dx} \sin x \right|_{x=\pi/2} = \cos \frac{\pi}{2} = 0.$$

## The Product Rule

A product of two functions is **not** differentiated by multiplying the derivatives. Use the **product rule** 乘积法则:

$$\frac{d}{dx}[uv] = u'v + uv'$$

"Derivative of the first times the second, plus the first times the derivative of the second."

Example:

$$\frac{d}{dx}(x^2 e^x) = 2x e^x + x^2 e^x.$$

Exam questions often build a new function from given pieces, e.g.  $k'(x) = (f(x))^2 g(x)$ , and ask you to combine rules while reading values from a table.

## The Quotient Rule

For a quotient, use the **quotient rule** 商法则:

$$\frac{d}{dx}\left[\frac{u}{v}\right] = \frac{u'v - uv'}{v^2}.$$

"Bottom times derivative of top, minus top times derivative of bottom, all over bottom squared." The order matters because of the minus sign, so write the numerator carefully.

Example:

$$\frac{d}{dx}\left(\frac{x}{\cos x}\right) = \frac{1 \cdot \cos x - x \cdot (-\sin x)}{\cos^2 x} = \frac{\cos x + x \sin x}{\cos^2 x}.$$

## Derivatives of Tangent, Cotangent, Secant, and Cosecant

The remaining trigonometric derivatives are not memorized separately –you **rewrite them with identities** 恒等式 and apply the quotient (or product) rule. For instance,  $\tan x = \frac{\sin x}{\cos x}$ , so the quotient rule gives

$$\frac{d}{dx} \tan x = \frac{\cos x \cos x - \sin x(-\sin x)}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x.$$

The same method (writing  $\cot x = \frac{\cos x}{\sin x}$ ,  $\sec x = \frac{1}{\cos x}$ ,  $\csc x = \frac{1}{\sin x}$ ) gives  $-\csc^2 x$ ,  $\sec x \tan x$ , and  $-\csc x \cot x$ .

**Higher-order derivatives.** Differentiating  $f'$  again gives the **second derivative** 二阶导数  $f''(x)$  (or  $\frac{d^2y}{dx^2}$ ) –the rate of change of the rate of change. An exam part like "Find  $k''(3)$ " just means differentiate twice, then substitute. You can also estimate a second derivative from a table by applying the average-rate-of-change method to the  $f'$  values.

## Exam tips

- The derivative is the **slope of the tangent** —the limit of the secant slope  $\frac{f(x+h)-f(x)}{h}$  as  $h \rightarrow 0$ .

- Memorise the rules: power, product, quotient, and the derivatives of  $\sin$ ,  $\cos$ ,  $e^x$ , and  $\ln x$ .
- **Differentiability implies continuity**, but not the reverse (a corner or cusp is continuous yet not differentiable).
- Distinguish average rate of change (secant slope over an interval) from instantaneous rate (the derivative at a point).
- Give a tangent-line equation as  $y - f(a) = f'(a)(x - a)$ .