


Applications of Integration

AP Calculus BC • Unit 8


AP Calculus BC



Applications of Integration

What we will cover

- 1 Average value of a function
- 2 Motion with integrals
- 3 Net change in context
- 4 Area between curves
- 5 Volumes by cross section
- 6 Volumes of revolution



1

Average value

Applications of Integration

Average value

Average value of a function

The **average value** 平均值 over $[a, b]$:

$$f_{\text{avg}} = \frac{1}{b-a} \int_a^b f(x) dx.$$

It is the **constant height** of a rectangle on $[a, b]$ with the **same area** as under f .

Do **not** confuse it with the **average rate of change** (which divides the **change in f** by the interval).

Average value of $f(x) = x^2$ on $[0, 3]$

$$\frac{1}{3} \int_0^3 x^2 dx = \frac{1}{3} \left[\frac{x^3}{3} \right]_0^3 = 3.$$

2

Motion

Applications of Integration

Motion

Motion, run backwards

Integration reverses Unit 4's links:

- **displacement** 位移 $= \int_{t_1}^{t_2} v(t) dt$
- **total distance** 总路程 $= \int_{t_1}^{t_2} |v(t)| dt$
- **position** $= x(t_1) + \int_{t_1}^t v(s) ds$

The displacement-vs-distance distinction (that absolute value) is a **classic graded point**.

Net change in an applied context

The **net change** 净变化 theorem is the everyday tool:

$$\text{final} = \text{initial} + \int_a^b (\text{rate}) dt.$$

"How much water is in the tank at $t = 5$ " =
start + $\int_0^5 (\text{in} - \text{out}) dt$.



bridge Suspension


3 Area between curves

Area between two curves

On $[a, b]$ with f on top and g below:

$$A = \int_a^b [f(x) - g(x)] dx.$$

Find the **intersections** first to set the limits, and always subtract **bottom from top**.




Worked example – the area between $y = x$ and $y = x^2$

Find the enclosed area

They cross at $x = 0$ and $x = 1$; on $(0, 1)$ the line is on top:

$$\begin{aligned} A &= \int_0^1 (x - x^2) dx = \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 \\ &= \frac{1}{2} - \frac{1}{3} = \frac{1}{6}. \end{aligned}$$



Horizontal slices, and more than two crossings

Slice in y
 $A = \int_c^d [x_{\text{right}}(y) - x_{\text{left}}(y)] dy$
Choose x - or y -slices to keep **one** top/bottom pair over the whole region.

Several crossings
 $A = \int_a^b |f(x) - g(x)| dx$
or **split into a sum** at each crossing – decide which curve is higher on each piece.

4 Volumes

Volumes with known cross sections

If a solid has a known **cross section** 横截面 perpendicular to an axis, the volume is the integral of its **area**:

$$V = \int_a^b A(x) dx.$$

The exam phrase “the base of a solid... cross sections are squares” signals exactly this.

Set the shape's key length equal to the slice length $s(x) = f(x) - g(x)$, then integrate.

square of side s : $A = s^2$

rectangle, height ks : $A = ks^2$

equilateral triangle: $A = \frac{\sqrt{3}}{4}s^2$

semicircle, diameter s : $A = \frac{\pi}{8}s^2$


Volumes of revolution – the disc method

rotate $y = f(x)$ about the x -axis

Revolving a region about an axis makes a **solid of revolution** 旋转体. If the region **touches** the axis, each slice is a **disc** 圆盘:

$$V = \pi \int_a^b [r(x)]^2 dx.$$

About the y -axis: express x as a function of y and integrate in y .



$$V = \pi \int_a^b [f(x)]^2 dx$$

5

Recap

Unit 8 – key takeaways

1 Average value

$\frac{1}{b-a} \int_a^b f$ – not the average rate of change

2 Motion

displacement integrates v ; distance integrates $|v|$

3 Net change

final = initial + $\int$ rate – watch signs and units

4 Area & volume

top minus bottom; $V = \int A(x) dx$; discs use πr^2


Check yourself

1 Which integral gives total distance, $\int v$ or $\int |v|$?

2 Find the area between $y = x$ and $y = x^2$.

3 Write the average value of f on $[a, b]$.

4 A solid's cross sections are squares of side $s(x)$. Give V .



Answers 1. $\int |v|$ 2. $\frac{1}{6}$ 3. $\frac{1}{b-a} \int_a^b f$ 4. $V = \int_a^b [s(x)]^2 dx$