


Infinite Sequences & Series

AP Calculus BC • Unit 10


AP Calculus BC



Infinite Sequences & Series

What we will cover

- 1 Convergence & partial sums
- 2 Geometric series
- 3 The convergence tests
- 4 Absolute vs conditional
- 5 Taylor polynomials & error
- 6 Power series & intervals



1 Convergence


Infinite Sequences & Series

Convergence

Convergent and divergent series

An **infinite series** 无穷级数 $\sum a_n$ is defined as the limit of its **partial sums** 部分和 $S_N = a_1 + \dots + a_N$.

Every convergence question is really a question about the limit of the partial sums.



Infinite Sequences & Series

Convergence

Geometric series – the one you can sum exactly

A **geometric series** 几何级数 has a constant ratio r . It converges **exactly** when $|r| < 1$:

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}.$$

It underlies **power series** later in the unit.

Sum $3 + \frac{3}{2} + \frac{3}{4} + \frac{3}{8} + \dots$

Here $a = 3$ and $r = \frac{1}{2}$, and $|r| < 1$:

$$\frac{a}{1-r} = \frac{3}{1-\frac{1}{2}} = 6.$$

2 The tests

Infinite Sequences & Series
└ The tests

Start here – the n th term test

If the terms do not shrink to zero, the sum cannot settle:

$$\lim_{n \rightarrow \infty} a_n \neq 0 \implies \text{diverges.}$$

Always check this **quick test** first.

✗ It is **only** a test for **divergence**

If $a_n \rightarrow 0$ the test is **inconclusive** – the series may **still** diverge

Infinite Sequences & Series
└ The tests

p -series and the harmonic series

$$\sum \frac{1}{n^p} \begin{cases} \text{converges} & p > 1 \\ \text{diverges} & p \leq 1 \end{cases}$$

The **integral test** 积分判别法 proves this: for positive, decreasing, continuous f , $\sum a_n$ and $\int_1^\infty f$ both converge or both diverge.

The **harmonic series** 调和级数 $\sum \frac{1}{n}$ ($p = 1$) **diverges** – even though its terms go to zero. A famous, must-know fact.

The p -series family is the standard **yardstick** for the comparison tests.

Infinite Sequences & Series
└ The tests

Comparison and alternating series

Direct comparison 直接比较
 $0 \leq a_n \leq b_n$ and $\sum b_n$ converges $\implies$ so does $\sum a_n$

Limit comparison 极限比较
 $\lim \frac{a_n}{b_n}$ finite and positive $\implies$ both do the **same** thing

Alternating series 交错级数
 b_n positive, **decreasing**, and $b_n \rightarrow 0 \implies$ converges

Ratio test 比值判别法
 $L = \lim \left| \frac{a_{n+1}}{a_n} \right|$: $L < 1$ converges, $L > 1$ diverges, $L = 1$ **inconclusive**

Infinite Sequences & Series
└ The tests

Worked example – the ratio test

Test $\sum \frac{n}{2^n}$

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{n+1}{2^{n+1}} \cdot \frac{2^n}{n} = \frac{n+1}{2n} \rightarrow \frac{1}{2} < 1$$

so the series **converges**.

The ratio test is the go-to for **factorials** or **n th powers** – and it is exactly how you find a **radius of convergence**.

Infinite Sequences & Series
└ The tests

Absolute vs conditional convergence

Absolutely 绝对收敛
 $\sum |a_n|$ converges – the **stronger** property

Conditionally 条件收敛
 $\sum a_n$ converges but $\sum |a_n|$ does not

The classic: $\sum \frac{(-1)^n}{n}$ converges **conditionally** – it relies on the **cancellation of signs**, since $\sum \frac{1}{n}$ diverges.

Alternating error bound

$$|S - S_N| \leq b_{N+1}$$

The error is **no larger** than the first omitted term.

3


Taylor

Taylor polynomials

A **Taylor polynomial** 泰勒多项式 approximates f near a centre a using its derivatives there:

$$P_n(x) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k.$$

Each added term matches **one more derivative**. Centred at $a = 0$ it is a **Maclaurin** polynomial.



The Lagrange error bound

How far can a Taylor polynomial be from the truth?

$$|R_n(x)| \leq \frac{\max |f^{(n+1)}(z)|}{(n+1)!} |x-a|^{n+1}.$$

Bound the $(n+1)$ th derivative on the interval, then compute – the standard way to **prove** an estimate is accurate enough.

Radius and interval of convergence

A **power series** 幂级数 $\sum c_n(x-a)^n$ converges within a **radius** 收敛半径 R of its centre. Find R with the **ratio test**, then test the **endpoints separately**.

The ratio test is **inconclusive at the endpoints** – that is why they need their own check.

Interval of $\sum \frac{x^n}{n}$

Ratio: $|x| \frac{n}{n+1} \rightarrow |x| < 1$, so $R = 1$. $x = -1$: alternating harmonic – converges.

$x = 1$: harmonic – diverges.

Interval = $[-1, 1)$.

The four Maclaurin series to memorise

$$e^x = \sum \frac{x^n}{n!}$$

$$\frac{1}{1-x} = \sum x^n$$

$$\sin x = \sum \frac{(-1)^n x^{2n+1}}{(2n+1)!}$$

$$\cos x = \sum \frac{(-1)^n x^{2n}}{(2n)!}$$

New series come from **manipulating** these – substituting, differentiating, integrating, multiplying.

Worked example – the substitution trick

Maclaurin series for e^{x^2}

Substitute x^2 for x in $e^x = \sum \frac{x^n}{n!}$:

$$e^{x^2} = \sum_{n=0}^{\infty} \frac{(x^2)^n}{n!} = 1 + x^2 + \frac{x^4}{2!} + \frac{x^6}{3!} + \dots$$

which converges for all x .

This substitution is **far faster** than differentiating e^{x^2} six times – which is exactly what the exam is testing.

4

Recap

Unit 10 – key takeaways

1 Convergence

the limit of the partial sums;
 check $a_n \rightarrow 0$ first

3 The tests

p -series yardstick; comparison,
 alternating, ratio

2 Geometric

$|r| < 1 \Rightarrow a/(1-r)$ – the one
 exact sum

4 Taylor

P_n matches derivatives at a ; ratio
 test gives R


Check yourself

1 Does $\sum \frac{1}{n}$ converge?

2 Sum $3 + \frac{3}{2} + \frac{3}{4} + \dots$

3 The ratio test gives $L = 1$. What follows?

4 Give the Maclaurin series for e^{x^2} .



Answers 1. no – it diverges 2. 6 3. inconclusive – use another test 4. $\sum x^{2n}/n!$