

6. The Taylor series for a function f about $x = 4$ is given by

$\sum_{n=1}^{\infty} \frac{(x-4)^{n+1}}{(n+1)3^n} = \frac{(x-4)^2}{2 \cdot 3} + \frac{(x-4)^3}{3 \cdot 3^2} + \frac{(x-4)^4}{4 \cdot 3^3} + \dots + \frac{(x-4)^{n+1}}{(n+1)3^n} + \dots$ and converges to $f(x)$ on its interval of convergence.

- A. Using the ratio test, find the interval of convergence of the Taylor series for f about $x = 4$. Justify your answer.
- B. Find the first three nonzero terms and the general term of the Taylor series for f' , the derivative of f , about $x = 4$.
- C. The Taylor series for f' described in part B is a geometric series. For all x in the interval of convergence of the Taylor series for f' , show that $f'(x) = \frac{x-4}{7-x}$.
- D. It is known that the radius of convergence of the Taylor series for f about $x = 4$ is the same as the radius of convergence of the Taylor series for f' about $x = 4$. Does the Taylor series for f' described in part B converge to $f'(x) = \frac{x-4}{7-x}$ at $x = 8$? Give a reason for your answer.

STOP
END OF EXAM

6. The function g has derivatives of all orders for all real numbers. The Maclaurin series for g is given by

$$g(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^n}{2e^n + 3} \text{ on its interval of convergence.}$$

- (a) State the conditions necessary to use the integral test to determine convergence of the series $\sum_{n=0}^{\infty} \frac{1}{e^n}$.

Use the integral test to show that $\sum_{n=0}^{\infty} \frac{1}{e^n}$ converges.

- (b) Use the limit comparison test with the series $\sum_{n=0}^{\infty} \frac{1}{e^n}$ to show that the series $g(1) = \sum_{n=0}^{\infty} \frac{(-1)^n}{2e^n + 3}$

converges absolutely.

- (c) Determine the radius of convergence of the Maclaurin series for g .

- (d) The first two terms of the series $g(1) = \sum_{n=0}^{\infty} \frac{(-1)^n}{2e^n + 3}$ are used to approximate $g(1)$. Use the alternating

series error bound to determine an upper bound on the error of the approximation.

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

STOP

END OF EXAM

$$\begin{aligned}f(0) &= 0 \\f'(0) &= 1 \\f^{(n+1)}(0) &= -n \cdot f^{(n)}(0) \text{ for all } n \geq 1\end{aligned}$$

6. A function f has derivatives of all orders for $-1 < x < 1$. The derivatives of f satisfy the conditions above.

The Maclaurin series for f converges to $f(x)$ for $|x| < 1$.

- (a) Show that the first four nonzero terms of the Maclaurin series for f are $x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4}$, and write the

general term of the Maclaurin series for f .

- (b) Determine whether the Maclaurin series described in part (a) converges absolutely, converges conditionally, or diverges at $x = 1$. Explain your reasoning.

- (c) Write the first four nonzero terms and the general term of the Maclaurin series for $g(x) = \int_0^x f(t) \, dt$.

- (d) Let $P_n\left(\frac{1}{2}\right)$ represent the n th-degree Taylor polynomial for g about $x = 0$ evaluated at $x = \frac{1}{2}$, where g is

the function defined in part (c). Use the alternating series error bound to show that

$$\left| P_4\left(\frac{1}{2}\right) - g\left(\frac{1}{2}\right) \right| < \frac{1}{500}.$$

STOP
END OF EXAM