


**Part B (AB or BC): Graphing calculator not allowed****Question 4****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.



The graph of the differentiable function  $f$ , shown for  $-6 \leq x \leq 7$ , has a horizontal tangent at  $x = -2$  and is linear for  $0 \leq x \leq 7$ . Let  $R$  be the region in the second quadrant bounded by the graph of  $f$ , the vertical line  $x = -6$ , and the  $x$ - and  $y$ -axes. Region  $R$  has area 12.

**Model Solution****Scoring**

- (a) The function  $g$  is defined by  $g(x) = \int_0^x f(t) dt$ . Find the values of  $g(-6)$ ,  $g(4)$ , and  $g(6)$ .

|                                                                                           |         |                |
|-------------------------------------------------------------------------------------------|---------|----------------|
| $g(-6) = \int_0^{-6} f(t) dt = -\int_{-6}^0 f(t) dt = -12$                                | $g(-6)$ | <b>1 point</b> |
| $g(4) = \int_0^4 f(t) dt = \frac{1}{2} \cdot 4 \cdot 2 = 4$                               | $g(4)$  | <b>1 point</b> |
| $g(6) = \int_0^6 f(t) dt = \frac{1}{2} \cdot 4 \cdot 2 - \frac{1}{2} \cdot 2 \cdot 1 = 3$ | $g(6)$  | <b>1 point</b> |

**Scoring notes:**

- Supporting work is not required for any of these values. However, any supporting work that is shown must be correct to earn the corresponding point.
- Special case: A response that explicitly presents  $g(x) = \int_{-6}^x f(t) dt$  does not earn the first point it would have otherwise earned. The response is eligible for all subsequent points for correct answers, or for consistent answers with supporting work.
  - Note:  $\int_{-6}^{-6} f(t) dt = 0$ ,  $\int_{-6}^4 f(t) dt = 16$ ,  $\int_{-6}^6 f(t) dt = 15$
- Labeled values may be presented in any order. Unlabeled values are read from left to right and from top to bottom as  $g(-6)$ ,  $g(4)$ , and  $g(6)$ , respectively. A response that presents only 1 or 2 values must label them to earn any points.

**Total for part (a) 3 points**

- (b) For the function  $g$  defined in part (a), find all values of  $x$  in the interval  $0 \leq x \leq 6$  at which the graph of  $g$  has a critical point. Give a reason for your answer.

|                                                               |                                 |                |
|---------------------------------------------------------------|---------------------------------|----------------|
| $g'(x) = f(x)$                                                | Fundamental Theorem of Calculus | <b>1 point</b> |
| $g'(x) = f(x) = 0 \Rightarrow x = 4$                          | Answer with reason              | <b>1 point</b> |
| Therefore, the graph of $g$ has a critical point at $x = 4$ . |                                 |                |

**Scoring notes:**

- The first point is earned for explicitly making the connection  $g' = f$  in this part.
  - A response that writes  $g'' = f'$  earns the first point but can only earn the second point by reasoning from  $f = 0$ .
- A response that does not earn the first point is eligible to earn the second point with an implied application of the FTC (e.g., “Because  $g'(4) = 0$ ,  $x = 4$  is a critical point”).
- A response that reports any additional critical points in  $0 < x < 6$  does not earn the second point.
  - Any presented critical point outside the interval  $0 < x < 6$  will not affect scoring.

**Total for part (b) 2 points**

- (c) The function  $h$  is defined by  $h(x) = \int_{-6}^x f'(t) dt$ . Find the values of  $h(6)$ ,  $h'(6)$ , and  $h''(6)$ . Show the work that leads to your answers.

|                                                                |                                      |                |
|----------------------------------------------------------------|--------------------------------------|----------------|
| $h(6) = \int_{-6}^6 f'(t) dt = f(6) - f(-6) = -1 - 0.5 = -1.5$ | Uses Fundamental Theorem of Calculus | <b>1 point</b> |
|                                                                | $h(6)$ with supporting work          | <b>1 point</b> |
| $h'(x) = f'(x)$ , so $h'(6) = f'(6) = -\frac{1}{2}$ .          | $h'(6)$                              | <b>1 point</b> |
| $h''(x) = f''(x)$ , so $h''(6) = f''(6) = 0$ .                 | $h''(6)$                             | <b>1 point</b> |

**Scoring notes:**

- Labeled values may be presented in any order.
- Unlabeled values are read from left to right and from top to bottom as  $h(6)$ ,  $h'(6)$ , and  $h''(6)$ , respectively. A response that presents only 1 or 2 values must label them in order to earn any points.
- A response of  $h(6) = -1.5$  does not earn either of the first 2 points. A response of  $h(6) = f(6) - f(-6)$  earns the first point but not yet the second point.
- A response of  $h(6) = -1 - 0.5$  is the minimum work required to earn both of the first 2 points.
- To earn the third point a response must state either  $h'(x) = f'(x)$  or  $h'(6) = f'(6)$ , and provide an answer of  $-\frac{1}{2}$ .
- The fourth point is earned for a response of  $h''(6) = 0$ , with or without supporting work.
- A response that has one or more linkage errors does not earn the first point it would have otherwise earned. For example,  $h'(x) = f'(6) = -\frac{1}{2}$  does not earn the third point but is eligible for the fourth point even in the presence of another linkage error, such as  $h''(x) = f''(6) = 0$ .

**Total for part (c)    4 points**

**Total for question 4    9 points**

**Part A (AB or BC): Graphing calculator required****Question 1****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

|                                |   |     |      |     |      |     |
|--------------------------------|---|-----|------|-----|------|-----|
| $t$<br>(seconds)               | 0 | 60  | 90   | 120 | 135  | 150 |
| $f(t)$<br>(gallons per second) | 0 | 0.1 | 0.15 | 0.1 | 0.05 | 0   |

A customer at a gas station is pumping gasoline into a gas tank. The rate of flow of gasoline is modeled by a differentiable function  $f$ , where  $f(t)$  is measured in gallons per second and  $t$  is measured in seconds since pumping began. Selected values of  $f(t)$  are given in the table.

**Model Solution****Scoring**

- (a) Using correct units, interpret the meaning of  $\int_{60}^{135} f(t) dt$  in the context of the problem. Use a right Riemann sum with the three subintervals  $[60, 90]$ ,  $[90, 120]$ , and  $[120, 135]$  to approximate the value of  $\int_{60}^{135} f(t) dt$ .

$\int_{60}^{135} f(t) dt$  represents the total number of gallons of gasoline pumped into the gas tank from time  $t = 60$  seconds to time  $t = 135$  seconds.

Interpretation with units **1 point**

$\int_{60}^{135} f(t) dt$   
 $\approx f(90)(90 - 60) + f(120)(120 - 90) + f(135)(135 - 120)$   
 $= (0.15)(30) + (0.1)(30) + (0.05)(15) = 8.25$

Form of Riemann sum **1 point**

Answer **1 point**

**Scoring notes:**

- To earn the first point the response must reference gallons of gasoline added/pumped and the time interval  $t = 60$  to  $t = 135$ .
- To earn the second point at least five of the six factors in the Riemann sum must be correct.
- If there is any error in the Riemann sum, the response does not earn the third point.
- A response of  $(0.15)(30) + (0.1)(30) + (0.05)(15)$  earns both the second and third points, unless there is a subsequent error in simplification, in which case the response would earn only the second point.

- A response that presents a correct value with accompanying work that shows the three products in the Riemann sum but does not show all six of the factors and/or the sum process, does not earn the second point but does earn the third point. For example, responses of either  $4.5 + 3.0 + 0.75$  or  $(0.15)(30)$ ,  $0.1(30)$ ,  $0.05(15) \rightarrow 8.25$  earn the third point but not the second.
- A response of  $f(90)(90 - 60) + f(120)(120 - 90) + f(135)(135 - 120) = 8.25$  earns both the second and the third points.
- A response that presents an answer of only 8.25 does not earn either the second or third point.
- A response that provides a completely correct left Riemann sum with accompanying work,  $f(60)(30) + f(90)(30) + f(120)(15) = 9$ , or  $(0.1)(30) + (0.15)(30) + (0.1)(15)$  earns 1 of the last 2 points. A response with any errors or missing factors in a left Riemann sum earns neither of the last 2 points.

**Total for part (a) 3 points****(b)** Must there exist a value of  $c$ , for  $60 < c < 120$ , such that  $f'(c) = 0$ ? Justify your answer. $f$  is differentiable.  $\Rightarrow f$  is continuous on  $[60, 120]$ . $f(120) - f(60) = 0$  **1 point**

$$\frac{f(120) - f(60)}{120 - 60} = \frac{0.1 - 0.1}{60} = 0$$

Answer with justification **1 point**

By the Mean Value Theorem, there must exist a  $c$ , for  $60 < c < 120$ , such that  $f'(c) = 0$ .

**Scoring notes:**

- To earn the first point a response must present either  $f(120) - f(60) = 0$ ,  $0.1 - 0.1 = 0$  (perhaps as the numerator of a quotient), or  $f(60) = f(120)$ .
- To earn the second point a response must:
  - have earned the first point,
  - state that  $f$  is continuous because  $f$  is differentiable (or equivalent), and
  - answer “yes” in some way.
- A response may reference either the Mean Value Theorem or Rolle’s Theorem.
- A response that references the Intermediate Value Theorem cannot earn the second point.

**Total for part (b) 2 points****(c)** The rate of flow of gasoline, in gallons per second, can also be modeled by

$$g(t) = \left(\frac{t}{500}\right) \cos\left(\left(\frac{t}{120}\right)^2\right) \text{ for } 0 \leq t \leq 150. \text{ Using this model, find the average rate of flow of}$$

gasoline over the time interval  $0 \leq t \leq 150$ . Show the setup for your calculations.

$$\frac{1}{150 - 0} \int_0^{150} g(t) dt$$

Average value formula **1 point**

$$= 0.0959967$$

Answer **1 point**

The average rate of flow of gasoline, in gallons per second, is 0.096 (or 0.095).

**Scoring notes:**

- The exact value of  $\frac{1}{150} \int_0^{150} g(t) dt$  is  $\frac{12}{125} \sin\left(\frac{25}{16}\right)$ .
- A response may present the average value formula in single or multiple steps. For example, the following response earns both points:  $\int_0^{150} g(t) dt = 14.399504$  so the average rate is 0.0959967.
- A response that presents the average value formula in multiple steps but provides incorrect or incomplete communication (e.g.,  $\int_0^{150} g(t) dt = \frac{14.399504}{150} = 0.0959967$ ) earns 1 out of 2 points.
- A response of  $\int_0^{150} g(t) dt = 0.0959967$  does not earn either point.
- Degree mode: A response that presents answers obtained by using a calculator in degree mode does not earn the first point it would have otherwise earned. The response is generally eligible for all subsequent points (unless no answer is possible in degree mode or the question is made simpler by using degree mode). In degree mode,  $\frac{1}{150} \int_0^{150} g(t) dt = 0.149981$  or 0.002618.

**Total for part (c) 2 points**

- (d) Using the model  $g$  defined in part (c), find the value of  $g'(140)$ . Interpret the meaning of your answer in the context of the problem.

|                                                                                                                                                         |                               |
|---------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------|
| $g'(140) \approx -0.004908$<br>$g'(140) = -0.005$ (or $-0.004$ )                                                                                        | $g'(140)$ <b>1 point</b>      |
| The rate at which gasoline is flowing into the tank is decreasing at a rate of 0.005 (or 0.004) gallon per second per second at time $t = 140$ seconds. | Interpretation <b>1 point</b> |

**Scoring notes:**

- The exact value of  $g'(140)$  is  $\frac{1}{500} \cos\left(\frac{49}{36}\right) - \frac{49}{9000} \sin\left(\frac{49}{36}\right)$ .
- The value of  $g'(140)$  may appear only in the interpretation.
- To be eligible for the second point a response must present some numerical value for  $g'(140)$ .
- To earn the second point the interpretation must include “the rate of flow of gasoline is changing at a rate of [the declared value of  $g'(140)$ ]” and “at  $t = 140$ ” (or equivalent).
- An interpretation of “decreasing at a rate of  $-0.005$ ” or “increasing at a rate of 0.005” does not earn the second point.
- Degree mode: In degree mode,  $g'(140) = 0.001997$  or 0.00187.

**Total for part (d) 2 points****Total for question 1 9 points**

**Part A (AB or BC): Graphing calculator required****Question 1****9 points****General Scoring Notes**

The model solution is presented using standard mathematical notation.

Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be correct to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

From 5 A.M. to 10 A.M., the rate at which vehicles arrive at a certain toll plaza is given by

$A(t) = 450\sqrt{\sin(0.62t)}$ , where  $t$  is the number of hours after 5 A.M. and  $A(t)$  is measured in vehicles per hour. Traffic is flowing smoothly at 5 A.M. with no vehicles waiting in line.

| Model Solution                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Scoring               |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| (a) Write, but do not evaluate, an integral expression that gives the total number of vehicles that arrive at the toll plaza from 6 A.M. ( $t = 1$ ) to 10 A.M. ( $t = 5$ ).                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                       |
| The total number of vehicles that arrive at the toll plaza from 6 A.M. to 10 A.M. is given by $\int_1^5 A(t) dt$ .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | Answer <b>1 point</b> |
| <b>Scoring notes:</b> <ul style="list-style-type: none"> <li>The response must be a definite integral with correct lower and upper limits to earn this point.</li> <li>Because <math> A(t)  = A(t)</math> for <math>1 \leq t \leq 5</math>, a response of <math>\int_1^5  450\sqrt{\sin(0.62t)}  dt</math> or <math>\int_1^5  A(t)  dt</math> earns the point.</li> <li>A response missing <math>dt</math> or using <math>dx</math> is eligible to earn the point.</li> <li>A response with a copy error in the expression for <math>A(t)</math> will earn the point only in the presence of <math>\int_1^5 A(t) dt</math>.</li> </ul> |                       |
| <b>Total for part (a) 1 point</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |                       |

- (b) Find the average value of the rate, in vehicles per hour, at which vehicles arrive at the toll plaza from 6 A.M. ( $t = 1$ ) to 10 A.M. ( $t = 5$ ).

$$\text{Average} = \frac{1}{5-1} \int_1^5 A(t) dt = 375.536966$$

Uses average value formula: **1 point**

$$\frac{1}{b-a} \int_a^b A(t) dt$$

The average rate at which vehicles arrive at the toll plaza from 6 A.M. to 10 A.M. is 375.537 (or 375.536) vehicles per hour.

Answer **1 point**

**Scoring notes:**

- The use of the average value formula, indicating that  $a = 1$  and  $b = 5$ , can be presented in single or multiple steps to earn the first point. For example, the following response earns both points:

$$\int_1^5 A(t) dt = 1502.147865, \text{ so the average value is } 375.536966.$$

- A response that presents a correct integral along with the correct average value, but provides incorrect or incomplete communication, earns 1 out of 2 points. For example, the following response earns 1 out of 2 points:  $\int_1^5 A(t) dt = 1502.147865 = 375.536966$ .

- The answer must be correct to three decimal places. For example,

$$\frac{1}{5-1} \int_1^5 A(t) dt = 375.536966 \approx 376 \text{ earns only the first point.}$$

- Degree mode: A response that presents answers obtained by using a calculator in degree mode does not earn the first point it would have otherwise earned. The response is generally eligible for all subsequent points (unless no answer is possible in degree mode or the question is made simpler by using degree mode). In degree mode,  $\frac{1}{4} \int_1^5 A(t) dt = 79.416068$ .

- Special case:  $\frac{1}{5} \int_1^5 A(t) dt = 300.429573$  earns 1 out of 2 points.

**Total for part (b) 2 points**

- (c) Is the rate at which vehicles arrive at the toll plaza at 6 A.M. ( $t = 1$ ) increasing or decreasing? Give a reason for your answer.

|                                                                                              |                    |                |
|----------------------------------------------------------------------------------------------|--------------------|----------------|
| $A'(1) = 148.947272$                                                                         | Considers $A'(1)$  | <b>1 point</b> |
| Because $A'(1) > 0$ , the rate at which the vehicles arrive at the toll plaza is increasing. | Answer with reason | <b>1 point</b> |

**Scoring notes:**

- The response need not present the value of  $A'(1)$ . The second line of the model solution earns both points.
- An incorrect value assigned to  $A'(1)$  earns the first point (but will not earn the second point).
- Without a reference to  $t = 1$ , the first point is earned by any of the following:
  - 148.947 accurate to the number of decimals presented, with zero up to three decimal places (i.e., 149, 148, 148.9, 148.95, or 148.94)
  - $A'(t) = 148.947$  by itself
- To be eligible for the second point, the first point must be earned.
- To earn the second point, there must be a reference to  $t = 1$ .
- Degree mode:  $A'(1) = 23.404311$

**Total for part (c) 2 points**

- (d) A line forms whenever  $A(t) \geq 400$ . The number of vehicles in line at time  $t$ , for  $a \leq t \leq 4$ , is given by  $N(t) = \int_a^t (A(x) - 400) dx$ , where  $a$  is the time when a line first begins to form. To the nearest whole number, find the greatest number of vehicles in line at the toll plaza in the time interval  $a \leq t \leq 4$ . Justify your answer.

|                                                                                                                                                                                                                                       |                       |                                   |     |   |     |           |   |           |        |         |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|-----------------------------------|-----|---|-----|-----------|---|-----------|--------|---------|
| $N'(t) = A(t) - 400 = 0$<br>$\Rightarrow A(t) = 400 \Rightarrow t = 1.469372, t = 3.597713$                                                                                                                                           | Considers $N'(t) = 0$ | 1 point                           |     |   |     |           |   |           |        |         |
|                                                                                                                                                                                                                                       | $t = a$ and $t = b$   | 1 point                           |     |   |     |           |   |           |        |         |
| $a = 1.469372$<br>$b = 3.597713$                                                                                                                                                                                                      |                       |                                   |     |   |     |           |   |           |        |         |
| <table><tr><td><math>t</math></td><td><math>N(t) = \int_a^t (A(x) - 400) dx</math></td></tr><tr><td><math>a</math></td><td>0</td></tr><tr><td><math>b</math></td><td>71.254129</td></tr><tr><td>4</td><td>62.338346</td></tr></table> | $t$                   | $N(t) = \int_a^t (A(x) - 400) dx$ | $a$ | 0 | $b$ | 71.254129 | 4 | 62.338346 | Answer | 1 point |
|                                                                                                                                                                                                                                       | $t$                   | $N(t) = \int_a^t (A(x) - 400) dx$ |     |   |     |           |   |           |        |         |
|                                                                                                                                                                                                                                       | $a$                   | 0                                 |     |   |     |           |   |           |        |         |
|                                                                                                                                                                                                                                       | $b$                   | 71.254129                         |     |   |     |           |   |           |        |         |
| 4                                                                                                                                                                                                                                     | 62.338346             |                                   |     |   |     |           |   |           |        |         |
|                                                                                                                                                                                                                                       | Justification         | 1 point                           |     |   |     |           |   |           |        |         |
| The greatest number of vehicles in line is 71.                                                                                                                                                                                        |                       |                                   |     |   |     |           |   |           |        |         |

**Scoring notes:**

- It is not necessary to indicate that  $A(t) = 400$  to earn the first point, although this statement alone would earn the first point.
- A response of “ $A(t) \geq 400$  when  $1.469372 \leq t \leq 3.597713$ ” will earn the first 2 points. A response of “ $A(t) \geq 400$ ” along with the presence of exactly one of the two numbers above will earn the first point, but not the second. A response of “ $A(t) \geq 400$ ” by itself will not earn either of the first 2 points.
- To earn the second point the values for  $a$  and  $b$  must be accurate to the number of decimals presented, with at least one and up to three decimal places. These may appear only in a candidates table, as limits of integration, or on a number line.
- A response with incorrect notation involving  $t$  or  $x$  is eligible to earn all 4 points.
- A response that does not earn the first point is still eligible for the remaining 3 points.
- To earn the third point, a response must present the greatest number of vehicles. This point is earned for answers of either 71 or 71.254\*\*\* only.
- A correct justification earns the fourth point, even if the third point is not earned because of a decimal presentation error.
- When using a Candidates Test, the response must include the values for  $N(a)$ ,  $N(b)$ , and  $N(4)$  to earn the fourth point. These values must be correct to the number of decimals presented, with up to three decimal places. (Correctly rounded integer values are acceptable.)
- Alternate solution for the third and fourth points:  
 For  $a \leq t \leq b$ ,  $A(t) \geq 400$ . For  $b \leq t \leq 4$ ,  $A(t) \leq 400$ .  
 Thus,  $N(t) = \int_a^t (A(x) - 400) dx$  is greatest at  $t = b$ .  
 $N(b) = 71.254129$ , and the greatest number of vehicles in line is 71.
- Degree mode: The response is only eligible to earn the first point because in degree mode  $A(t) < 400$ .

---

**Total for part (d)      4 points**

---

**Total for question 1      9 points**