7.7 Calculating Equilibrium Concentrations

Name: _____ Class: _____ Date: _____ **Total: 12 marks**

KEY WORDS ICE table ICE • small-x approximation • equilibrium concentration • unknown • valid • percent

Objective

Build the skills to answer exam questions on **calculating equilibrium concentrations** : ICE tables with x, and the small-x approximation when K is small.

You must be able to:

- set up an ICE table with an unknown change x
- write the K expression in terms of x
- use the **small-x approximation** when K is very small, and check it (less than 5%)
- find each equilibrium concentration from x

1 Key knowledge and worked examples

■ Solving for x

step	what to do
1	ICE table: the changes are $-x$, $+x$, $+2x$, etc., from the coefficients
2	put the equilibrium row into the K expression
3	if K is very small, $c - x \approx c$ (this makes the algebra easy)
4	solve for x, then find every concentration
5	check: $\dfrac{x}{c} \times 100\%$ should be under 5%

Worked example: a weak dissociation. $\mathrm{AB} \rightleftharpoons \mathrm{A} + \mathrm{B}$, $K_c = 4.0 \times 10^{-6}$, $[\mathrm{AB}]_0 = 0.10$ M.

1. $\dfrac{x^2}{0.10 - x} \approx \dfrac{x^2}{0.10} = 4.0 \times 10^{-6}$, so $x^2 = 4.0 \times 10^{-7}$ and $x = 6.3 \times 10^{-4}$ M.
2. Check: $\dfrac{6.3 \times 10^{-4}}{0.10} = 0.63\%$, under 5%: **the approximation is fine**.

exercises.lol exercises.com.cn


> **Answer shapes:** show the ICE table, the expression in x, the approximation and the check.

2 Practice

2.1 0.50 M of A is placed in a flask: $\mathrm{A(g)} \rightleftharpoons 2\mathrm{B(g)}$. Write the equilibrium concentrations of A and B in terms of x, and the K_c expression in terms of x. [2]

	A	B
I	0.50	0
C		
E		

2.2 Explain why $0.20 - x$ can be replaced by 0.20 when K_c is very small. [2]



2.3 Solve $\dfrac{x^2}{0.10} = 1.0 \times 10^{-5}$ for x. [2]




exercises.lol exercises.com.cn

3 Exam-style questions

3.1 0.200 mol of PCl_5 is placed in a 1.00 L flask: $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$, $K_c = 1.8 \times 10^{-5}$ at this temperature.

(a) Write the equilibrium concentrations in terms of x, and the K_c expression in terms of x. [2]



(b) Use the small-x approximation to find $[Cl_2]$ at equilibrium. [2]

(c) Show that the approximation is valid. [2]

exercises.lol exercises.com.cn