

Past-paper walk-through

Introduction to Entropy ■ 9.1

eXercise

Questions in this set

- 2025 Q3
- 2024 Q2
- 2019 Q1

Reading the mark scheme

- **B1** a mark that stands on its own – the point is either made or not.
- **M1** a *method* mark: the right approach, even if the number is wrong.
- **A1** an *answer* mark, and it usually needs its M mark first.
- **C1** a working mark on the way to the answer.
- **//** separates answers the examiner accepts equally.
- **ecf** = error carried forward: a later part is marked on your own earlier answer.
- **owtte** = “or words to that effect” – the wording need not match.

Question 3

2025 ■ Q3


White phosphorus is composed of P_4 molecules with a tetrahedral structure, as shown in the diagram on the left. Each P atom is bonded to the other three P atoms by single bonds, as shown in the incomplete Lewis diagram on the right.

[Left: a ball-and-stick tetrahedral structure of P_4 , showing four P atoms (spheres) at the vertices of a tetrahedron connected by bonds (sticks).]

[Right: an incomplete Lewis diagram showing four P atoms arranged in a tetrahedral projection (one P at top, one P in the middle/front, two P at the bottom left and right), connected by single bonds forming the P_4 skeleton, with no lone pairs drawn yet.]

(a) In the box in part A, complete the Lewis diagram for P_4 by drawing the nonbonding electrons.


Question 3



Question 3

Mass of P_4O_{10}	0.100 g
Mass of H_2O	100.0 g
Initial temperature	22.00° C
Final temperature	22.38° C
Molar mass of P_4O_{10}	283.9 g/mol
Specific heat of H_2O	4.18 J/(g·°C)

Question 3



Question 3


2025 ■ Q3

White phosphorus is composed of P_4 molecules with a tetrahedral structure, as shown in the diagram on the left. Each P atom is bonded to the other three P atoms by single bonds, as shown in the incomplete Lewis diagram on the right.

[Left: a ball-and-stick tetrahedral structure of P_4 , showing four P atoms (spheres) at the vertices of a tetrahedron connected by bonds (sticks).]

[Right: an incomplete Lewis diagram showing four P atoms arranged in a tetrahedral projection (one P at top, one P in the middle/front, two P at the bottom left and right), connected by single bonds forming the P_4 skeleton, with no lone pairs drawn yet.]

(b) The reaction of white phosphorus with oxygen to form $P_4O_{10}(s)$ is thermodynamically favorable at 298 K. The reaction is represented by equation 1.



Question 3

(b)(i) The entropy change of the reaction, ΔS° , is negative. Using particle-level reasoning, explain why the entropy decreases as the reaction progresses.

(b)(ii) The enthalpy change of the reaction, ΔH° , is also negative. A student claims that the favorability of the reaction is driven by enthalpy and **not** by entropy. Is the student's claim correct? Justify your answer by using the relationship between ΔG° , ΔH° , and ΔS° .

Question 3


2025 ■ Q3

White phosphorus is composed of P_4 molecules with a tetrahedral structure, as shown in the diagram on the left. Each P atom is bonded to the other three P atoms by single bonds, as shown in the incomplete Lewis diagram on the right.

[Left: a ball-and-stick tetrahedral structure of P_4 , showing four P atoms (spheres) at the vertices of a tetrahedron connected by bonds (sticks).]

[Right: an incomplete Lewis diagram showing four P atoms arranged in a tetrahedral projection (one P at top, one P in the middle/front, two P at the bottom left and right), connected by single bonds forming the P_4 skeleton, with no lone pairs drawn yet.]

(c) $P_4O_{10}(s)$ reacts exothermically with water to form phosphoric acid, as represented by equation 2.



Question 3

A chemist uses a calorimetry experiment to determine the enthalpy change for the reaction, as represented by the following diagram.

[Calorimeter diagram: a coffee-cup-style calorimeter labeled "Thermometer" (inserted through the insulating cover), "Stirrer" (inserted alongside the thermometer), "Insulating Cover" (the lid), "Inner Vessel" (the cup holding the solution, shaded), and "Outer Vessel" (the outer supporting cup).]

The chemist carries out the calorimetry experiment and records the following information.

Quantity	Value	—	—	Mass of P_4O_{10}	0.100 g	Mass of H_2O	100.0 g
Initial temperature	22.00°C	Final temperature	22.38°C	Molar mass of P_4O_{10}	283.9 g/mol	Specific heat of H_2O	4.18 J/(g · °C)

(c)(i) Calculate the amount of heat, q , released during the experiment, in kJ. Assume that the specific heat of the solution is the same as that of water.

Question 3

(c)(ii) Calculate the value of $\Delta H_{\text{rxn}}^{\circ}$ for equation 2 in $\text{kJ/mol}_{\text{rxn}}$. Include the sign in your answer.

Question 3

2025 ■ Q3

White phosphorus is composed of P_4 molecules with a tetrahedral structure, as shown in the diagram on the left. Each P atom is bonded to the other three P atoms by single bonds, as shown in the incomplete Lewis diagram on the right.

[Left: a ball-and-stick tetrahedral structure of P_4 , showing four P atoms (spheres) at the vertices of a tetrahedron connected by bonds (sticks).]

[Right: an incomplete Lewis diagram showing four P atoms arranged in a tetrahedral projection (one P at top, one P in the middle/front, two P at the bottom left and right), connected by single bonds forming the P_4 skeleton, with no lone pairs drawn yet.]

(d) The chemist weighed out 0.100 g P_4O_{10} of and 100.0 g of H_2O to perform a second trial. In the second trial, some of the solid P_4O_{10} stuck to the weighing paper and was not transferred to the calorimeter. Given that P_4O_{10} is the limiting reactant,

Question 3

would ΔT for the second trial be greater than, less than, or equal to the value in the first trial? Justify your answer.

$\text{P}_4(\text{s})$ also reacts readily with $\text{Cl}_2(\text{g})$ to produce phosphorus trichloride, $\text{PCl}_3(\text{g})$, which in turn reacts with $\text{Cl}_2(\text{g})$ in an equilibrium process to produce $\text{PCl}_5(\text{g})$. The reactions are represented by equations 3 and 4.



Question 3


2025 ■ Q3

White phosphorus is composed of P_4 molecules with a tetrahedral structure, as shown in the diagram on the left. Each P atom is bonded to the other three P atoms by single bonds, as shown in the incomplete Lewis diagram on the right.

[Left: a ball-and-stick tetrahedral structure of P_4 , showing four P atoms (spheres) at the vertices of a tetrahedron connected by bonds (sticks).]

[Right: an incomplete Lewis diagram showing four P atoms arranged in a tetrahedral projection (one P at top, one P in the middle/front, two P at the bottom left and right), connected by single bonds forming the P_4 skeleton, with no lone pairs drawn yet.]

(e) Calculate the standard enthalpy of formation of $PCl_5(g)$ represented by equation 5.



The following particle-level diagram represents the contents of the vessel in an equilibrium mixture at 546 K involving equation 4.

Question 3

[Particle diagram: a box containing multiple particles at random positions representing an equilibrium mixture; a key indicates two-atom particles (grey-grey) = Cl_2 , four-atom tetrahedral-looking particles (grey with three white) = PCl_3 , and six-atom particles (grey with five white) = PCl_5 . The box shows several Cl_2 particles (pairs of grey circles), several PCl_3 particles (grey center with three white circles), and several PCl_5 particles (grey center with five white circles) scattered throughout.]

Question 3

2025 ■ Q3

White phosphorus is composed of P_4 molecules with a tetrahedral structure, as shown in the diagram on the left. Each P atom is bonded to the other three P atoms by single bonds, as shown in the incomplete Lewis diagram on the right.

[Left: a ball-and-stick tetrahedral structure of P_4 , showing four P atoms (spheres) at the vertices of a tetrahedron connected by bonds (sticks).]

[Right: an incomplete Lewis diagram showing four P atoms arranged in a tetrahedral projection (one P at top, one P in the middle/front, two P at the bottom left and right), connected by single bonds forming the P_4 skeleton, with no lone pairs drawn yet.]

(f) Equation 4 for the reaction that occurs is shown.



(f)(i) If each particle in the diagram represents a partial pressure of 1.00 atm, what is the value of K_p for the equilibrium mixture at 546 K?

Question 3

(f)(ii) Does the value of K_p increase, decrease, or remain the same when the temperature is increased to 596 K? Justify your answer based on ΔH_2° .

Solution 3

(a)

Mark scheme

- Complete Lewis diagram for P_4 : the four P atoms form a tetrahedron, each bonded to the other three P atoms by single bonds, and each P atom bears one lone (nonbonding) pair of electrons — giving each P three bonding pairs plus one lone pair, satisfying the octet rule.

Solution 3

(b)

Mark scheme

- No separate response required — this is the context/stem presenting equation 1 ($\text{P}_4(\text{s}) + 5 \text{O}_2(\text{g}) \rightarrow \text{P}_4\text{O}_{10}(\text{s})$), thermodynamically favorable at 298 K, that parts (i) and (ii) below refer to.

(b)(i)

Mark scheme

- Because gas particles are more dispersed (have more accessible microstates) than solids, the entropy decreases as the reactants (which include a gas, O_2) convert into the solid product (P_4O_{10}).

Solution 3

(b)(ii)

Mark scheme

- Yes, the student's claim is correct. Given $\Delta G_{rxn}^{\circ} = \Delta H_{rxn}^{\circ} - T\Delta S_{rxn}^{\circ}$, the reaction must have $\Delta G_{rxn}^{\circ} < 0$ to be favorable. Because the reaction is exothermic, $\Delta H_{rxn}^{\circ} < 0$ and enthalpy contributes to favorability. $\Delta S_{rxn}^{\circ} < 0$, so entropy does NOT contribute to favorability — i.e., enthalpy alone drives it.

Solution 3

(c)

Mark scheme

- No separate response required — this is the context/stem describing the calorimetry experiment for the exothermic reaction of $\text{P}_4\text{O}_{10}(\text{s})$ with water (equation 2: $\text{P}_4\text{O}_{10}(\text{s}) + 6 \text{H}_2\text{O}(\text{l}) \rightarrow 4 \text{H}_3\text{PO}_4(\text{aq})$) that parts (i) and (ii) below use.

Solution 3

(c)(i)

Mark scheme

- $q = mc\Delta T = (100.1 \text{ g})(4.18 \text{ J/(g}\cdot^{\circ}\text{C)})(22.38^{\circ}\text{C} - 22.00^{\circ}\text{C}) = 160 \text{ J} = 0.16 \text{ kJ}$
(reported to the correct number of significant figures).

Solution 3

(c)(ii)

Mark scheme

■ $q_{rxn} = -q_{surr} = -0.16 \text{ kJ}.$

$$\Delta H_{rxn}^{\circ} = \frac{-0.16 \text{ kJ}}{0.100 \text{ g P}_4\text{O}_{10}} \times \frac{283.9 \text{ g P}_4\text{O}_{10}}{1 \text{ mol P}_4\text{O}_{10}} \times \frac{1 \text{ mol P}_4\text{O}_{10}}{1 \text{ mol}_{rxn}} = -450 \text{ kJ/mol}_{rxn}.$$

The correct (negative) sign must be included.

Solution 3

(d)

Mark scheme

- Less than. If less P_4O_{10} is present (some stuck to the weighing paper), less thermal energy will be transferred to the water during the reaction, causing the temperature increase (ΔT) to be less than it was with the full 0.100 g of P_4O_{10} .

Solution 3

(e)

Mark scheme

- Using Hess's law:

$$\Delta H_{f,\text{PCl}_5(g)}^{\circ} = \frac{1}{4}\Delta H_1^{\circ} + \Delta H_2^{\circ} = \frac{1}{4}(-1148) + (-88) = -375 \text{ kJ/mol.}$$

Solution 3

(f)

Mark scheme

- No separate response required — this is the context/stem presenting equation 4 ($\text{PCl}_3(g) + \text{Cl}_2(g) \rightleftharpoons \text{PCl}_5(g)$, $\Delta H_2^\circ = -88 \text{ kJ/mol}_{\text{rxn}}$) that parts (i) and (ii) below refer to.

Solution 3

(f)(i)

Mark scheme

- Reading partial pressures of 1.00 atm per particle from the diagram (4 PCl_5 , 2 PCl_3 , 6 Cl_2 particles): $K_p = \frac{P_{\text{PCl}_5}}{(P_{\text{PCl}_3})(P_{\text{Cl}_2})} = \frac{4.00}{(2.00)(6.00)} = \frac{1}{3} = 0.333$.

Solution 3


(f)(ii)

Mark scheme

- Decrease. The negative value of ΔH_2° indicates the reaction is exothermic. Because exothermic reactions favor reactant formation at higher temperature (the equilibrium shifts toward reactants as heat is added), the value of K_p decreases.

Question 2

2024 ■ Q2



2. A chemical reaction between maleic acid ($\text{H}_2\text{C}_4\text{H}_2\text{O}_4$) and sodium bicarbonate (NaHCO_3) occurs in the presence of water to produce carbon dioxide and sodium maleate ($\text{Na}_2\text{C}_4\text{H}_2\text{O}_4$), as represented by the following equation.

(a)(i) A student combines equal masses of $\text{H}_2\text{C}_4\text{H}_2\text{O}_4(s)$ chunks and $\text{NaHCO}_3(s)$ chunks with sufficient water at 20.0°C . The student determines that 0.0114 mol of $\text{CO}_2(g)$ is produced after the reaction goes to completion.

Calculate the number of grams of $\text{CO}_2(g)$ produced.

(a)(ii) The $\text{CO}_2(g)$ produced from the reaction at 20.0°C was collected and found to have a pressure of 1.25 atm . Calculate the volume of $\text{CO}_2(g)$, in liters.

Question 2

2024 ■ Q2



2. A chemical reaction between maleic acid ($\text{H}_2\text{C}_4\text{H}_2\text{O}_4$) and sodium bicarbonate (NaHCO_3) occurs in the presence of water to produce carbon dioxide and sodium maleate ($\text{Na}_2\text{C}_4\text{H}_2\text{O}_4$), as represented by the following equation.

(b)(i) The student performs a second experiment that is identical to the first except that the student grinds the chunks of $\text{H}_2\text{C}_4\text{H}_2\text{O}_4(s)$ and $\text{NaHCO}_3(s)$ into powder before combining the powder with water.

What happens to the surface area of the reactants when the student grinds the chunks into powder?

Question 2

(b)(ii) The rate-determining step for the overall reaction is the dissolving of the solids. Would the time required for the dissolving of the solids in the second experiment be longer than, shorter than, or the same as the time required in the first experiment?

Justify your answer based on the collisions between particles.

(b)(iii) When the reaction is complete, will the volume of $\text{CO}_2(g)$ at the end of the second experiment be greater than, less than, or equal to the volume at the end of the first experiment? Justify your answer.

Question 2

2024 ■ Q2



2. A chemical reaction between maleic acid ($\text{H}_2\text{C}_4\text{H}_2\text{O}_4$) and sodium bicarbonate (NaHCO_3) occurs in the presence of water to produce carbon dioxide and sodium maleate ($\text{Na}_2\text{C}_4\text{H}_2\text{O}_4$), as represented by the following equation.

(c) The student conducts additional trials of the experiment and produces the following data table.

Trial	Mass of $\text{H}_2\text{C}_4\text{H}_2\text{O}_4$ (grams)	Mass of NaHCO_3 (grams)	Moles of CO_2 Produced (mol)
3	1.543	1.251	0.01489
4	1.543	1.686	0.02007

Based on the student's data, identify the limiting reactant in trial 3. Justify your answer.

Question 2

2024 ■ Q2



2. A chemical reaction between maleic acid ($\text{H}_2\text{C}_4\text{H}_2\text{O}_4$) and sodium bicarbonate (NaHCO_3) occurs in the presence of water to produce carbon dioxide and sodium maleate ($\text{Na}_2\text{C}_4\text{H}_2\text{O}_4$), as represented by the following equation.

(d) The reaction has a value of ΔS° greater than zero. Using particle-level reasoning, explain why the entropy increases as the reaction progresses.

Question 2

2024 ■ Q2



2. A chemical reaction between maleic acid ($\text{H}_2\text{C}_4\text{H}_2\text{O}_4$) and sodium bicarbonate (NaHCO_3) occurs in the presence of water to produce carbon dioxide and sodium maleate ($\text{Na}_2\text{C}_4\text{H}_2\text{O}_4$), as represented by the following equation.

(e) The student notices that the temperature of the reaction mixture decreases as the reaction takes place and correctly determines that the reaction is endothermic.

The student claims that the reaction is thermodynamically favorable at all temperatures because $\Delta S_{\text{rxn}}^{\circ} > 0$ and the reaction is endothermic. Do you agree or disagree with the student's claim? Justify your answer.



Question 2

2024 ■ Q2



2. A chemical reaction between maleic acid ($\text{H}_2\text{C}_4\text{H}_2\text{O}_4$) and sodium bicarbonate (NaHCO_3) occurs in the presence of water to produce carbon dioxide and sodium maleate ($\text{Na}_2\text{C}_4\text{H}_2\text{O}_4$), as represented by the following equation.

(f) Next, the student investigates the acid-base behavior of maleic acid. The student notes that maleic acid is a diprotic acid. The two acid dissociation processes that occur are represented by the following equations.



Question 2

Calculate the $\text{p}K_{\text{a}2}$ value for the $\text{HC}_4\text{H}_2\text{O}_4^-$ ion.

Question 2

2024 ■ Q2



2. A chemical reaction between maleic acid ($\text{H}_2\text{C}_4\text{H}_2\text{O}_4$) and sodium bicarbonate (NaHCO_3) occurs in the presence of water to produce carbon dioxide and sodium maleate ($\text{Na}_2\text{C}_4\text{H}_2\text{O}_4$), as represented by the following equation.

(g) A buffer solution with a pH of 7.00 is prepared using $\text{C}_4\text{H}_2\text{O}_4^{2-}$ and $\text{HC}_4\text{H}_2\text{O}_4^-$. Calculate the ratio

$$\frac{[\text{C}_4\text{H}_2\text{O}_4^{2-}]}{[\text{HC}_4\text{H}_2\text{O}_4^-]}$$

in this solution.

Solution 2

(a)(i)**Mark scheme**

$$\blacksquare 0.0114 \text{ mol CO}_2 \times \frac{44.01 \text{ g}}{1 \text{ mol}} = 0.502 \text{ g CO}_2.$$

Solution 2

(a)(ii)

Mark scheme

- Using $PV = nRT$:

$$V = \frac{nRT}{P} = \frac{(0.0114 \text{ mol})(0.08206 \text{ L} \cdot \text{atm}/(\text{mol} \cdot \text{K}))(293 \text{ K})}{1.25 \text{ atm}} = 0.219 \text{ L.}$$

Solution 2

(b)(i)

Mark scheme

- Claim: the surface area of the solid reactants increases when the chunks are ground into powder.

(b)(ii)

Mark scheme

- Shorter than. The powdered solids have a larger surface area than the solid chunks, so collisions between water molecules and the solid surface particles occur more frequently, resulting in a faster rate of dissolution and a shorter time to dissolve the solids.

Solution 2

(b)(iii)

Mark scheme

- Equal to. Both experiments begin with the same amount (mass) of reactants, so they produce the same number of moles of $\text{CO}_2(g)$ under the same conditions of pressure and temperature; therefore the final volume of CO_2 will be the same regardless of particle size (rate changes, but the final amount/volume does not).

Solution 2

(c) Mark scheme

- NaHCO_3 is the limiting reactant (accept either justification): changing the mass of NaHCO_3 alters the amount of CO_2 produced, OR NaHCO_3 has the smaller theoretical yield of CO_2 :

$$1.543 \text{ g H}_2\text{C}_4\text{H}_2\text{O}_4 \times \frac{1 \text{ mol}}{116.07 \text{ g}} \times \frac{2 \text{ mol CO}_2}{1 \text{ mol acid}} = 0.02659 \text{ mol CO}_2 \text{ (if acid were limiting)}$$

$$\text{vs. } 1.251 \text{ g NaHCO}_3 \times \frac{1 \text{ mol}}{84.01 \text{ g}} \times \frac{2 \text{ mol CO}_2}{2 \text{ mol NaHCO}_3} = 0.01489 \text{ mol CO}_2$$

(if NaHCO_3 limiting) — the smaller value (NaHCO_3) is the actual limiting reactant, matching the trial's observed 0.01489 mol CO_2 .

Solution 2

(d)

Mark scheme

- The entropy change is positive because the aqueous/solid reactants produce 2 moles of gas (CO_2) particles, according to the balanced equation. Gases are far more dispersed (occupy many more microstates) than condensed (solid/aqueous) phases, so the entropy of the products is greater than that of the reactants.

Solution 2

(e)

Mark scheme

- Disagree. Accept either justification: (1) The reaction is endothermic ($\Delta H > 0$) with a positive entropy change ($\Delta S > 0$), so it is only thermodynamically favorable at a high enough temperature where the magnitude of $-T\Delta S$ exceeds ΔH — NOT at all temperatures. (2) For a reaction to be thermodynamically favorable ($\Delta G < 0$) at ALL temperatures, it must be exothermic ($\Delta H < 0$) AND have $\Delta S > 0$ — this reaction is endothermic, so it fails that requirement.

Solution 2

(f)

Mark scheme

- Given $K_{a_2} = 8.5 \times 10^{-7}$ for the second dissociation of maleic acid:
 $\text{p}K_{a_2} = -\log(8.5 \times 10^{-7}) = 6.07.$

Solution 2

(g)

Mark scheme

- Using the Henderson–Hasselbalch equation: $\text{pH} = \text{p}K_{a_2} + \log \frac{[\text{C}_4\text{H}_2\text{O}_4^{2-}]}{[\text{HC}_4\text{H}_2\text{O}_4^-]}$.

Solving: $\frac{[\text{C}_4\text{H}_2\text{O}_4^{2-}]}{[\text{HC}_4\text{H}_2\text{O}_4^-]} = 10^{(\text{pH} - \text{p}K_{a_2})} = 10^{(7.00 - 6.07)} = 8.5$.

Question 1

2019 ■ Q1

[Complete Lewis electron-dot diagram of urea: $\text{H}-\text{N}(\text{H})-\text{C}(=\text{O})-\text{N}(\text{H})-\text{H}$, with the carbon double-bonded to oxygen (two lone pairs on O), and each nitrogen bonded to two H atoms and bearing one lone pair, and the central carbon bonded to both nitrogens]

The compound urea, H_2NCONH_2 , is widely used in chemical fertilizers. The complete Lewis electron-dot diagram for the urea molecule is shown above.

(a) Identify the hybridization of the valence orbitals of the carbon atom in the urea molecule.

Question 1



Polystyrene
Cup




Thermometer



Stirring
Rod



Bottle of
Urea





Balance



Distilled
Water

Question 1



Question 1

	S° (J/(mol·K))
$\text{H}_2\text{NCONH}_2(s)$	104.6
$\text{H}_2\text{NCONH}_2(aq)$	?

Question 1

2019 ■ Q1

[Complete Lewis electron-dot diagram of urea: $\text{H}-\text{N}(\text{H})-\text{C}(=\text{O})-\text{N}(\text{H})-\text{H}$, with the carbon double-bonded to oxygen (two lone pairs on O), and each nitrogen bonded to two H atoms and bearing one lone pair, and the central carbon bonded to both nitrogens]


The compound urea, H_2NCONH_2 , is widely used in chemical fertilizers. The complete Lewis electron-dot diagram for the urea molecule is shown above.

(b) Urea has a high solubility in water, due in part to its ability to form hydrogen bonds. A urea molecule and four water molecules are represented in the box below. Draw ONE dashed line (—) to indicate a possible location of a hydrogen bond between a water molecule and the urea molecule.

[Diagram: a box containing one urea molecule (drawn with spheres: white = O, gray = H, black = C, hatched = N) surrounded by four separate water molecules, each

Question 1

water molecule drawn as one white (O) sphere bonded to two gray (H) spheres;
legend: white circle = O, gray circle = H, black circle = C, hatched circle = N]



The dissolution of urea is represented by the equation above. A student determines that 5.39 grams of H_2NCONH_2 (molar mass 60.06 g/mol) can dissolve in water to make 5.00 mL of a saturated solution at 20.°C.

Question 1

2019 ■ Q1

[Complete Lewis electron-dot diagram of urea: $\text{H}-\text{N}(\text{H})-\text{C}(=\text{O})-\text{N}(\text{H})-\text{H}$, with the carbon double-bonded to oxygen (two lone pairs on O), and each nitrogen bonded to two H atoms and bearing one lone pair, and the central carbon bonded to both nitrogens]

The compound urea, H_2NCONH_2 , is widely used in chemical fertilizers. The complete Lewis electron-dot diagram for the urea molecule is shown above.

(c) Calculate the concentration of urea, in mol/L, in the saturated solution at $20.^{\circ}\text{C}$.

Question 1

2019 ■ Q1

[Complete Lewis electron-dot diagram of urea: $\text{H}-\text{N}(\text{H})-\text{C}(=\text{O})-\text{N}(\text{H})-\text{H}$, with the carbon double-bonded to oxygen (two lone pairs on O), and each nitrogen bonded to two H atoms and bearing one lone pair, and the central carbon bonded to both nitrogens]

The compound urea, H_2NCONH_2 , is widely used in chemical fertilizers. The complete Lewis electron-dot diagram for the urea molecule is shown above.

(d) The student also determines that the concentration of urea in a saturated solution at 25°C is 19.8 M . Based on this information, is the dissolution of urea endothermic or exothermic? Justify your answer in terms of Le Chatelier's principle.

[Diagram: row of laboratory equipment icons, labelled left to right: Polystyrene Cup, Thermometer, Stirring Rod, Bottle of Urea, Balance, Distilled Water]

Question 1

2019 ■ Q1

[Complete Lewis electron-dot diagram of urea: $\text{H—N(H)—C(=O)—N(H)—H}$, with the carbon double-bonded to oxygen (two lone pairs on O), and each nitrogen bonded to two H atoms and bearing one lone pair, and the central carbon bonded to both nitrogens]

The compound urea, H_2NCONH_2 , is widely used in chemical fertilizers. The complete Lewis electron-dot diagram for the urea molecule is shown above.

(e) The equipment shown above is provided so that the student can determine the value of the molar heat of solution for urea. Knowing that the specific heat of the solution is $4.18 \text{ J/(g}\cdot^\circ\text{C)}$, list the specific measurements that are required to be made during the experiment.

$S^\circ \text{ (J/(mol}\cdot\text{K))}$	—	—	$\text{H}_2\text{NCONH}_2(\text{s})$	104.6	$\text{H}_2\text{NCONH}_2(\text{aq})$	?
--	---	---	--------------------------------------	-------	---------------------------------------	---

Question 1

2019 ■ Q1

[Complete Lewis electron-dot diagram of urea: $\text{H}-\text{N}(\text{H})-\text{C}(=\text{O})-\text{N}(\text{H})-\text{H}$, with the carbon double-bonded to oxygen (two lone pairs on O), and each nitrogen bonded to two H atoms and bearing one lone pair, and the central carbon bonded to both nitrogens]

The compound urea, H_2NCONH_2 , is widely used in chemical fertilizers. The complete Lewis electron-dot diagram for the urea molecule is shown above.

(f) The entropy change for the dissolution of urea, $\Delta S_{\text{soln}}^\circ$, is $70.1 \text{ J}/(\text{mol}\cdot\text{K})$ at 25°C . Using the information in the table above, calculate the absolute molar entropy, S° , of aqueous urea.

Question 1

2019 ■ Q1

[Complete Lewis electron-dot diagram of urea: $\text{H}-\text{N}(\text{H})-\text{C}(=\text{O})-\text{N}(\text{H})-\text{H}$, with the carbon double-bonded to oxygen (two lone pairs on O), and each nitrogen bonded to two H atoms and bearing one lone pair, and the central carbon bonded to both nitrogens]

The compound urea, H_2NCONH_2 , is widely used in chemical fertilizers. The complete Lewis electron-dot diagram for the urea molecule is shown above.

(g) Using particle-level reasoning, explain why $\Delta S_{\text{soln}}^\circ$ is positive for the dissolution of urea in water.

Question 1

2019 ■ Q1

[Complete Lewis electron-dot diagram of urea: $\text{H}-\text{N}(\text{H})-\text{C}(=\text{O})-\text{N}(\text{H})-\text{H}$, with the carbon double-bonded to oxygen (two lone pairs on O), and each nitrogen bonded to two H atoms and bearing one lone pair, and the central carbon bonded to both nitrogens]

The compound urea, H_2NCONH_2 , is widely used in chemical fertilizers. The complete Lewis electron-dot diagram for the urea molecule is shown above.

(h) The student claims that ΔS° for the process contributes to the thermodynamic favorability of the dissolution of urea at 25°C . Use the thermodynamic information above to support the student's claim.

Solution 1

(a)

Mark scheme

- sp^2 . The carbon atom in urea is bonded to three groups (two N atoms and one O atom, one bond of which is a double bond) with no lone pairs, giving trigonal-planar sp^2 hybridization. 1 point for stating sp^2 .

Solution 1

(b)

Mark scheme

- A dashed line should connect a hydrogen atom of a water molecule to a nitrogen or oxygen atom of urea, OR an oxygen atom of a water molecule to a hydrogen atom of urea (an H-bond donor H drawn near an H-bond acceptor lone pair). 1 point is earned for any correctly placed dashed line matching this pattern.

Solution 1

(c)

Mark scheme

- Moles of urea: $5.39 \text{ g H}_2\text{NCONH}_2 \times (1 \text{ mol} / 60.06 \text{ g}) = 0.0897 \text{ mol}$ (1 point, may be implicit). Molarity: $0.0897 \text{ mol} / 0.00500 \text{ L} = 17.9 \text{ M}$ (1 point for the correct molarity).

Solution 1

(d)

Mark scheme

- Endothermic. The increased solubility at the higher temperature (19.8 M at 25°C vs. 17.9 M at 20.°C) implies the dissolution of urea is endothermic. By Le Chatelier's principle: heating a saturated solution stresses the equilibrium, and that stress is counteracted by more urea dissolving (the endothermic direction is favored). 1 point for the correct answer (endothermic) with this justification.

Solution 1

(e)

Mark scheme

- To determine the molar heat of solution calorimetrically, the student must measure: the mass of urea and the mass of water used (1 point); and the initial temperature of the water and the final temperature of the resulting solution (1 point). With the given specific heat $4.18 \text{ J}/(\text{g} \cdot ^\circ\text{C})$, these give $q = mc\Delta T$, which is then divided by moles of urea for the molar heat of solution.

Solution 1

(f)

Mark scheme

- $\Delta S^{\circ}_{\text{soln}} = S^{\circ}(\text{H}_2\text{NCONH}_2(\text{aq})) - S^{\circ}(\text{H}_2\text{NCONH}_2(\text{s})); 70.1 \text{ J}/(\text{mol} \cdot \text{K}) = S^{\circ}(\text{aq}) - 104.6 \text{ J}/(\text{mol} \cdot \text{K}); S^{\circ}(\text{H}_2\text{NCONH}_2(\text{aq})) = 174.7 \text{ J}/(\text{mol} \cdot \text{K}).$ 1 point for the correct answer.

Solution 1

(g)

Mark scheme

- Urea molecules in aqueous solution have a much greater number of possible arrangements/microstates (free to move and interact throughout the solution) than in the ordered solid, so dissolution increases the number of accessible microstates, giving a positive $\Delta S^{\circ}_{\text{soln}}$. 1 point for a correct particle-level explanation along these lines.

Solution 1

(h)

Mark scheme

- Thermodynamic favorability at standard conditions is determined by the sign of ΔG° , where $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$. Since ΔS° is positive, the $-T\Delta S^\circ$ term is negative, making ΔG° smaller (more negative) and thus making the dissolution more thermodynamically favorable — supporting the student's claim. 1 point for this reasoning.