

9.7 Coupled Reactions

Name: _____ Class: _____ Date: _____

Total: 9 marks

Objective

Build the skills to answer exam questions on **coupled reactions**.

You must be able to:

- add a favorable reaction to an unfavorable one to drive it
- add the ΔG values
- decide whether the coupled reaction is favorable overall

1 Worked examples

Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

■ Coupling reactions

An **unfavorable** reaction ($\Delta G > 0$) can be driven by **coupling** it to a **favorable** one ($\Delta G < 0$) that shares a species, if the total ΔG is negative.

■ Adding the ΔG values

Like Hess's law, when reactions are added their ΔG values add:

$$\Delta G_{\text{total}} = \Delta G_1 + \Delta G_2.$$

■ A worked coupling

Reaction 1: $\Delta G = +30$ kJ (unfavorable). Reaction 2: $\Delta G = -50$ kJ (favorable).
Coupled: $\Delta G = +30 + (-50) = -20$ kJ — now favorable.

■ Biological example

Cells couple the very favorable hydrolysis of ATP ($\Delta G \ll 0$) to otherwise-unfavorable reactions to make them go — the basis of metabolism.

2 Practice

Now apply the methods above.

2.1 When two reactions are added, how do their ΔG values combine? [1]2.2 Reaction A (+20 kJ) is coupled to B (−35 kJ). Find the total ΔG . [1]

2.3 Is the coupled reaction in 2.2 favorable? [1]

3 Exam-style questions

3.1 An unfavorable reaction can be driven by coupling it to a reaction with [1]

A	B
C	D

A a larger positive ΔG B a sufficiently negative ΔG C zero ΔG

D a higher activation energy

3.2 Reaction 1 ($\Delta G = +25$ kJ) is coupled with reaction 2 ($\Delta G = -40$ kJ).(a) Find the total ΔG . [2]

(b) State whether the coupled process is favorable. [1]

3.3 Explain how cells use ATP hydrolysis to drive unfavorable reactions. [2]