


5. Hydrogen gas and iodine gas react to form hydrogen iodide at an elevated temperature, as represented by the following equation.



(a) Write the expression for the equilibrium constant, K_c , for this reaction.

(b) $\text{H}_2(g)$ and $\text{I}_2(g)$ are added to a previously evacuated container and allowed to react.


- (i) At a certain time, the value of the reaction quotient, Q , is 0.67. The following particle diagram is an incomplete representation of the system at this time. The diagram shows the relative number of $\text{H}_2(g)$ and $\text{I}_2(g)$ molecules, but the $\text{HI}(g)$ molecules are not included. Draw the number of $\text{HI}(g)$ molecules needed to complete the diagram so that it accurately represents the system.



GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 5** on this page.

- (ii) A student monitors the number of moles of $\text{HI}(g)$ over time. Hypothesize an experimental change that could have been applied to the system in the rigid container at time t to result in the change in the number of moles of $\text{HI}(g)$ shown in the graph. Assume that the student did not add more $\text{HI}(g)$ to the system.



- (iii) After equilibrium is established, the mixture is transferred to a larger container at constant temperature. As a result, would the number of moles of $\text{HI}(g)$ increase, decrease, or remain the same? Justify your answer.

GO ON TO THE NEXT PAGE.