


1. Answer the following questions about the element Si and some of its compounds.

(a) The mass spectrum of a pure sample of Si is shown below.



(i) How many protons and how many neutrons are in the nucleus of an atom of the most abundant isotope of Si?

(ii) Write the ground-state electron configuration of Si.

Two compounds that contain Si are SiO_2 and SiH_4 .

(b) At 161 K, SiH_4 boils but SiO_2 remains as a solid. Using principles of interparticle forces, explain the difference in boiling points.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

At high temperatures, SiH_4 decomposes to form solid silicon and hydrogen gas.

(c) Write a balanced equation for the reaction.

A table of absolute entropies of some substances is given below.

Substance	S° (J/(mol · K))
$\text{H}_2(g)$	131
$\text{Si}(s)$	18
$\text{SiH}_4(g)$	205

(d) Explain why the absolute molar entropy of $\text{Si}(s)$ is less than that of $\text{H}_2(g)$.


(e) Calculate the value, in J/(mol · K), of ΔS° for the reaction.

(f) The reaction is thermodynamically favorable at all temperatures. Explain why the reaction occurs only at high temperatures.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

(g) A partial photoelectron spectrum of pure Si is shown below. On the spectrum, draw the missing peak that corresponds to the electrons in the $3p$ sublevel.



(h) Using principles of atomic structure, explain why the first ionization energy of Ge is lower than that of Si.

(i) A single photon with a wavelength of $4.00 \times 10^{-7} \text{ m}$ is absorbed by the Si sample. Calculate the energy of the photon in joules.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.