


2. Answer the following questions about ascorbic acid (vitamin C).


A. A student combusts a sample of ascorbic acid, $C_xH_yO_z$, to determine its chemical composition. The only products of the reaction are 0.2400 mol of CO_2 and 2.883 g of H_2O .

- Calculate the number of moles of H_2O produced.
- The mole ratio of carbon (C) to oxygen (O) is 1:1 in ascorbic acid. Based on this information and your answer to part A (i), determine the empirical formula of ascorbic acid.

B. Ascorbic acid, $HAsc(aq)$, acts as a weak acid, as shown in the equation.



The following titration curve was produced when a 10.0 mL sample of $HAsc(aq)$ was titrated using 0.0550 M $NaOH(aq)$.



- Calculate the molar concentration of the ascorbic acid solution.
- From the titration curve, determine the approximate pK_a of ascorbic acid.
- What is the value of the ratio $\frac{[Asc^-]}{[HAsc]}$ when the pH of the solution is 4.7?


C. Dehydroascorbic acid (DHAsc) can be produced by reacting ascorbic acid with the triiodide ion, I_3^- , as represented by the following equation.



The student runs three trials of the reaction with different initial concentrations of $HAsc$ and I_3^- , producing the following data.

Trial	$[HAsc] (M)$	$[I_3^-] (M)$	Initial Rate of DHAsc Formation (M/s)
1	0.450	1.200	2.457×10^{-4}
2	0.450	0.600	1.229×10^{-4}
3	0.900	1.200	4.914×10^{-4}


- The rate law for the reaction is $rate = k[HAsc][I_3^-]$. Explain how the data in the table support the conclusion that the reaction is first order with respect to $[HAsc]$.
 - Calculate the value of the rate constant, k , for the reaction. Include units with your answer.
- D. The triiodide ion, I_3^- , is significantly more soluble in water than elemental iodine, I_2 , is. Identify an intermolecular force between I_3^- and water that is **not** present between I_2 and water, which could explain the difference in solubility. Lewis diagrams for I_2 and I_3^- are provided.



Begin your response to **QUESTION 1** on this page.**CHEMISTRY****SECTION II****Time—1 hour and 45 minutes****7 Questions**


Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

For each question, show your work for each part in the space provided after that part. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.



1. A student is studying the reaction between lactic acid, $\text{C}_3\text{H}_6\text{O}_3$, and sodium hydroxide, NaOH , as represented in the balanced equation above.


(a) The structural formula of lactic acid is shown in the following diagram. Circle the hydrogen atom that most readily participates in the chemical reaction with sodium hydroxide.




- (b) The student begins the experiment by dissolving 10.22 g of sodium hydroxide (molar mass 40.00 g/mol) in enough water to produce 500. mL of solution. Calculate the molarity of the sodium hydroxide solution.

GO ON TO THE NEXT PAGE.Continue your response to **QUESTION 1** on this page.

The student uses the sodium hydroxide solution from part (b), a buret, a pH meter, and a 100 mL Erlenmeyer flask to titrate a 25.0 mL sample of lactic acid solution. The student's data are shown in the following graph.



- (c) Use the information in the graph to determine the approximate $\text{p}K_a$ of lactic acid. _____



- (d) The preceding diagram represents the relative amounts of major species in a sample of the solution in the flask at one point during the titration. (Note that water molecules are omitted.)


- (i) Draw an X on the preceding titration curve at a point in the titration where the reaction mixture would be represented by this diagram.
(ii) Justify your answer.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

Experiment	Mass of NaOH(<i>s</i>) (grams)	Volume of Solution (mL)	Titration Curve
1	10.22	500.	Already shown on graph
2	20.44	500.	?

- (iii) The student repeats the experiment but uses a solution of NaOH(*aq*) with twice the concentration, as shown in the preceding table. On the following graph, draw the titration curve that would be expected for experiment 2.



- (e) In a third experiment, the student investigates the enthalpy of the reaction between lactic acid and sodium hydroxide. The student combines 100.0 mL of a 0.500 *M* lactic acid solution at 20.0°C with 100.0 mL of a 0.500 *M* NaOH solution at 20.0°C in a calorimeter. The final temperature of the resulting combined solution is 23.2°C. Assume that the density of each solution before combining is 1.00 g/mL and that the specific heat capacity of the combined solution is 4.2 J / (g · °C).

- (i) Calculate the quantity of heat produced in the reaction, in J.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

- (ii) Calculate the molar enthalpy of reaction, in kJ / mol_{rxn}. Include the sign in your answer.

- (iii) The student claims that if heat is lost from the calorimeter to the surrounding air during the reaction, then the experimental value of the molar enthalpy of reaction will be smaller in magnitude than the actual value. Do you agree or disagree with the student's claim? Justify your answer.

GO ON TO THE NEXT PAGE.

2. A chemical reaction between maleic acid ($\text{H}_2\text{C}_4\text{H}_2\text{O}_4$) and sodium bicarbonate (NaHCO_3) occurs in the presence of water to produce carbon dioxide and sodium maleate ($\text{Na}_2\text{C}_4\text{H}_2\text{O}_4$), as represented by the following equation.



- (a) A student combines equal masses of $\text{H}_2\text{C}_4\text{H}_2\text{O}_4(s)$ chunks and $\text{NaHCO}_3(s)$ chunks with sufficient water at 20.0°C . The student determines that 0.0114 mol of $\text{CO}_2(g)$ is produced after the reaction goes to completion.

(i) Calculate the number of grams of $\text{CO}_2(g)$ produced.

(ii) The $\text{CO}_2(g)$ produced from the reaction at 20.0°C was collected and found to have a pressure of 1.25 atm . Calculate the volume of $\text{CO}_2(g)$, in liters.

- (b) The student performs a second experiment that is identical to the first except that the student grinds the chunks of $\text{H}_2\text{C}_4\text{H}_2\text{O}_4(s)$ and $\text{NaHCO}_3(s)$ into powder before combining the powder with water.

(i) What happens to the surface area of the reactants when the student grinds the chunks into powder?

(ii) The rate-determining step for the overall reaction is the dissolving of the solids. Would the time required for the dissolving of the solids in the second experiment be longer than, shorter than, or the same as the time required in the first experiment? Justify your answer based on the collisions between particles.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.

- (iii) When the reaction is complete, will the volume of $\text{CO}_2(g)$ at the end of the second experiment be greater than, less than, or equal to the volume at the end of the first experiment? Justify your answer.

The student conducts additional trials of the experiment and produces the following data table.

Trial	Mass of $\text{H}_2\text{C}_4\text{H}_2\text{O}_4$ (grams)	Mass of NaHCO_3 (grams)	Moles of CO_2 Produced (mol)
3	1.543	1.251	0.01489
4	1.543	1.686	0.02007

- (c) Based on the student's data, identify the limiting reactant in trial 3. Justify your answer.

- (d) The reaction has a value of ΔS° greater than zero. Using particle-level reasoning, explain why the entropy increases as the reaction progresses.



GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.

The student notices that the temperature of the reaction mixture decreases as the reaction takes place and correctly determines that the reaction is endothermic.

- (e) The student claims that the reaction is thermodynamically favorable at all temperatures because $\Delta S_{rxn}^{\circ} > 0$ and the reaction is endothermic. Do you agree or disagree with the student's claim? Justify your answer.

Next, the student investigates the acid-base behavior of maleic acid. The student notes that maleic acid is a diprotic acid. The two acid dissociation processes that occur are represented by the following equations.



- (f) Calculate the $\text{p}K_{a2}$ value for the $\text{HC}_4\text{H}_2\text{O}_4^-$ ion.

- (g) A buffer solution with a pH of 7.00 is prepared using $\text{C}_4\text{H}_2\text{O}_4^{2-}$ and $\text{HC}_4\text{H}_2\text{O}_4^-$. Calculate the ratio

$$\frac{[\text{C}_4\text{H}_2\text{O}_4^{2-}]}{[\text{HC}_4\text{H}_2\text{O}_4^-]} \text{ in this solution.}$$

GO ON TO THE NEXT PAGE.

7. A student conducts a chromatography experiment and needs to prepare 100.0 mL of 0.340 *M* NaCl(*aq*) to use as the solvent.

- (a) Calculate the mass of solid NaCl (molar mass 58.44 g/mol) needed to prepare the 100.0 mL of 0.340 *M* NaCl(*aq*).

- (b) In the following table, briefly list the additional steps necessary to prepare the 100.0 mL of 0.340 *M* NaCl(*aq*) solution using only materials selected from the choices given. Assume that all appropriate safety measures are already in place. Not all materials in the list may be needed.


- | | | |
|--------------|-----------------------------|------------------------------|
| • Solid NaCl | • Distilled water | • Weighing paper and scoop |
| • Balance | • 100.0 mL volumetric flask | • 50.0 mL graduated cylinder |
| • Pipet | • 150 mL beakers | • Chromatography paper |

Step	Step Description and Materials Used
1.	Use the weighing paper and scoop to measure the correct mass of solid NaCl on the balance.
2.	
3.	Swirl the mixture to dissolve the solid NaCl.
4.	
5.	Stopper and invert the mixture several times to ensure that the mixture is homogeneous.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 7** on this page.

The student uses the $\text{NaCl}(aq)$ solvent to separate a mixture of compounds X and Y in a chromatography experiment. After 30 minutes, the student removes the chromatography paper from the chamber. The results of the experiment are shown.



- (c) A second student conducts the same chromatography experiment but removes the chromatography paper from the chamber after 15 minutes instead of 30 minutes. Predict the effect, if any, this would have on the separation distance between compounds X and Y in the new experiment. Explain your reasoning.

STOP

END OF EXAM

CHEMISTRY

SECTION II

Time—1 hour and 45 minutes

7 Questions

Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

For each question, show your work for each part in the space provided after that part. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.


Answer the following questions related to manganese compounds.

(a) Manganese has several common oxidation states.

(i) Write the complete electron configuration for an Mn atom in the ground state.

(ii) When manganese forms cations, electrons are lost from which subshell first? Identify both the number and letter associated with the subshell.

A student performs an experiment to produce a manganese salt of unknown composition, $\text{Mn}_x\text{Cl}_y(aq)$, and determine its empirical formula. The student places a sample of $\text{Mn}(s)$ in a beaker containing excess $\text{HCl}(aq)$, as represented by the following equation.



GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

The student heats the resulting mixture until only $\text{Mn}_x\text{Cl}_y(s)$ remains in the beaker. The data are given in the following table.


Mass of empty beaker	60.169 g
Mass of beaker and $\text{Mn}(s)$	61.262 g
Mass of beaker and Mn_xCl_y after heating to constant mass	62.673 g

- (b) Calculate the mass of Cl in the sample of $\text{Mn}_x\text{Cl}_y(s)$ remaining in the beaker.
- (c) Calculate the number of moles of Cl in the sample of $\text{Mn}_x\text{Cl}_y(s)$ remaining in the beaker.
- (d) The student determines that 0.0199 mol of Mn was used in the experiment. Use the data to determine the empirical formula of the $\text{Mn}_x\text{Cl}_y(s)$.
- (e) The student repeats the experiment using the same amounts of Mn and HCl and notices that some of the Mn_xCl_y splatters out of the beaker as it is heated to dryness. Will the number of moles of Cl calculated for this trial be greater than, less than, or equal to the number calculated in part (c)? Justify your answer.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

- (f) Another compound of manganese, MnO_2 , is used in alkaline batteries, represented by the following diagram. Some half-reactions are given in the table.



Reduction Half-Reaction	E° (V)
$\text{Zn}^{2+}(aq) + 2 e^- \rightarrow \text{Zn}(s)$	-0.76
$\text{ZnO}(s) + \text{H}_2\text{O}(l) + 2 e^- \rightarrow \text{Zn}(s) + 2 \text{OH}^-(aq)$	-1.28
$2 \text{MnO}_2(s) + \text{H}_2\text{O}(l) + 2 e^- \rightarrow \text{Mn}_2\text{O}_3(s) + 2 \text{OH}^-(aq)$	0.15

- (i) Based on the half-reactions given in the table, write the balanced net ionic equation for the reaction that has the greatest thermodynamic favorability.
- (ii) Calculate the value of E_{cell}° for the overall reaction.
- (iii) Calculate the value of ΔG° in kJ/mol_{rxn}.
- (iv) A student claims that the total mass of an alkaline battery decreases as the battery operates because the anode loses mass. Do you agree with the student's claim? Justify your answer.

GO ON TO THE NEXT PAGE.

Answer the following questions about the compounds NH_2Cl and NCl_3 . The Lewis electron-dot diagrams of the two compounds are shown.



(a) Calculate the number of moles of NH_2Cl (molar mass 51.48 g/mol) present in 1.0 L of a solution in which the concentration of NH_2Cl is 0.0016 g/L.

(b) NH_2Cl is highly soluble in water, whereas NCl_3 is nearly insoluble. Explain this observation in terms of the types and relative strengths of the intermolecular forces between each of the solutes and water.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 4** on this page.


(c) The value of $\Delta H_{\text{vaporization}}^\circ$ for $\text{NCl}_3(l)$ is 32.9 kJ/mol. Calculate the amount of energy required to vaporize a 15.0 g sample of NCl_3 (molar mass 120.36 g/mol).

GO ON TO THE NEXT PAGE.

CHEMISTRY**Section II****Time—1 hour and 45 minutes****7 Questions****YOU MAY USE YOUR CALCULATOR FOR THIS SECTION.**



Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

Write your response in the space provided following each question. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.




1. The compound urea, H_2NCONH_2 , is widely used in chemical fertilizers. The complete Lewis electron-dot diagram for the urea molecule is shown above.

- (a) Identify the hybridization of the valence orbitals of the carbon atom in the urea molecule.
- (b) Urea has a high solubility in water, due in part to its ability to form hydrogen bonds. A urea molecule and four water molecules are represented in the box below. Draw ONE dashed line (----) to indicate a possible location of a hydrogen bond between a water molecule and the urea molecule.



The dissolution of urea is represented by the equation above. A student determines that 5.39 grams of H_2NCONH_2 (molar mass 60.06 g/mol) can dissolve in water to make 5.00 mL of a saturated solution at 20.°C.

- (c) Calculate the concentration of urea, in mol/L, in the saturated solution at 20.°C.
- (d) The student also determines that the concentration of urea in a saturated solution at 25°C is 19.8 *M*. Based on this information, is the dissolution of urea endothermic or exothermic? Justify your answer in terms of Le Chatelier's principle.



- (e) The equipment shown above is provided so that the student can determine the value of the molar heat of solution for urea. Knowing that the specific heat of the solution is 4.18 J/(g·°C), list the specific measurements that are required to be made during the experiment.


	S° (J/(mol·K))
$\text{H}_2\text{NCONH}_2(s)$	104.6
$\text{H}_2\text{NCONH}_2(aq)$	?

- (f) The entropy change for the dissolution of urea, $\Delta S^\circ_{\text{soln}}$, is 70.1 J/(mol·K) at 25°C. Using the information in the table above, calculate the absolute molar entropy, S° , of aqueous urea.
- (g) Using particle-level reasoning, explain why $\Delta S^\circ_{\text{soln}}$ is positive for the dissolution of urea in water.
- (h) The student claims that ΔS° for the process contributes to the thermodynamic favorability of the dissolution of urea at 25°C. Use the thermodynamic information above to support the student's claim.

CHEMISTRY**Section II****Time—1 hour and 45 minutes****7 Questions****YOU MAY USE YOUR CALCULATOR FOR THIS SECTION.**

Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

Write your response in the space provided following each question. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.



1. A student performs an experiment to determine the value of the enthalpy change, $\Delta H_{\text{rxn}}^\circ$, for the oxidation-reduction reaction represented by the balanced equation above.


- (a) Determine the oxidation number of Cl in NaOCl.
 (b) Calculate the number of grams of $\text{Na}_2\text{S}_2\text{O}_3$ needed to prepare 100.00 mL of 0.500 M $\text{Na}_2\text{S}_2\text{O}_3(aq)$.

In the experiment, the student uses the solutions shown in the table below.

Solution	Concentration (M)	Volume (mL)
$\text{Na}_2\text{S}_2\text{O}_3(aq)$	0.500	5.00
$\text{NaOCl}(aq)$	0.500	5.00
$\text{NaOH}(aq)$	0.500	5.00

- (c) Using the balanced equation for the oxidation-reduction reaction and the information in the table above, determine which reactant is the limiting reactant. Justify your answer.

The solutions, all originally at 20.0°C, are combined in an insulated calorimeter. The temperature of the reaction mixture is monitored, as shown in the graph below.




- (d) According to the graph, what is the temperature change of the reaction mixture?
 (e) The mass of the reaction mixture inside the calorimeter is 15.21 g.
 (i) Calculate the magnitude of the heat energy, in joules, that is released during the reaction. Assume that the specific heat of the reaction mixture is 3.94 J/(g·°C) and that the heat absorbed by the calorimeter is negligible.
 (ii) Using the balanced equation for the oxidation-reduction reaction and your answer to part (c), calculate the value of the enthalpy change of the reaction, $\Delta H_{\text{rxn}}^\circ$, in kJ/mol_{rxn}. Include the appropriate algebraic sign with your answer.


The student repeats the experiment, but this time doubling the volume of each of the reactants, as shown in the table below.

Solution	Concentration (M)	Volume (mL)
$\text{Na}_2\text{S}_2\text{O}_3(aq)$	0.500	10.0
$\text{NaOCl}(aq)$	0.500	10.0
$\text{NaOH}(aq)$	0.500	10.0

- (f) The magnitude of the enthalpy change, $\Delta H_{\text{rxn}}^\circ$, in kJ/mol_{rxn}, calculated from the results of the second experiment is the same as the result calculated in part (e)(ii). Explain this result.
 (g) Write the balanced net ionic equation for the given reaction.



3. To determine the molar mass of an unknown metal, M, a student reacts iodine with an excess of the metal to form the water-soluble compound MI_2 , as represented by the equation above. The reaction proceeds until all of the I_2 is consumed. The $\text{MI}_2(aq)$ solution is quantitatively collected and heated to remove the water, and the product is dried and weighed to constant mass. The experimental steps are represented below, followed by a data table.



Data for Unknown Metal Lab	
Mass of beaker	125.457 g
Mass of beaker + metal M	126.549 g
Mass of beaker + metal M + I_2	127.570 g
Mass of MI_2 , first weighing	1.284 g
Mass of MI_2 , second weighing	1.284 g

- (a) Given that the metal M is in excess, calculate the number of moles of I_2 that reacted.
- (b) Calculate the molar mass of the unknown metal M.

The student hypothesizes that the compound formed in the synthesis reaction is ionic.

- (c) Propose an experimental test the student could perform that could be used to support the hypothesis. Explain how the results of the test would support the hypothesis if the substance was ionic.

The student hypothesizes that Br_2 will react with metal M more vigorously than I_2 did because Br_2 is a liquid at room temperature.

- (d) Explain why I_2 is a solid at room temperature whereas Br_2 is a liquid. Your explanation should clearly reference the types and relative strengths of the intermolecular forces present in each substance.

While cleaning up after the experiment, the student wishes to dispose of the unused solid I_2 in a responsible manner. The student decides to convert the solid I_2 to $\text{I}^-(aq)$ anion. The student has access to three solutions, $\text{H}_2\text{O}_2(aq)$, $\text{Na}_2\text{S}_2\text{O}_3(aq)$, and $\text{Na}_2\text{S}_4\text{O}_6(aq)$, and the standard reduction table shown below.

Half reaction	E° (V)
$\text{S}_4\text{O}_6^{2-}(aq) + 2 e^- \rightarrow 2 \text{S}_2\text{O}_3^{2-}(aq)$	0.08
$\text{I}_2(s) + 2 e^- \rightarrow 2 \text{I}^-(aq)$	0.54
$\text{O}_2(g) + 2 \text{H}^+(aq) + 2 e^- \rightarrow \text{H}_2\text{O}_2(aq)$	0.68

- (e) Which solution should the student add to $\text{I}_2(s)$ to reduce it to $\text{I}^-(aq)$? Circle your answer below. Justify your answer, including a calculation of E° for the overall reaction.



- (f) Write the balanced net-ionic equation for the reaction between I_2 and the solution you selected in part (e).