

2. A chemical reaction between maleic acid ($\text{H}_2\text{C}_4\text{H}_2\text{O}_4$) and sodium bicarbonate (NaHCO_3) occurs in the presence of water to produce carbon dioxide and sodium maleate ($\text{Na}_2\text{C}_4\text{H}_2\text{O}_4$), as represented by the following equation.



- (a) A student combines equal masses of $\text{H}_2\text{C}_4\text{H}_2\text{O}_4(s)$ chunks and $\text{NaHCO}_3(s)$ chunks with sufficient water at 20.0°C . The student determines that 0.0114 mol of $\text{CO}_2(g)$ is produced after the reaction goes to completion.

(i) Calculate the number of grams of $\text{CO}_2(g)$ produced.

- (ii) The $\text{CO}_2(g)$ produced from the reaction at 20.0°C was collected and found to have a pressure of 1.25 atm . Calculate the volume of $\text{CO}_2(g)$, in liters.

- (b) The student performs a second experiment that is identical to the first except that the student grinds the chunks of $\text{H}_2\text{C}_4\text{H}_2\text{O}_4(s)$ and $\text{NaHCO}_3(s)$ into powder before combining the powder with water.

(i) What happens to the surface area of the reactants when the student grinds the chunks into powder?

- (ii) The rate-determining step for the overall reaction is the dissolving of the solids. Would the time required for the dissolving of the solids in the second experiment be longer than, shorter than, or the same as the time required in the first experiment? Justify your answer based on the collisions between particles.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.

- (iii) When the reaction is complete, will the volume of $\text{CO}_2(g)$ at the end of the second experiment be greater than, less than, or equal to the volume at the end of the first experiment? Justify your answer.

The student conducts additional trials of the experiment and produces the following data table.

Trial	Mass of $\text{H}_2\text{C}_4\text{H}_2\text{O}_4$ (grams)	Mass of NaHCO_3 (grams)	Moles of CO_2 Produced (mol)
3	1.543	1.251	0.01489
4	1.543	1.686	0.02007

- (c) Based on the student's data, identify the limiting reactant in trial 3. Justify your answer.

- (d) The reaction has a value of ΔS° greater than zero. Using particle-level reasoning, explain why the entropy increases as the reaction progresses.



GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.

The student notices that the temperature of the reaction mixture decreases as the reaction takes place and correctly determines that the reaction is endothermic.

- (e) The student claims that the reaction is thermodynamically favorable at all temperatures because $\Delta S_{rxn}^{\circ} > 0$ and the reaction is endothermic. Do you agree or disagree with the student's claim? Justify your answer.

Next, the student investigates the acid-base behavior of maleic acid. The student notes that maleic acid is a diprotic acid. The two acid dissociation processes that occur are represented by the following equations.



- (f) Calculate the $\text{p}K_{a_2}$ value for the $\text{HC}_4\text{H}_2\text{O}_4^-$ ion.

- (g) A buffer solution with a pH of 7.00 is prepared using $\text{C}_4\text{H}_2\text{O}_4^{2-}$ and $\text{HC}_4\text{H}_2\text{O}_4^-$. Calculate the ratio

$$\frac{[\text{C}_4\text{H}_2\text{O}_4^{2-}]}{[\text{HC}_4\text{H}_2\text{O}_4^-]} \text{ in this solution.}$$

GO ON TO THE NEXT PAGE.

Answer the following questions about an experiment in which $\text{CaCO}_3(s)$ is combined with $\text{HCl}(aq)$, represented by the following balanced equation.



(a) Write the balanced net ionic equation for the reaction.

A student performs an investigation to study factors that affect the rate of the reaction. In each trial the student combines 50.0 mL of $\text{HCl}(aq)$ at 21.2°C with 1.00 g of $\text{CaCO}_3(s)$ and measures the time required for the reaction to go to completion. The data are given in the following table.

Trial	Concentration of $\text{HCl}(aq)$ (M)	Particle Size of $\text{CaCO}_3(s)$	Time of Reaction (s)
1	1.00	Fine powder	67
2	1.00	Small chunks	112
3	1.00	Large chunk	342
4	3.00	Fine powder	22
5	3.00	Small chunks	227
6	3.00	Large chunk	114

(b) The student correctly identifies that trial 5 is inconsistent with the other trials. Explain why the student's claim is correct using the data in the table.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 3** on this page.

- (c) Based on the reaction conditions and the collisions that occur between particles, explain the reason for the difference in the reaction times for trial 2 and trial 3.
- (d) The student claims that the reaction is zero order with respect to $\text{HCl}(aq)$. Do you agree or disagree with the student's claim? Justify your answer using the student's data.
- (e) The $\text{HCl}(aq)$ was present in excess in all trials of the experiment. Determine the molarity of the $\text{HCl}(aq)$ in the beaker after the reaction is complete in trial 2. Assume that the volume of the mixture remains constant at 50.0 mL throughout the trial. (The molar mass of CaCO_3 is 100.09 g/mol.)

GO ON TO THE NEXT PAGE.

Continue your response to QUESTION 6 on this page.



In order to measure the enthalpy of the reaction shown, the student repeats trial 1 by mixing 50.0 mL of $\text{HCl}(aq)$ with 1.00 g of $\text{CaCO}_3(s)$ using a coffee cup calorimeter. The student records the temperature of the system every 20 seconds. The data are given in the following table.

Time (s)	Measured Temperature of Solution ($^{\circ}\text{C}$)
0	21.20
20	21.51
40	21.70
60	21.85
80	21.90
100	21.90

(f) Is the reaction endothermic or exothermic? Justify your answer using the information in the table.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 3** on this page.

(g) Based on the experimental data, the mass of the system is 51.0 g, and the specific heat of the reaction mixture is $4.0 \text{ J} / (\text{g} \cdot ^\circ\text{C})$.

(i) Calculate the magnitude of heat transfer, q , in joules.


(ii) Calculate the enthalpy of reaction in units of kJ/mol_{rxn} . Include the algebraic sign on your answer.

GO ON TO THE NEXT PAGE.


CHEMISTRY**Section II****7 Questions****Time—1 hour and 45 minutes****YOU MAY USE YOUR CALCULATOR FOR THIS SECTION.**

Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

Write your response in the space provided following each question. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.




1. Carbon tetrachloride, $\text{CCl}_4(g)$, can be synthesized according to the reaction represented above. A chemist runs the reaction at a constant temperature of 120°C in a rigid 25.0 L container.
 - (a) Chlorine gas, $\text{Cl}_2(g)$, is initially present in the container at a pressure of 0.40 atm.
 - (i) How many moles of $\text{Cl}_2(g)$ are in the container?
 - (ii) How many grams of carbon disulfide, $\text{CS}_2(g)$, are needed to react completely with the $\text{Cl}_2(g)$?
 - (b) At 30°C the reaction is thermodynamically favorable, but no reaction is observed to occur. However, at 120°C , the reaction occurs at an observable rate.
 - (i) Explain how the higher temperature affects the collisions between the reactant molecules so that the reaction occurs at an observable rate at 120°C .
 - (ii) The graph below shows a distribution for the collision energies of reactant molecules at 120°C . Draw a second curve on the graph that shows the distribution for the collision energies of reactant molecules at 30°C .




(c) S_2Cl_2 is a product of the reaction.

- (i) In the box below, complete the Lewis electron-dot diagram for the S_2Cl_2 molecule by drawing in all of the electron pairs.



- (ii) What is the approximate value of the Cl-S-S bond angle in the S_2Cl_2 molecule that you drew in part (c)(i) ? (If the two Cl-S-S bond angles are not equal, include both angles.)

(d) $\text{CCl}_4(g)$ can also be produced by reacting $\text{CHCl}_3(g)$ with $\text{Cl}_2(g)$ at 400°C , as represented by the equation below.




At the completion of the reaction a chemist successfully separates the $\text{CCl}_4(g)$ from the $\text{HCl}(g)$ by cooling the mixture to 70°C , at which temperature the $\text{CCl}_4(g)$ condenses while the $\text{HCl}(g)$ remains in the gaseous state.

- (i) Identify all types of intermolecular forces present in $\text{HCl}(l)$.
- (ii) What can be inferred about the relative strengths of the intermolecular forces in $\text{CCl}_4(l)$ and $\text{HCl}(l)$? Justify your answer in terms of the information above.



2. A student designs an experiment to study the reaction between NaHCO_3 and $\text{HC}_2\text{H}_3\text{O}_2$. The reaction is represented by the equation above. The student places 2.24 g of NaHCO_3 in a flask and adds 60.0 mL of 0.875 M $\text{HC}_2\text{H}_3\text{O}_2$. The student observes the formation of bubbles and that the flask gets cooler as the reaction proceeds.
- (a) Identify the reaction represented above as an acid-base reaction, precipitation reaction, or redox reaction. Justify your answer.
- (b) Based on the information above, identify the limiting reactant. Justify your answer with calculations.
- (c) The student observes that the bubbling is rapid at the beginning of the reaction and gradually slows as the reaction continues. Explain this change in the reaction rate in terms of the collisions between reactant particles.
- (d) In thermodynamic terms, a reaction can be driven by enthalpy, entropy, or both.
- (i) Considering that the flask gets cooler as the reaction proceeds, what drives the chemical reaction between $\text{NaHCO}_3(s)$ and $\text{HC}_2\text{H}_3\text{O}_2(aq)$? Answer by drawing a circle around one of the choices below.
- Enthalpy only Entropy only Both enthalpy and entropy
- (ii) Justify your selection in part (d)(i) in terms of ΔG° .
- (e) The HCO_3^- ion has three carbon-to-oxygen bonds. Two of the carbon-to-oxygen bonds have the same length and the third carbon-to-oxygen bond is longer than the other two. The hydrogen atom is bonded to one of the oxygen atoms. In the box below, draw a Lewis electron-dot diagram (or diagrams) for the HCO_3^- ion that is (are) consistent with the given information.

- (f) A student prepares a solution containing equimolar amounts of $\text{HC}_2\text{H}_3\text{O}_2$ and $\text{NaC}_2\text{H}_3\text{O}_2$. The pH of the solution is measured to be 4.7. The student adds two drops of 3.0 M $\text{HNO}_3(aq)$ and stirs the sample, observing that the pH remains at 4.7. Write a balanced, net-ionic equation for the reaction between $\text{HNO}_3(aq)$ and the chemical species in the sample that is responsible for the pH remaining at 4.7.



7. The half-life ($t_{1/2}$) of the catalyzed isomerization of *cis*-2-butene gas to produce *trans*-2-butene gas, represented above, was measured under various conditions, as shown in the table below.

Trial Number	Initial $P_{\text{cis-2-butene}}$ (torr)	V (L)	T (K)	$t_{1/2}$ (s)
1	300.	2.00	350.	100.
2	600.	2.00	350.	100.
3	300.	4.00	350.	100.
4	300.	2.00	365	50.

- (a) The reaction is first order. Explain how the data in the table are consistent with a first-order reaction.
- (b) Calculate the rate constant, k , for the reaction at 350. K. Include appropriate units with your answer.
- (c) Is the initial rate of the reaction in trial 1 greater than, less than, or equal to the initial rate in trial 2? Justify your answer.
- (d) The half-life of the reaction in trial 4 is less than the half-life in trial 1. Explain why, in terms of activation energy.

STOP

END OF EXAM