


4. A scientist is investigating the properties of a mixture of CH_3OH and H_2CO . The scientist generates the following diagram to represent the mixture.



- A. Identify the hybridization of the valence orbitals of the C atom in the H_2CO molecule.
- B. In the diagram provided, draw a SINGLE dashed line (---) to represent a strong hydrogen-bonding attraction between one CH_3OH molecule and one H_2CO molecule in the mixture.
- C. The scientist plans to cool a gaseous mixture of CH_3OH and H_2CO to form a liquid mixture and finds data on the two compounds. The data are summarized in the table.

Substance	Melting Point (K)	Boiling Point (K)	Enthalpy of Vaporization (kJ/mol)
CH_3OH	176	338	37.6
H_2CO	181	254	24.2

- i. Propose a temperature to which the mixture should be cooled such that CH_3OH and H_2CO will both be liquids.
- ii. The scientist analyzes the mixture after it is cooled and determines that 8.59 g of $\text{CH}_3\text{OH}(l)$ is present. Calculate the amount of thermal energy, in kJ, that was removed to condense the 8.59 g of CH_3OH (molar mass 32.04 g/mol) at its boiling point.

. Answer the following questions related to $\text{HBr}(l)$ and $\text{HF}(l)$.

- (a) In the following table, list all of the types of intermolecular forces present in pure samples of $\text{HBr}(l)$ and $\text{HF}(l)$.

Liquid	$\text{HBr}(l)$	$\text{HF}(l)$
Intermolecular forces present		

- (b) The enthalpy of vaporization, $\Delta H_{\text{vap}}^{\circ}$, for each liquid is provided in the following table.

Liquid	$\text{HBr}(l)$	$\text{HF}(l)$
$\Delta H_{\text{vap}}^{\circ}$	17.3 kJ/mol	25.2 kJ/mol

- (i) Based on the types and relative strengths of intermolecular forces, explain why $\Delta H_{\text{vap}}^{\circ}$ of $\text{HF}(l)$ is greater than that of $\text{HBr}(l)$.

- (ii) Calculate the amount of thermal energy, in kJ, required to vaporize 6.85 g of $\text{HF}(l)$.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 6** on this page.

- (c) Based on the arrangement of electrons in the Br and F atoms, explain why the bond length in an HBr molecule is greater than that in an HF molecule.

GO ON TO THE NEXT PAGE.

CHEMISTRY**SECTION II****Time—1 hour and 45 minutes****7 Questions****YOU MAY USE YOUR CALCULATOR FOR THIS SECTION.**

Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

For each question, show your work for each part in the space provided after that part. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.

4. A student reacts 0.300 g of methyl salicylate ($\text{C}_8\text{H}_8\text{O}_3$) with a stoichiometric amount of a strong base. This product is then acidified to produce salicylic acid crystals ($\text{HC}_7\text{H}_5\text{O}_3$).

- (a) For every 1 mole of $\text{C}_8\text{H}_8\text{O}_3$ (molar mass 152.15 g/mol) reactant used, 1 mole of salicylic acid crystals ($\text{HC}_7\text{H}_5\text{O}_3$, molar mass 138.12 g/mol) is produced. Calculate the maximum mass, in grams, of $\text{HC}_7\text{H}_5\text{O}_3$ that could be produced in this reaction.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

As part of the experimental procedure to purify the $\text{HC}_7\text{H}_5\text{O}_3$ crystals after the reaction is complete, the crystals are filtered from the reaction mixture, rinsed with distilled water, and dried. Some physical properties of $\text{HC}_7\text{H}_5\text{O}_3$ are given in the following table.

Properties of Salicylic Acid ($\text{HC}_7\text{H}_5\text{O}_3$)	
Melting point	159°C
Solubility in H_2O at 25°C	2.2 g/L
Specific heat capacity	1.17 J/(g·°C)
Heat of fusion	27.1 kJ/mol


(b) The student's experiment results in an 87% yield of dry $\text{HC}_7\text{H}_5\text{O}_3$. The student suggests that some of the $\text{HC}_7\text{H}_5\text{O}_3$ crystals dissolved in the distilled water during the rinsing step. Is the student's claim consistent with the calculated percent yield value? Justify your answer.

(c) Given the physical properties in the table, calculate the quantity of heat that must be absorbed to increase the temperature of a 0.105 g sample of dry $\text{HC}_7\text{H}_5\text{O}_3$ (molar mass 138.12 g/mol) crystals from 25°C to the melting point of 159°C and melt the crystals completely.


GO ON TO THE NEXT PAGE.

Continue your response to QUESTION 7 on this page.

The structures and melting points for methyl salicylate and salicylic acid are shown.




Methyl Salicylate
Melting Point: -9°C



Salicylic Acid
Melting Point: 159°C

- (d) The same three types of intermolecular forces (London dispersion forces, dipole-dipole interactions, and hydrogen bonding) exist among molecules of each substance. Explain why the melting point of salicylic acid is higher than that of methyl salicylate.


The student titrates 20.0 mL of 0.0100 M $\text{HC}_7\text{H}_5\text{O}_3(aq)$ with 0.0200 M NaOH, using a probe to monitor the pH of the solution. The data are plotted producing the following titration curve.





GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

- (e) Using the information in the graph, estimate the pK_a of $HC_7H_5O_3$. _____
- (f) When the pH of the titration mixture is 4.00, is there a higher concentration of the weak acid, $HC_7H_5O_3$, or its conjugate base, $C_7H_5O_3^-$, in the flask? Justify your answer.
- (g) The student researches benzoic acid ($HC_7H_5O_2$) and finds that it has similar properties to salicylic acid ($HC_7H_5O_3$). The K_a for benzoic acid is 6.3×10^{-5} . Calculate the value of pK_a for benzoic acid.
- (h) The student performs a second titration, this time titrating 20.0 mL of a 0.0100 M benzoic acid solution with 0.0200 M NaOH. Sketch the curve that would result from this titration of benzoic acid on the following graph, which already shows the original curve from the titration of 20.0 mL of 0.0100 M salicylic acid. The initial pH of the benzoic acid solution is 3.11.

**GO ON TO THE NEXT PAGE.**

Answer the following questions about the compounds NH_2Cl and NCl_3 . The Lewis electron-dot diagrams of the two compounds are shown.



- (a) Calculate the number of moles of NH_2Cl (molar mass 51.48 g/mol) present in 1.0 L of a solution in which the concentration of NH_2Cl is 0.0016 g/L.

- (b) NH_2Cl is highly soluble in water, whereas NCl_3 is nearly insoluble. Explain this observation in terms of the types and relative strengths of the intermolecular forces between each of the solutes and water.

GO ON TO THE NEXT PAGE.

Name:

AP Chemistry: 2022


Continue your response to **QUESTION 4** on this page.

- (c) The value of $\Delta H_{\text{vaporization}}^{\circ}$ for $\text{NCl}_3(l)$ is 32.9 kJ/mol. Calculate the amount of energy required to vaporize a 15.0 g sample of NCl_3 (molar mass 120.36 g/mol).

GO ON TO THE NEXT PAGE.

7. Aluminum metal can be recycled from scrap metal by melting the metal to evaporate impurities.

- (a) Calculate the amount of heat needed to purify 1.00 mole of Al originally at 298 K by melting it. The melting point of Al is 933 K. The molar heat capacity of Al is 24 J/(mol·K), and the heat of fusion of Al is 10.7 kJ/mol.
- (b) The equation for the overall process of extracting Al from Al_2O_3 is shown below. Which requires less energy, recycling existing Al or extracting Al from Al_2O_3 ? Justify your answer with a calculation.



STOP

END OF EXAM