


Question 7

The following information applies to parts A and B.

7. Sodium metal reacts with oxygen gas according to Equation 1.



Standard enthalpies of formation are provided in Table 1.

Table 1

Substance	ΔH_f° (kJ/mol)
Na(s)	0
O ₂ (g)	0
Na ₂ O(s)	?

Part A

Based on the information given, **calculate** the value of ΔH_f° for Na₂O(s). Show the work that leads to your answer.

Part B

An 18.4 g sample of Na(s) reacts with 12.8 g of O₂(g) according to Equation 1. **Calculate** the total amount of heat, in kilojoules, released during this process. Show the work that leads to your answer.

Part C

Lattice enthalpy can be defined as the energy required to separate an ionic crystal into gaseous ions. Rb₂O(s) and Na₂O(s) have similar crystal structures, and their lattice enthalpies are given in Table 2.



Table 2

Compound	Lattice Enthalpy (kJ/mol)
Na ₂ O(s)	2481
Rb ₂ O(s)	2163


Using Coulomb's law, **explain** why the lattice enthalpy of Rb₂O(s) is smaller than that of Na₂O(s).

STOP
END OF EXAM

3. White phosphorus is composed of P_4 molecules with a tetrahedral structure, as shown in the diagram on the left. Each P atom is bonded to the other three P atoms by single bonds, as shown in the incomplete Lewis diagram on the right.




- A. In the box in part A, complete the Lewis diagram for P_4 by drawing the nonbonding electrons.
- B. The reaction of white phosphorus with oxygen to form $P_4O_{10}(s)$ is thermodynamically favorable at 298 K. The reaction is represented by equation 1.




- The entropy change of the reaction, ΔS° , is negative. Using particle-level reasoning, explain why the entropy decreases as the reaction progresses.
- The enthalpy change of the reaction, ΔH° , is also negative. A student claims that the favorability of the reaction is driven by enthalpy and **not** by entropy. Is the student's claim correct? Justify your answer by using the relationship between ΔG° , ΔH° , and ΔS° .

$P_4O_{10}(s)$ reacts exothermically with water to form phosphoric acid, as represented by equation 2.



A chemist uses a calorimetry experiment to determine the enthalpy change for the reaction, as represented by the following diagram.





- C. The chemist carries out the calorimetry experiment and records the following information.

Mass of P_4O_{10}	0.100 g
Mass of H_2O	100.0 g
Initial temperature	22.00° C
Final temperature	22.38° C
Molar mass of P_4O_{10}	283.9 g/mol
Specific heat of H_2O	4.18 J/(g·°C)


- Calculate the amount of heat, q , released during the experiment, in kJ. Assume that the specific heat of the solution is the same as that of water.
- Calculate the value of ΔH°_{rxn} for equation 2 in kJ/mol_{rxn}. Include the sign in your answer.

D. The chemist weighed out 0.100 g P_4O_{10} of and 100.0 g of H_2O to perform a second trial. In the second trial, some of the solid P_4O_{10} stuck to the weighing paper and was not transferred to the calorimeter. Given that P_4O_{10} is the limiting reactant, would ΔT for the second trial be greater than, less than, or equal to the value in the first trial? Justify your answer.


$\text{P}_4(\text{s})$ also reacts readily with $\text{Cl}_2(\text{g})$ to produce phosphorus trichloride, $\text{PCl}_3(\text{g})$, which in turn reacts with $\text{Cl}_2(\text{g})$ in an equilibrium process to produce $\text{PCl}_5(\text{g})$. The reactions are represented by equations 3 and 4.




E. Calculate the standard enthalpy of formation of $\text{PCl}_5(\text{g})$ represented by equation 5.



The following particle-level diagram represents the contents of the vessel in an equilibrium mixture at 546 K involving equation 4.




F. Equation 4 for the reaction that occurs is shown.



- If each particle in the diagram represents a partial pressure of 1.00 atm, what is the value of K_p for the equilibrium mixture at 546 K?
- Does the value of K_p increase, decrease, or remain the same when the temperature is increased to 596 K? Justify your answer based on ΔH_2° .

2. In the gas phase, AlCl_3 is a molecular substance. A reaction of gaseous AlCl_3 at high temperature is represented by the following balanced equation.





(a) How many grams of $\text{Cl}(\text{g})$ can be formed from 1.25 mol of $\text{AlCl}_3(\text{g})$?

Additional reactions that involve Al or Cl are shown in the following table.



Reaction Number	Equation	$\Delta H_{\text{rxn}}^\circ$ (kJ/mol _{rxn})
2	$\text{Al}(\text{s}) + \frac{3}{2} \text{Cl}_2(\text{g}) \rightarrow \text{AlCl}_3(\text{g})$	-583
3	$\text{Al}(\text{s}) \rightarrow \text{Al}(\text{g})$	+326
4	$\text{Cl}_2(\text{g}) \rightarrow 2 \text{Cl}(\text{g})$	+243

(b) Calculate the value of ΔH_1° , in kJ/mol_{rxn}, for reaction 1 above using reactions 2, 3, and 4.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.(c) A potential energy diagram for Cl_2 is shown in the following graph.(i) Based on the graph, what is the bond length, in picometers, for Cl_2 ? _____(ii) A student finds that the average $\text{Al} - \text{Cl}$ bond length is 220 picometers and the average bond energy is 425 kJ/mol. Draw the potential energy curve for the average $\text{Al} - \text{Cl}$ bond on the preceding graph.(d) Three proposed Lewis diagrams for the $\text{AlCl}_3(g)$ molecule are shown.(i) The $\text{AlCl}_3(g)$ molecule has a trigonal planar geometry. Which diagram (1, 2, or 3) can be eliminated based on geometry? Justify your choice based on VSEPR theory.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.(ii) Which of the three diagrams is the best representation for the bonding in AlCl_3 ? Justify your choice based on formal charges. AlCl_3 is known to dimerize reversibly in the gas phase. The dimerization equilibrium is represented by the following equation.(e) Write the expression for the equilibrium constant, K_p , for this reaction.A particle-level diagram of an equilibrium mixture of $\text{AlCl}_3(g)$ and $\text{Al}_2\text{Cl}_6(g)$ at 400°C in a 25 L closed container is shown.(f) Using the particle-level diagram, calculate the value of K_p for the reaction if the total pressure in the container is 22.1 atm.

GO ON TO THE NEXT PAGE.

2. Answer the following questions relating to the chemistry of the halogens.

- (a) The molecular formulas of diatomic bromine, chlorine, fluorine, and iodine are written below. Circle the formula of the molecule that has the longest bond length. Justify your choice in terms of atomic structure.



A chemistry teacher wants to prepare Br_2 . The teacher has access to the following three reagents: $\text{NaBr}(aq)$, $\text{Cl}_2(g)$, and $\text{I}_2(s)$.


Half-Reaction	E° at 25°C (V)
$\text{Br}_2 + 2 e^- \rightarrow 2 \text{Br}^-$	1.07
$\text{Cl}_2 + 2 e^- \rightarrow 2 \text{Cl}^-$	1.36
$\text{I}_2 + 2 e^- \rightarrow 2 \text{I}^-$	0.53

- (b) Using the data in the table above, write the balanced equation for the thermodynamically favorable reaction that will produce Br_2 when the teacher combines two of the reagents. Justify that the reaction is thermodynamically favorable by calculating the value of E° for the reaction.

Br_2 and Cl_2 can react to form the compound BrCl .

- (c) The boiling point of Br_2 is 332 K, whereas the boiling point of BrCl is 278 K. Explain this difference in boiling point in terms of all the intermolecular forces present between molecules of each substance.

The compound BrCl can decompose into Br_2 and Cl_2 , as represented by the balanced chemical equation below.



A 0.100 mole sample of pure $\text{BrCl}(g)$ is placed in a previously evacuated, rigid 2.00 L container at 298 K. Eventually the system reaches equilibrium according to the equation above.

- (d) Calculate the pressure in the container before equilibrium is established.
- (e) Write the expression for the equilibrium constant, K_{eq} , for the decomposition of BrCl .

After the system has reached equilibrium, 42 percent of the original BrCl sample has decomposed.


- (f) Determine the value of K_{eq} for the decomposition reaction of BrCl at 298 K.
- (g) Calculate the bond energy of the Br-Cl bond, in kJ/mol, using ΔH° for the reaction ($1.6 \text{ kJ/mol}_{\text{rxn}}$) and the information in the following table.

Bond	Bond Energy (kJ/mol)
Br – Br	193
Cl – Cl	243
Br – Cl	?

7. Aluminum metal can be recycled from scrap metal by melting the metal to evaporate impurities.

- (a) Calculate the amount of heat needed to purify 1.00 mole of Al originally at 298 K by melting it. The melting point of Al is 933 K. The molar heat capacity of Al is $24 \text{ J/(mol}\cdot\text{K)}$, and the heat of fusion of Al is 10.7 kJ/mol .


- (b) The equation for the overall process of extracting Al from Al_2O_3 is shown below. Which requires less energy, recycling existing Al or extracting Al from Al_2O_3 ? Justify your answer with a calculation.




STOP

END OF EXAM

6. A student places a mixture of plastic beads consisting of polypropylene (PP) and polyvinyl chloride (PVC) in a 1.0 L beaker containing distilled water. After stirring the contents of the beaker vigorously, the student observes that the beads of one type of plastic sink to the bottom of the beaker and the beads of the other type of plastic float on the water. The chemical structures of PP and PVC are represented by the diagrams below, which show segments of each polymer.




PP




PVC

- (a) Given that the spacing between polymer chains in PP and PVC is similar, the beads that sink are made of which polymer? Explain.

PP is synthesized from propene, C_3H_6 , and PVC is synthesized from vinyl chloride, $\text{C}_2\text{H}_3\text{Cl}$. The structures of the molecules are shown below.




Propene



Vinyl Chloride (chloroethene)

- (b) The boiling point of liquid propene (226 K) is lower than the boiling point of liquid vinyl chloride (260 K). Account for this difference in terms of the types and strengths of intermolecular forces present in each liquid.

In a separate experiment, the student measures the enthalpies of combustion of propene and vinyl chloride. The student determines that the combustion of 2.00 mol of vinyl chloride releases 2300 kJ of energy, according to the equation below.



- (c) Using the table of standard enthalpies of formation below, determine whether the combustion of 2.00 mol of propene releases more, less, or the same amount of energy that 2.00 mol of vinyl chloride releases. Justify your answer with a calculation. The balanced equation for the combustion of 2.00 mol of propene is $2 \text{C}_3\text{H}_6(g) + 9 \text{O}_2(g) \rightarrow 6 \text{CO}_2(g) + 6 \text{H}_2\text{O}(g)$.

Substance	$\text{C}_2\text{H}_3\text{Cl}(g)$	$\text{C}_3\text{H}_6(g)$	$\text{CO}_2(g)$	$\text{H}_2\text{O}(g)$	$\text{HCl}(g)$	$\text{O}_2(g)$
Standard Enthalpy of Formation (kJ/mol)	37	21	-394	-242	-92	0