


1. Answer the following questions about magnesium.


A. An incomplete mass spectrum for magnesium is shown in the diagram.



The percent abundance of magnesium-24 is 79%. The percent abundances of the other two natural isotopes of magnesium, magnesium-25 and magnesium-26, are approximately equal.

- i. Complete the mass spectrum in part A by drawing thick lines in the appropriate locations to represent the percent abundance of magnesium-25 and magnesium-26.
- ii. Describe the difference in atomic structure that accounts for the difference in mass between magnesium-25 and magnesium-26.

A student prepares a $1.85 \times 10^{-3} M$ solution of $\text{Mg}(\text{NO}_3)_2(aq)$ in beaker 1 and a $2.80 \times 10^{-4} M$ solution of $\text{NaOH}(aq)$ in beaker 2, as shown.




B. The particle diagram shown represents a magnesium ion, Mg^{2+} , in beaker 1. A sodium ion, Na^+ , in beaker 2 has a weaker attraction to water than the Mg^{2+} does. Explain this phenomenon using Coulomb's law and each of the following.


- i. The relative charge of the ions
- ii. The relative radii of the ions

C. Calculate the pH of the solution in beaker 2.

- D. A student combines 35.00 mL of $1.85 \times 10^{-3} M \text{Mg}(\text{NO}_3)_2(aq)$ with 50.00 mL of $2.80 \times 10^{-4} M \text{NaOH}(aq)$, as shown in the diagram. Calculate $[\text{Mg}^{2+}]$ after the two solutions are combined but before any reaction takes place. (Assume that volumes are additive.)





- E. The dissolution of magnesium hydroxide is represented by the following equation.




- Write the expression for the solubility product constant, K_{sp} .
 - After the two solutions are combined in beaker 3 as described in part D, but before any reaction takes place, $[\text{OH}^{-}] = 1.65 \times 10^{-4} M$. Using your answer to part D, calculate the value of the reaction quotient, Q .
 - Using the reaction quotient, Q , predict whether a precipitate should form as the mixture in beaker 3 approaches equilibrium. Justify your answer.
- F. In a separate experiment, the student adds $\text{HNO}_3(aq)$ to decrease the pH of a saturated solution containing undissolved $\text{Mg}(\text{OH})_2(s)$. Does the amount of undissolved $\text{Mg}(\text{OH})_2(s)$ increase, decrease, or remain the same as the $\text{HNO}_3(aq)$ is added? Justify your answer.

3. White phosphorus is composed of P_4 molecules with a tetrahedral structure, as shown in the diagram on the left. Each P atom is bonded to the other three P atoms by single bonds, as shown in the incomplete Lewis diagram on the right.




- A. In the box in part A, complete the Lewis diagram for P_4 by drawing the nonbonding electrons.
- B. The reaction of white phosphorus with oxygen to form $P_4O_{10}(s)$ is thermodynamically favorable at 298 K. The reaction is represented by equation 1.




- The entropy change of the reaction, ΔS° , is negative. Using particle-level reasoning, explain why the entropy decreases as the reaction progresses.
- The enthalpy change of the reaction, ΔH° , is also negative. A student claims that the favorability of the reaction is driven by enthalpy and **not** by entropy. Is the student's claim correct? Justify your answer by using the relationship between ΔG° , ΔH° , and ΔS° .

$\text{P}_4\text{O}_{10}(s)$ reacts exothermically with water to form phosphoric acid, as represented by equation 2.



A chemist uses a calorimetry experiment to determine the enthalpy change for the reaction, as represented by the following diagram.




C. The chemist carries out the calorimetry experiment and records the following information.

Mass of P_4O_{10}	0.100 g
Mass of H_2O	100.0 g
Initial temperature	22.00° C
Final temperature	22.38° C
Molar mass of P_4O_{10}	283.9 g/mol
Specific heat of H_2O	4.18 J/(g·°C)


- Calculate the amount of heat, q , released during the experiment, in kJ. Assume that the specific heat of the solution is the same as that of water.
- Calculate the value of $\Delta H^\circ_{\text{rxn}}$ for equation 2 in kJ/mol_{rxn}. Include the sign in your answer.

- D. The chemist weighed out 0.100 g P_4O_{10} of and 100.0 g of H_2O to perform a second trial. In the second trial, some of the solid P_4O_{10} stuck to the weighing paper and was not transferred to the calorimeter. Given that P_4O_{10} is the limiting reactant, would ΔT for the second trial be greater than, less than, or equal to the value in the first trial? Justify your answer.


$\text{P}_4(\text{s})$ also reacts readily with $\text{Cl}_2(\text{g})$ to produce phosphorus trichloride, $\text{PCl}_3(\text{g})$, which in turn reacts with $\text{Cl}_2(\text{g})$ in an equilibrium process to produce $\text{PCl}_5(\text{g})$. The reactions are represented by equations 3 and 4.




- E. Calculate the standard enthalpy of formation of $\text{PCl}_5(\text{g})$ represented by equation 5.



The following particle-level diagram represents the contents of the vessel in an equilibrium mixture at 546 K involving equation 4.



- F. Equation 4 for the reaction that occurs is shown.



- If each particle in the diagram represents a partial pressure of 1.00 atm, what is the value of K_p for the equilibrium mixture at 546 K?
- Does the value of K_p increase, decrease, or remain the same when the temperature is increased to 596 K? Justify your answer based on ΔH_2° .


5. Hydrogen gas and iodine gas react to form hydrogen iodide at an elevated temperature, as represented by the following equation.



(a) Write the expression for the equilibrium constant, K_c , for this reaction.

(b) $\text{H}_2(g)$ and $\text{I}_2(g)$ are added to a previously evacuated container and allowed to react.


- (i) At a certain time, the value of the reaction quotient, Q , is 0.67. The following particle diagram is an incomplete representation of the system at this time. The diagram shows the relative number of $\text{H}_2(g)$ and $\text{I}_2(g)$ molecules, but the $\text{HI}(g)$ molecules are not included. Draw the number of $\text{HI}(g)$ molecules needed to complete the diagram so that it accurately represents the system.



GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 5** on this page.

- (ii) A student monitors the number of moles of $\text{HI}(g)$ over time. Hypothesize an experimental change that could have been applied to the system in the rigid container at time t to result in the change in the number of moles of $\text{HI}(g)$ shown in the graph. Assume that the student did not add more $\text{HI}(g)$ to the system.



- (iii) After equilibrium is established, the mixture is transferred to a larger container at constant temperature. As a result, would the number of moles of $\text{HI}(g)$ increase, decrease, or remain the same? Justify your answer.

GO ON TO THE NEXT PAGE.

2. In the gas phase, AlCl_3 is a molecular substance. A reaction of gaseous AlCl_3 at high temperature is represented by the following balanced equation.



- (a) How many grams of $\text{Cl}(g)$ can be formed from 1.25 mol of $\text{AlCl}_3(g)$?


Additional reactions that involve Al or Cl are shown in the following table.

Reaction Number	Equation	ΔH_{rxn}° (kJ/mol _{rxn})
2	$\text{Al}(s) + \frac{3}{2} \text{Cl}_2(g) \rightarrow \text{AlCl}_3(g)$	-583
3	$\text{Al}(s) \rightarrow \text{Al}(g)$	+326
4	$\text{Cl}_2(g) \rightarrow 2 \text{Cl}(g)$	+243

- (b) Calculate the value of ΔH_1° , in kJ/mol_{rxn}, for reaction 1 above using reactions 2, 3, and 4.

GO ON TO THE NEXT PAGE.

(c) A potential energy diagram for Cl_2 is shown in the following graph.



(i) Based on the graph, what is the bond length, in picometers, for Cl_2 ? _____

(ii) A student finds that the average $\text{Al} - \text{Cl}$ bond length is 220 picometers and the average bond energy is 425 kJ/mol. Draw the potential energy curve for the average $\text{Al} - \text{Cl}$ bond on the preceding graph.

(d) Three proposed Lewis diagrams for the $\text{AlCl}_3(g)$ molecule are shown.




Diagram 1




Diagram 2




Diagram 3


(i) The $\text{AlCl}_3(g)$ molecule has a trigonal planar geometry. Which diagram (1, 2, or 3) can be eliminated based on geometry? Justify your choice based on VSEPR theory.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.


- (ii) Which of the three diagrams is the best representation for the bonding in AlCl_3 ? Justify your choice based on formal charges.

AlCl_3 is known to dimerize reversibly in the gas phase. The dimerization equilibrium is represented by the following equation.




- (e) Write the expression for the equilibrium constant, K_p , for this reaction.

A particle-level diagram of an equilibrium mixture of $\text{AlCl}_3(g)$ and $\text{Al}_2\text{Cl}_6(g)$ at 400°C in a 25 L closed container is shown.



- (f) Using the particle-level diagram, calculate the value of K_p for the reaction if the total pressure in the container is 22.1 atm.

GO ON TO THE NEXT PAGE.



Methanol vapor decomposes to form carbon monoxide gas and hydrogen gas at high temperatures in the presence of a platinum catalyst, as represented by the balanced chemical equation given.

(a) Are the hydrogen atoms oxidized or are they reduced in the forward reaction? Justify your answer in terms of oxidation numbers.

(b) In the following box, draw the complete Lewis electron-dot diagram for the carbon monoxide molecule in which every atom obeys the octet rule. Show all bonding and nonbonding valence electrons.



(c) The values of the standard molar entropies of the compounds involved in the reaction are given in the following table.

Substance	S° (J/(K·mol))
$\text{CH}_3\text{OH}(g)$	240.
$\text{CO}(g)$	198
$\text{H}_2(g)$	131


(i) Use the data in the table to calculate the value of the standard entropy change, ΔS° , in J/(K·mol_{rxn}), for the reaction.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.

- (ii) Calculate the value of ΔG° , in $\text{kJ/mol}_{\text{rxn}}$, for the reaction at 375 K. Assume that ΔH° and ΔS° are independent of temperature.


The following particle-level diagram shows a representative sample of the equilibrium mixture represented by the equation given.



- (d) Use information from the particle diagram to calculate the partial pressure of CO at equilibrium when the total pressure of the equilibrium mixture is 12.0 atm.
- (e) Write the expression for the equilibrium constant, K_p , for the reaction.

GO ON TO THE NEXT PAGE.

Continue your response to QUESTION 2 on this page.



The reaction system represented by the equation is allowed to achieve equilibrium at a different temperature. The following table gives the partial pressure of each species in the equilibrium mixture.

Substance	Partial Pressure at Different Temperature
$\text{CH}_3\text{OH}(g)$	2.7 atm
$\text{CO}(g)$	4.2 atm
$\text{H}_2(g)$	8.4 atm

(f) Use the information in the table to calculate the value of the equilibrium constant, K_p , at the new temperature.


(g) The volume of the container is rapidly doubled with no change in temperature. As equilibrium is re-established, does the number of moles of $\text{CH}_3\text{OH}(g)$ increase, decrease, or remain the same? Justify your answer by comparing the value of the reaction quotient, Q , with the value of the equilibrium constant, K_p .

GO ON TO THE NEXT PAGE.

CHEMISTRY**SECTION II****Time—1 hour and 45 minutes****7 Questions****YOU MAY USE YOUR CALCULATOR FOR THIS SECTION.**

Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.

For each question, show your work for each part in the space provided after that part. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.



1. Methanoic acid, HCOOH , ionizes according to the equation above.



(a) Write the expression for the equilibrium constant, K_a , for the reaction.

(b) Calculate the pH of a 0.25 M solution of HCOOH .

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

(c) In the box below, complete the Lewis electron-dot diagram for HCOOH. Show all bonding and nonbonding valence electrons.



(d) In aqueous solution, the compound H_2NNH_2 reacts according to the equation above. A 50.0 mL sample of 0.25 M $\text{H}_2\text{NNH}_2(aq)$ is combined with a 50.0 mL sample of 0.25 M $\text{HCOOH}(aq)$.


(i) Write the balanced net ionic equation for the reaction that occurs when H_2NNH_2 is combined with HCOOH .

(ii) Is the resulting solution acidic, basic, or neutral? Justify your answer.


GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

When a catalyst is added to a solution of $\text{HCOOH}(aq)$, the reaction represented by the following equation occurs.



(e) Is the reaction a redox reaction? Justify your answer.



(f) The reaction occurs in a rigid 4.3 L vessel at 25°C , and the total pressure is monitored, as shown in the graph above. The vessel originally did not contain any gas. Calculate the number of moles of $\text{CO}_2(g)$ produced in the reaction. (Assume that the amount of $\text{CO}_2(g)$ dissolved in the solution is negligible.)

(g) After the reaction has proceeded for several minutes, does the amount of catalyst increase, decrease, or remain the same? Justify your answer.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

2. Answer the following questions relating to the chemistry of the halogens.

- (a) The molecular formulas of diatomic bromine, chlorine, fluorine, and iodine are written below. Circle the formula of the molecule that has the longest bond length. Justify your choice in terms of atomic structure.



A chemistry teacher wants to prepare Br_2 . The teacher has access to the following three reagents: $\text{NaBr}(aq)$, $\text{Cl}_2(g)$, and $\text{I}_2(s)$.


Half-Reaction	E° at 25°C (V)
$\text{Br}_2 + 2 e^- \rightarrow 2 \text{Br}^-$	1.07
$\text{Cl}_2 + 2 e^- \rightarrow 2 \text{Cl}^-$	1.36
$\text{I}_2 + 2 e^- \rightarrow 2 \text{I}^-$	0.53

- (b) Using the data in the table above, write the balanced equation for the thermodynamically favorable reaction that will produce Br_2 when the teacher combines two of the reagents. Justify that the reaction is thermodynamically favorable by calculating the value of E° for the reaction.

Br_2 and Cl_2 can react to form the compound BrCl .

- (c) The boiling point of Br_2 is 332 K, whereas the boiling point of BrCl is 278 K. Explain this difference in boiling point in terms of all the intermolecular forces present between molecules of each substance.

The compound BrCl can decompose into Br_2 and Cl_2 , as represented by the balanced chemical equation below.




A 0.100 mole sample of pure $\text{BrCl}(g)$ is placed in a previously evacuated, rigid 2.00 L container at 298 K. Eventually the system reaches equilibrium according to the equation above.

- (d) Calculate the pressure in the container before equilibrium is established.
- (e) Write the expression for the equilibrium constant, K_{eq} , for the decomposition of BrCl .

After the system has reached equilibrium, 42 percent of the original BrCl sample has decomposed.

- (f) Determine the value of K_{eq} for the decomposition reaction of BrCl at 298 K.
- (g) Calculate the bond energy of the Br-Cl bond, in kJ/mol, using ΔH° for the reaction (1.6 kJ/mol_{rxn}) and the information in the following table.

Bond	Bond Energy (kJ/mol)
Br – Br	193
Cl – Cl	243
Br – Cl	?



3. At high temperatures, $\text{N}_2(\text{g})$ and $\text{O}_2(\text{g})$ can react to produce nitrogen monoxide, $\text{NO}(\text{g})$, as represented by the equation above.

- (a) Write the expression for the equilibrium constant, K_p , for the forward reaction.
- (b) A student injects $\text{N}_2(\text{g})$ and $\text{O}_2(\text{g})$ into a previously evacuated, rigid vessel and raises the temperature of the vessel to 2000°C . At this temperature the initial partial pressures of $\text{N}_2(\text{g})$ and $\text{O}_2(\text{g})$ are 6.01 atm and 1.61 atm, respectively. The system is allowed to reach equilibrium. The partial pressure of $\text{NO}(\text{g})$ at equilibrium is 0.122 atm. Calculate the value of K_p .

Nitrogen monoxide, $\text{NO}(\text{g})$, can undergo further reactions to produce acids such as HNO_2 , a weak acid with a K_a of 4.0×10^{-4} and a $\text{p}K_a$ of 3.40.

- (c) A student is asked to make a buffer solution with a pH of 3.40 by using 0.100 M $\text{HNO}_2(\text{aq})$ and 0.100 M $\text{NaOH}(\text{aq})$.
- (i) Explain why the addition of 0.100 M $\text{NaOH}(\text{aq})$ to 0.100 M $\text{HNO}_2(\text{aq})$ can result in the formation of a buffer solution. Include the net ionic equation for the reaction that occurs when the student adds the $\text{NaOH}(\text{aq})$ to the $\text{HNO}_2(\text{aq})$.
- (ii) Determine the volume, in mL, of 0.100 M $\text{NaOH}(\text{aq})$ the student should add to 100. mL of 0.100 M $\text{HNO}_2(\text{aq})$ to make a buffer solution with a pH of 3.40. Justify your answer.
- (d) A second student makes a buffer by dissolving 0.100 mol of $\text{NaNO}_2(\text{s})$ in 100. mL of 1.00 M $\text{HNO}_2(\text{aq})$. Which is more resistant to changes in pH when a strong acid or a strong base is added, the buffer made by the second student or the buffer made by the first student in part (c)? Justify your answer.
- (e) A new buffer is made using $\text{HNO}_2(\text{aq})$ as one of the ingredients. A particulate representation of a small representative portion of the buffer solution is shown below. (Cations and water molecules are not shown.) Is the pH of the buffer represented in the diagram greater than, less than, or equal to 3.40? Justify your answer.

