


4 points

Question 7: Short Answer

A For a correct circled atom.

Point 01

Accept one of the following:



(Because of resonance, the two C–O bonds on the right are equivalent.)

B (i) For the correct calculated value:

Point 02

$$K_b = \frac{[\text{HC}_2\text{H}_3\text{O}_3][\text{OH}^-]}{[\text{C}_2\text{H}_3\text{O}_3^-]} = \frac{(1.3 \times 10^{-5})(1.3 \times 10^{-5})}{(2.5 - 1.3 \times 10^{-5})} \approx \frac{(1.3 \times 10^{-5})^2}{(2.5)} = 6.8 \times 10^{-11}$$

(ii) For the correct calculated value, consistent with part B (i):

Point 03

$$K_a = \frac{K_w}{K_b} = \frac{1.0 \times 10^{-14}}{6.8 \times 10^{-11}} = 1.5 \times 10^{-4}$$

C For the correct answer and a valid justification:

Point 04

Agree. H_3O^+ is consumed in step 1 and regenerated in step 2, which is consistent with the behavior of a catalyst.

4 points

Question 5: Short Answer

(a) (i) For the correct calculated value:

1 point

$$n = \frac{PV}{RT} = \frac{(7.45 \text{ atm})(6.00 \text{ L})}{(0.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}})(296 \text{ K})} = 1.84 \text{ mol}$$

(ii) For the correct calculated value:

1 point

Accept one of the following:

• $\frac{P_1}{T_1} = \frac{P_2}{T_2}$

$$P_2 = \frac{(P_1)(T_2)}{T_1} = \frac{(7.45 \text{ atm})(271 \text{ K})}{296 \text{ K}} = 6.82 \text{ atm}$$


• $P = \frac{nRT}{V} = \frac{(1.84 \text{ mol})(0.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}})(271 \text{ K})}{6.00 \text{ L}} = 6.82 \text{ atm}$

Total for part (a) 2 points

(b) For a correct drawing:

1 point

The drawing should show three water molecules with a hydrogen atom (dark circle) oriented towards the Cl^- ion.



(c) For the correct answer and a valid justification:

1 point

HNO_2 . The diagram shows most of the molecules in their un-ionized form, indicating a weak acid with a K_a value less than 1, which is consistent with HNO_2 .

Total for question 5 4 points


Question 1: Long Answer

10 points

- (a) For the correct expression: 1 point

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{HCOO}^-]}{[\text{HCOOH}]}$$

- (b) For the correct calculated concentration of H_3O^+ : 1 point



Let $[\text{H}_3\text{O}^+] = x$, then $1.8 \times 10^{-4} = \frac{x^2}{(0.25 - x)}$


Assume $x \ll 0.25$, then $1.8 \times 10^{-4} = \frac{x^2}{0.25} \Rightarrow x = 0.0067 \text{ M}$

- For the correct calculated value of pH: 1 point


$$\text{pH} = -\log[\text{H}_3\text{O}^+] = -\log(0.0067) = 2.17$$

Total for part (b) 2 points

- (c) For the correct diagram: 1 point



- (d) (i) For the correct balanced equation (state symbols not required): 1 point



- (ii) For the correct answer and a valid justification: 1 point

Acidic. The K_a of H_2NNH_3^+ is greater than the K_b of HCOO^- , so the production of $\text{H}_3\text{O}^+(aq)$ occurs to a greater extent than the production of $\text{OH}^-(aq)$.

Total for part (d) 2 points

- (e) For the correct answer and a valid justification: 1 point

Accept one of the following:

- Yes. The oxidation number of hydrogen changes from +1 in HCOOH to zero in H_2 .
- Yes. The oxidation number of carbon changes from +2 in HCOOH to +4 in CO_2 .

- (f) For the correct calculated value of the pressure of CO_2 (may be implicit): 1 point

$$24 \text{ atm total} \times 1 \text{ atm CO}_2 / 2 \text{ atm of product} = 12 \text{ atm CO}_2$$

- For the correct calculated number of moles of CO_2 : 1 point

$$PV = nRT$$

$$n = \frac{PV}{RT} = \frac{(12 \text{ atm})(4.3 \text{ L})}{(0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1})(298 \text{ K})} = 2.1 \text{ mol CO}_2$$

Total for part (f) 2 points

- (g) For the correct answer and a valid justification: 1 point

It would remain the same. In a catalyzed reaction the net amount of catalyst is constant.


Total for question 1 10 points

AP[®] Chemistry

Scoring Guidelines

2019 SCORING GUIDELINES

Question 1




The compound urea, H_2NCONH_2 , is widely used in chemical fertilizers. The complete Lewis electron-dot diagram for the urea molecule is shown above.

- (a) Identify the hybridization of the valence orbitals of the carbon atom in the urea molecule.

 sp^2


1 point is earned for the correct answer.

- (b) Urea has a high solubility in water, due in part to its ability to form hydrogen bonds. A urea molecule and four water molecules are represented in the box below. Draw ONE dashed line (---) to indicate a possible location of a hydrogen bond between a water molecule and the urea molecule.



A dashed line should connect a hydrogen atom in water to a nitrogen or oxygen atom in urea or an oxygen atom in water to a hydrogen atom in urea. One possible correct response is shown above.

1 point is earned for a correct answer.



The dissolution of urea is represented by the equation above. A student determines that 5.39 grams of H_2NCONH_2 (molar mass 60.06 g/mol) can dissolve in water to make 5.00 mL of a saturated solution at 20.°C.

2019 SCORING GUIDELINES


Question 1 (continued)

- (c) Calculate the concentration of urea, in mol/L, in the saturated solution at 20.°C.

$5.39 \text{ g H}_2\text{NCONH}_2 \times \frac{1 \text{ mol}}{60.06 \text{ g}} = 0.0897 \text{ mol}$ $\frac{0.0897 \text{ mol}}{0.00500 \text{ L}} = 17.9 \text{ M}$	1 point is earned for the correct number of moles of urea (may be implicit). 1 point is earned for the correct molarity.
--	--

- (d) The student also determines that the concentration of urea in a saturated solution at 25°C is 19.8 M. Based on this information, is the dissolution of urea endothermic or exothermic? Justify your answer in terms of Le Chatelier's principle.

The increased solubility at the higher temperature implies that the dissolution of urea is endothermic. If a saturated solution of urea is heated, then the equilibrium system is stressed. The stress is counteracted by the endothermic dissolution of more urea.	1 point is earned for the correct answer with an appropriate justification.
---	---



- (e) The equipment shown above is provided so that the student can determine the value of the molar heat of solution for urea. Knowing that the specific heat of the solution is 4.18 J/(g·°C), list the specific measurements that are required to be made during the experiment.

mass of urea, mass of water, initial temperature of water, final temperature of solution	1 point is earned for the masses. 1 point is earned for the temperatures.
--	---

2019 SCORING GUIDELINES

Question 1 (continued)

	S° (J/(mol·K))
$\text{H}_2\text{NCONH}_2(\text{s})$	104.6
$\text{H}_2\text{NCONH}_2(\text{aq})$	?

- (f) The entropy change for the dissolution of urea,
- $\Delta S^\circ_{\text{soln}}$
- , is 70.1 J/(mol·K) at 25°C. Using the information in the table above, calculate the absolute molar entropy,
- S°
- , of aqueous urea.

$\Delta S^\circ_{\text{soln}} = S^\circ(\text{H}_2\text{NCONH}_2(\text{aq})) - S^\circ(\text{H}_2\text{NCONH}_2(\text{s}))$ $70.1 \text{ J/(mol·K)} = S^\circ(\text{H}_2\text{NCONH}_2(\text{aq})) - 104.6 \text{ J/(mol·K)}$ $S^\circ(\text{H}_2\text{NCONH}_2(\text{aq})) = 174.7 \text{ J/(mol·K)}$	1 point is earned for the correct answer.
--	---

- (g) Using particle-level reasoning, explain why
- $\Delta S^\circ_{\text{soln}}$
- is positive for the dissolution of urea in water.

Urea molecules in solution have a greater number of possible arrangements than in solid urea. This increased number of arrangements corresponds to a positive $\Delta S^\circ_{\text{soln}}$.	1 point is earned for a correct explanation.
--	--

- (h) The student claims that
- ΔS°
- for the process contributes to the thermodynamic favorability of the dissolution of urea at 25°C. Use the thermodynamic information above to support the student's claim.

Thermodynamic favorability for a process at standard conditions is determined by the sign of ΔG° , with $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$. Since ΔS° is positive, the $T\Delta S^\circ$ term makes the value of ΔG° smaller and thus makes the dissolution more thermodynamically favorable.	1 point is earned for the correct answer.
--	---

2019 SCORING GUIDELINES

Question 2

Answer the following questions relating to the chemistry of the halogens.

- (a) The molecular formulas of diatomic bromine, chlorine, fluorine, and iodine are written below. Circle the formula of the molecule that has the longest bond length. Justify your choice in terms of atomic structure.




I₂ has the longest bond length because the radius of the I atom is greater than the radii of the other halogen atoms. Thus, the distance between the nuclei of atoms in I₂ is greater than it is in smaller halogens.

1 point is earned for circling I₂ and providing a valid explanation.

A chemistry teacher wants to prepare Br₂. The teacher has access to the following three reagents: NaBr(aq), Cl₂(g), and I₂(s).

Half-Reaction	E° at 25°C (V)
Br ₂ + 2 e ⁻ → 2 Br ⁻	1.07
Cl ₂ + 2 e ⁻ → 2 Cl ⁻	1.36
I ₂ + 2 e ⁻ → 2 I ⁻	0.53

- (b) Using the data in the table above, write the balanced equation for the thermodynamically favorable reaction that will produce Br₂ when the teacher combines two of the reagents. Justify that the reaction is thermodynamically favorable by calculating the value of E° for the reaction.



1 point is earned for the correct balanced equation.

$$E^\circ = E^\circ(\text{reduced species}) - E^\circ(\text{oxidized species}) \\ = 1.36 \text{ V} - 1.07 \text{ V} = +0.29 \text{ V}.$$

Because E° for the reaction has a positive value, the reaction is thermodynamically favorable.

1 point is earned for the correct calculation of E°.

Br₂ and Cl₂ can react to form the compound BrCl.

- (c) The boiling point of Br₂ is 332 K, whereas the boiling point of BrCl is 278 K. Explain this difference in boiling point in terms of all the intermolecular forces present between molecules of each substance.

2019 SCORING GUIDELINES


Question 2 (continued)

The only intermolecular attractions in Br₂(l) are London forces, while those in BrCl(l) include both London forces and dipole-dipole forces. However, due to the greater polarizability of the electron cloud of Br₂ compared to that of BrCl, the London forces in Br₂(l) are stronger than the combined intermolecular forces in BrCl(l). Thus, the boiling point of Br₂(l) is greater than that of BrCl(l).

1 point is earned for identifying the intermolecular forces in each substance.

1 point is earned for a valid explanation.

The compound BrCl can decompose into Br₂ and Cl₂, as represented by the balanced chemical equation below.



A 0.100 mole sample of pure BrCl(g) is placed in a previously evacuated, rigid 2.00 L container at 298 K. Eventually the system reaches equilibrium according to the equation above.

- (d) Calculate the pressure in the container before equilibrium is established.

$$P = \frac{nRT}{V} = \frac{(0.100 \text{ mol})(0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1})(298 \text{ K})}{2.00 \text{ L}} = 1.22 \text{ atm}$$

1 point is earned for a correct pressure with consistent units.


- (e) Write the expression for the equilibrium constant, K_{eq}, for the decomposition of BrCl.

$$K_{eq} = \frac{[\text{Br}_2][\text{Cl}_2]}{[\text{BrCl}]^2} \quad \text{or} \quad K_{eq} = \frac{P_{\text{Br}_2} P_{\text{Cl}_2}}{(P_{\text{BrCl}})^2}$$

1 point is earned for a correct equilibrium expression.

After the system has reached equilibrium, 42 percent of the original BrCl sample has decomposed.

- (f) Determine the value of K_{eq} for the decomposition reaction of BrCl at 298 K.



I	1.22	0	0
C	-2x	+x	+x
E	0.71	0.26	0.26

$$P_{\text{BrCl decomposed}} = (0.42)(1.22 \text{ atm}) = 0.51 \text{ atm}$$

$$2x = 0.51 \text{ atm} \Rightarrow x = 0.26 \text{ atm}$$

$$K_{eq} = \frac{(0.26)(0.26)}{(0.71)^2} = 0.13$$

Note: The solution is in terms of pressures. Solutions in terms of molar concentrations also earn full credit.

1 point is earned for correct stoichiometric values at equilibrium.

1 point is earned for a consistent value of K_{eq}.

2019 SCORING GUIDELINES

Question 2 (continued)

Calculate the bond energy of the Br-Cl bond, in kJ/mol, using ΔH° for the reaction (1.6 kJ/mol information in the following table.

Bond	Bond Energy (kJ/mol)
Br – Br	193
Cl – Cl	243
Br – Cl	?

$$\Delta H^\circ = \sum (\text{bond energies})_{\text{broken}} - \sum (\text{bond energies})_{\text{formed}}$$

$$\text{J/mol} = 2(\text{Br-Cl bond energy}) - (193 \text{ kJ/mol} + 243 \text{ kJ/mol})$$

$$-36 + 1.6 \text{ kJ/mol} = 2(\text{Br-Cl bond energy})$$


$$\text{Cl bond energy} = 219 \text{ kJ/mol}$$

1 point is earned for a correct calculation of the Br-Cl bond energy.


2019 SCORING GUIDELINES

Question 3

A student is given 50.0 mL of a solution of Na_2CO_3 of unknown concentration. To determine the concentration of the solution, the student mixes the solution with excess 1.0 M $\text{Ca}(\text{NO}_3)_2(aq)$, causing a precipitate to form. The balanced equation for the reaction is shown below.




Write the net ionic equation for the reaction that occurs when the solutions of Na_2CO_3 and $\text{Ca}(\text{NO}_3)_2$ are mixed.



1 point is earned for the correct equation.

The diagram below is incomplete. Draw in the species needed to accurately represent the major ionic species remaining in the solution after the reaction has been completed.



The drawing shows one Ca^{2+} ion.

1 point is earned for drawing a Ca^{2+} ion.

2019 SCORING GUIDELINES

Question 3 (continued)

The student filters and dries the precipitate of CaCO_3 (molar mass 100.1 g/mol) and records the data in the table below.

Volume of Na_2CO_3 solution	50.0 mL
Volume of 1.0 M $\text{Ca}(\text{NO}_3)_2$ added	100.0 mL
Mass of CaCO_3 precipitate collected	0.93 g

Determine the number of moles of Na_2CO_3 in the original 50.0 mL of solution.

$0.93 \text{ g CaCO}_3 \times \frac{1 \text{ mol CaCO}_3}{100.1 \text{ g}} = 0.0093 \text{ mol CaCO}_3$	1 point is earned for the correct answer.
$0.0093 \text{ mol CaCO}_3 \times \frac{1 \text{ mol Na}_2\text{CO}_3}{1 \text{ mol CaCO}_3} = 0.0093 \text{ mol Na}_2\text{CO}_3$	


The student realizes that the precipitate was not completely dried and claims that as a result, the calculated Na_2CO_3 molarity is too low. Do you agree with the student's claim? Justify your answer.

Disagree. The presence of water in the solid will cause the measured mass of the precipitate to be greater than the actual mass of CaCO_3 . As a result, the calculated number of moles of CaCO_3 and moles of Na_2CO_3 will be greater than the actual moles present. Therefore the calculated concentration of $\text{Na}_2\text{CO}_3(aq)$ will be too high.	1 point is earned for the correct answer with valid justification.
---	--

After the precipitate forms and is filtered, the liquid that passed through the filter is tested to see if it can conduct electricity. What would be observed? Justify your answer.

The liquid conducts electricity because ions ($\text{Na}^+(aq)$, $\text{CO}_3^{2-}(aq)$, and $\text{NO}_3^-(aq)$) are present in the solution.	1 point is earned for the correct answer with valid justification.
--	--

The student decides to determine the molarity of the same Na_2CO_3 solution using a second method. When is dissolved in water, $\text{CO}_3^{2-}(aq)$ hydrolyzes to form $\text{HCO}_3^-(aq)$, as shown by the following equation.



2019 SCORING GUIDELINES

Question 3 (continued)

The student decides to first determine $[\text{OH}^-]$ in the solution, then use that result to calculate the initial concentration of $\text{CO}_3^{2-}(aq)$.

- (i) Identify a laboratory method (not titration) that the student could use to collect data to determine $[\text{OH}^-]$ in the solution.

Determine the pH of the solution using a pH meter.	1 point is earned for identifying a valid method.
--	---

- (ii) Explain how the student could use the measured value in part (f)(i) to calculate the initial concentration of $\text{CO}_3^{2-}(aq)$. (Do not do any numerical calculations.)

First determine $[\text{OH}^-]$ using $\text{pOH} = 14 - \text{pH}$, then $[\text{OH}^-] = 10^{-\text{pOH}}$.

use the K_b expression and an ICE table (see example below) to determine $[\text{CO}_3^{2-}]$ and $[\text{HCO}_3^-]$ at equilibrium. The initial concentration of CO_3^{2-} , c_i , is equal to the sum of the equilibrium concentrations of CO_3^{2-} and HCO_3^- .

	$\text{CO}_3^{2-}(aq) + \text{H}_2\text{O}(l)$	$\rightleftharpoons$	$\text{HCO}_3^-(aq) + \text{OH}^-(aq)$
	c_i	---	0
	$-x$	---	$+x$
	$c_i - x$	---	x

$$K_b = \frac{(x)(x)}{c_i - x} \Rightarrow c_i = \frac{(x)(x)}{K_b} + x$$

1 point is earned for a valid method of determining $[\text{OH}^-]$ from the measured value.

1 point is earned for a valid method of determining the initial concentration of CO_3^{2-} .

In the original Na_2CO_3 solution at equilibrium, is the concentration of $\text{HCO}_3^-(aq)$ greater than, less than, or equal to the concentration of $\text{CO}_3^{2-}(aq)$? Justify your answer.

Less than. The small value of K_b , 2.1×10^{-4} , indicates the reactants are favored.	1 point is earned for the correct answer with a valid justification.
---	--

The student needs to make a $\text{CO}_3^{2-}/\text{HCO}_3^-$ buffer. Is the Na_2CO_3 solution suitable for making a buffer with a pH of 6? Explain why or why not.

The Na_2CO_3 solution is not suitable. The $\text{p}K_a$ of HCO_3^- is 10.32. Buffers are effective when the required pH is approximately equal to the $\text{p}K_a$ of the weak acid. An acid with a $\text{p}K_a$ of 10.32 is not appropriate to prepare a buffer with a pH of 6.	1 point is earned for the correct answer with a valid explanation.
--	--

2019 SCORING GUIDELINES

Question 4

dent is doing experiments with $\text{CO}_2(g)$. Originally, a sample of the gas is in a rigid container at 299 K and 0.70 atm. The student increases the temperature of the $\text{CO}_2(g)$ in the container to 425 K.

Describe the effect of raising the temperature on the motion of the $\text{CO}_2(g)$ molecules.

average speed of the molecules increases as temperature increases.

1 point is earned for the correct answer.

Calculate the pressure of the $\text{CO}_2(g)$ in the container at 425 K.

Both the volume and the number of molecules are constant, therefore

$$\frac{P_1}{T_1} = \frac{P_2}{T_2} \Rightarrow \frac{0.70 \text{ atm}}{299 \text{ K}} = \frac{P_2}{425 \text{ K}} \Rightarrow P_2 = 0.99 \text{ atm}$$

1 point is earned for the correct answer.

In terms of kinetic molecular theory, briefly explain why the pressure of the $\text{CO}_2(g)$ in the container changes as it is heated to 425 K.

Faster-moving gas particles collide more frequently with the walls of the container, thus increasing the pressure.

ster-moving gas particles collide more forcefully with the walls of the container, thus increasing the pressure.

1 point is earned for a correct explanation.

The student measures the actual pressure of the $\text{CO}_2(g)$ in the container at 425 K and observes that it is less than the pressure predicted by the ideal gas law. Explain this observation.


The attractive forces between CO_2 molecules result in a pressure that is lower than that predicted by the ideal gas law.

1 point is earned for a correct explanation.

2019 SCORING GUIDELINES

Question 5

The complete photoelectron spectrum of an element in its ground state is represented below.



Based on the spectrum,

(i) write the ground-state electron configuration of the element, and

$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2$ or $[\text{Ar}] 4s^2$

1 point is earned for the correct answer.

(ii) identify the element.

Ca

1 point is earned for the correct answer.

Calculate the wavelength, in meters, of electromagnetic radiation needed to remove an electron from the valence shell of an atom of the element.

Energy (E) required = $0.980 \times 10^{-18} \text{ J}$

$$E = h\nu = \frac{hc}{\lambda} \Rightarrow \lambda = \frac{hc}{E}$$

$$\lambda = \frac{(6.626 \times 10^{-34} \text{ Js})(2.998 \times 10^8 \text{ ms}^{-1})}{0.980 \times 10^{-18} \text{ J}}$$

$$\lambda = 2.03 \times 10^{-7} \text{ m}$$


1 point is earned for the correct identification of the energy required to remove an electron from the valence shell (may be implicit).

1 point is earned for calculating the correct wavelength.




2019 SCORING GUIDELINES

Question 6

Nitrogen dioxide, $\text{NO}_2(\text{g})$, is produced as a byproduct of the combustion of fossil fuels in internal combustion engines. At elevated temperatures $\text{NO}_2(\text{g})$ decomposes according to the equation below.



The concentration of a sample of $\text{NO}_2(\text{g})$ is monitored as it decomposes and is recorded on the graph directly below. The two graphs that follow it are derived from the original data.



Explain how the graphs indicate that the reaction is second order.

The linear graph of $\frac{1}{[\text{NO}_2]}$ vs. time indicates a second-order reaction.

1 point is earned for the correct answer.

2019 SCORING GUIDELINES

Question 6 (continued)

Write the rate law for the decomposition of $\text{NO}_2(\text{g})$.



$$\text{rate} = k[\text{NO}_2]^2$$

1 point is earned for the correct answer.

Consider two possible mechanisms for the decomposition reaction.

- (i) Is the rate law described by mechanism I shown below consistent with the rate law you wrote in part (b)? Justify your answer.

Mechanism I




Step 1 is slow, therefore it is the rate-determining step of this mechanism. The rate law of this elementary reaction is $\text{rate} = k[\text{NO}_2][\text{NO}_2] = k[\text{NO}_2]^2$, which is consistent with the second-order rate law in part (b).

1 point is earned for the correct answer with justification.

- (ii) Is the rate law described by mechanism II shown below consistent with the rate law you wrote in part (b)? Justify your answer.

Mechanism II




Step 2 is slow; therefore, it is the rate-determining step of this mechanism. The rate law of this elementary reaction is $\text{rate} = k[\text{N}_2\text{O}_4]$. Because N_2O_4 is an intermediate, it cannot appear in the rate law of the overall reaction. Because $K_{eq} = \frac{[\text{N}_2\text{O}_4]}{[\text{NO}_2]^2}$ in step 1, $[\text{N}_2\text{O}_4] = K_{eq}[\text{NO}_2]^2$. Then, substituting $K_{eq}[\text{NO}_2]^2$ for $[\text{N}_2\text{O}_4]$ in the rate law of step 2 gives $\text{rate} = (k K_{eq})[\text{NO}_2]^2$, which is consistent with the rate law in part (b).

1 point is earned for the correct answer with justification.

2019 SCORING GUIDELINES

Question 7




A student dissolved a 0.139 g sample of oxalic acid, $\text{H}_2\text{C}_2\text{O}_4$, in water in an Erlenmeyer flask. Then the student titrated the $\text{H}_2\text{C}_2\text{O}_4$ solution in the flask with a solution of KMnO_4 , which has a dark purple color. The balanced chemical equation for the reaction that occurred during the titration is shown above.

Identify the species that was reduced in the titration reaction. Justify your answer in terms of oxidation numbers.

MnO_4^- is reduced to Mn^{2+} as the oxidation number of Mn changes from +7 to +2, indicating a gain of 5 electrons.

1 point is earned for the correct answer with justification.

The student used a 50.0 mL buret to add the $\text{KMnO}_4(aq)$ to the $\text{H}_2\text{C}_2\text{O}_4(aq)$ until a faint lavender color was observed in the flask, an indication that the end point of the titration had been reached. The initial and final volume readings of the solution in the buret are shown below. Write down the initial reading and the final reading and use them to determine the volume of $\text{KMnO}_4(aq)$ that was added during the titration.



$$29.55 \text{ mL} - 3.35 \text{ mL} = 26.20 \text{ mL}$$

1 point is earned for the correct answer.

2019 SCORING GUIDELINES

Question 7 (continued)

Given that the concentration of $\text{KMnO}_4(aq)$ was 0.0235 M, calculate the number of moles of MnO_4^- ions that completely reacted with the $\text{H}_2\text{C}_2\text{O}_4$.

$$(0.02620 \text{ L})(0.0235 \text{ mol/L}) = 0.000616 \text{ mol}$$

1 point is earned for the correct answer.

The student proposes to perform another titration using a 0.139 g sample of $\text{H}_2\text{C}_2\text{O}_4$, but this time using 0.00143 M $\text{KMnO}_4(aq)$ in the buret. Would this titrant concentration be a reasonable choice to use if the student followed the same procedure and used the same equipment as before? Justify your response.

No. The 0.00143 M titrant solution is so diluted that volume of titrant needed to reach the end point would be much greater than the 50 mL capacity of the buret.

1 point is earned for the correct answer with appropriate justification.

2018


AP[®] CollegeBoard

AP Chemistry

Scoring Guidelines

2018 SCORING GUIDELINES

Question 1



A student performs an experiment to determine the value of the enthalpy change, $\Delta H_{\text{rxn}}^\circ$, for the oxidation-reduction reaction represented by the balanced equation above.

Determine the oxidation number of Cl in NaOCl.

+1

1 point is earned for the correct answer.

Calculate the number of grams of $\text{Na}_2\text{S}_2\text{O}_3$ needed to prepare 100.00 mL of 0.500 M $\text{Na}_2\text{S}_2\text{O}_3$.

$$100.00 \text{ mL} \times \frac{0.500 \text{ mol Na}_2\text{S}_2\text{O}_3}{1000 \text{ mL}} \times \frac{158.10 \text{ g Na}_2\text{S}_2\text{O}_3}{1 \text{ mol Na}_2\text{S}_2\text{O}_3}$$

$$= 7.90 \text{ g Na}_2\text{S}_2\text{O}_3$$

1 point is earned for the correct number of moles of $\text{Na}_2\text{S}_2\text{O}_3$ (may be implicit).

1 point is earned for the correct calculation of mass of $\text{Na}_2\text{S}_2\text{O}_3$ consistent with the number of moles.

In the experiment, the student uses the solutions shown in the table below.

Solution	Concentration (M)	Volume (mL)
$\text{Na}_2\text{S}_2\text{O}_3(aq)$	0.500	5.00
$\text{NaOCl}(aq)$	0.500	5.00
$\text{NaOH}(aq)$	0.500	5.00

Using the balanced equation for the oxidation-reduction reaction and the information in the table above, determine which reactant is the limiting reactant. Justify your answer.

NaOCl is the limiting reactant.


Given that equal numbers of moles of each reactant were present initially, it follows from the coefficients of the reactants in the balanced equation that NaOCl will be depleted first.

1 point is earned for identifying limiting reactant and providing a valid justification.

2018 SCORING GUIDELINES

Question 1 (continued)

The solutions, all originally at 20.0°C, are combined in an insulated calorimeter. The temperature of the reaction mixture is monitored, as shown in the graph below.



According to the graph, what is the temperature change of the reaction mixture?

From the graph the final temperature is 32.5°C.

$$\Delta T = T_f - T_i = 32.5^\circ\text{C} - 20.0^\circ\text{C} = 12.5^\circ\text{C}$$

1 point is earned for the correct value ΔT

The mass of the reaction mixture inside the calorimeter is 15.21 g.

- (i) Calculate the magnitude of the heat energy, in joules, that is released during the reaction. Assume that the specific heat of the reaction mixture is 3.94 J/(g·°C) and that the heat absorbed by the calorimeter is negligible.

$$q = mc\Delta T$$

$$= (15.21 \text{ g})(3.94 \text{ J/(g}\cdot^\circ\text{C)})(12.5^\circ\text{C}) = 749 \text{ J}$$

1 point is earned for the correct calculation q consistent with the ΔT value from part (i)

- (ii) Using the balanced equation for the oxidation-reduction reaction and your answer to part (c), calculate the value of the enthalpy change of the reaction, $\Delta H_{\text{rxn}}^\circ$, in kJ/mol_{rxn}. Include the appropriate algebraic sign with your answer.

2018 SCORING GUIDELINES

Question 1 (continued)

$\text{NaOCl} = 5.00 \text{ mL} \times \frac{0.500 \text{ mol NaOCl}}{1000 \text{ mL NaOCl}} = 0.00250 \text{ mol NaOCl}$ $= 0.00250 \text{ mol NaOCl} \times \frac{1 \text{ mol}_{\text{rxn}}}{4 \text{ mol NaOCl}} = 0.000625 \text{ mol}_{\text{rxn}}$ $\Delta H_{\text{rxn}}^{\circ} = \frac{-0.749 \text{ kJ}}{0.000625 \text{ mol}_{\text{rxn}}} = -1.20 \times 10^3 \text{ kJ/mol}_{\text{rxn}}$	1 point is earned for correctly calculating the value of mol consistent with the limiting reactant in part 1 point is earned for a negative $\Delta H_{\text{rxn}}^{\circ}$ obtained by dividing the calculated value of q by the calculated value of mol
---	---

The student repeats the experiment, but this time doubling the volume of each of the reactants, as shown in the table below.

Solution	Concentration (M)	Volume (mL)
$\text{Na}_2\text{S}_2\text{O}_3(aq)$	0.500	10.0
$\text{NaOCl}(aq)$	0.500	10.0
$\text{NaOH}(aq)$	0.500	10.0

The magnitude of the enthalpy change, $\Delta H_{\text{rxn}}^{\circ}$, in $\text{kJ/mol}_{\text{rxn}}$, calculated from the results of the second experiment is the same as the result calculated in part (e)(ii). Explain this result.


By doubling the volumes, the number of moles of the reactants are doubled, which doubles the amount of energy produced. Therefore the amount of heat per mole will remain the same. OR In the second experiment, $\Delta H_{\text{rxn}}^{\circ} = \frac{2mc\Delta T}{2n} = \frac{mc\Delta T}{n} = \Delta H_{\text{rxn}}^{\circ}$. Thus the magnitude is the same as calculated in the first experiment.	1 point is earned for a valid explanation.
--	---

Write the balanced net ionic equation for the given reaction.

$\text{S}_3^{2-}(aq) + 4 \text{ OCl}^{-}(aq) + 2 \text{ OH}^{-}(aq) \rightarrow 2 \text{ SO}_4^{2-}(aq) + 4 \text{ Cl}^{-}(aq) + \text{H}_2\text{O}(l)$	1 point is earned for the correct net ionic equation.
---	--


2018 SCORING GUIDELINES

Question 2




A student investigates the reactions of nitrogen oxides. One of the reactions in the investigation requires an equimolar mixture of $\text{NO}(g)$ and $\text{NO}_2(g)$, which the student produces by using the reaction represented above.

- () The particle-level representation of the equimolar mixture of $\text{NO}(g)$ and $\text{NO}_2(g)$ in the flask at the completion of the reaction between $\text{NO}(g)$ and $\text{O}_2(g)$ is shown below in the box on the right. In the box below on the left, draw the particle-level representation of the reactant mixture of $\text{NO}(g)$ and O_2 would yield the product mixture shown in the box on the right. In your drawing, represent oxygen atoms and nitrogen atoms as indicated below.



See sample student response above. (8 molecules of NO and 2 molecules of O_2)	1 point is earned for correctly representing molecules of NO and O 1 point is earned for correctly representing atom conservation.
--	--

The student reads in a reference text that $\text{NO}(g)$ and $\text{NO}_2(g)$ will react as represented by the equation below. Thermodynamic data for the reaction are given in the table below the equation.



ΔH_{298}°	ΔS_{298}°	ΔG_{298}°
$-40.4 \text{ kJ/mol}_{\text{rxn}}$	$-138.5 \text{ J/(K}\cdot\text{mol}_{\text{rxn}})$	$0.87 \text{ kJ/mol}_{\text{rxn}}$

2018 SCORING GUIDELINES

Question 2 (continued)

The student begins with an equimolar mixture of NO(g) and $\text{NO}_2\text{(g)}$ in a rigid reaction vessel and the mixture reaches equilibrium at 298 K.

- (i) Calculate the value of the equilibrium constant, K , for the reaction at 298 K.

$$\begin{aligned}\Delta G^\circ &= -RT \ln K \\ K &= e^{-\Delta G^\circ / RT} \\ &= e^{-\frac{870 \text{ J/mol}}{(8.314 \text{ J mol}^{-1} \text{ K}^{-1})(298 \text{ K})}} \\ K &= 0.70\end{aligned}$$

1 point is earned for a correct calculation of K .

- (ii) If both P_{NO} and P_{NO_2} in the vessel are initially 1.0 atm, will $P_{\text{N}_2\text{O}_3}$ at equilibrium be equal to 1.0 atm? Justify your answer.

No, the pressure will not equal 1.0 atm.

$P_{\text{N}_2\text{O}_3}$ would only equal 1.0 atm if the reaction goes to completion.

The value of K indicates that a substantial amount of reactants will be present at equilibrium.

1 point is earned for a correct choice and valid justification based on the value of


The student hypothesizes that increasing the temperature will increase the amount of $\text{N}_2\text{O}_3\text{(g)}$ in the equilibrium mixture. Indicate whether you agree or disagree with the hypothesis. Justify your answer.

Disagree.

Because the reaction is exothermic, increasing the temperature of the reaction will favor the formation of the reactants (according to Le Chatelier's principle).

1 point is earned for the correct choice and a correct justification.

g) reacts with water to form nitrous acid, $\text{HNO}_2\text{(aq)}$, a compound involved in the production of acid rain. The reaction is represented below.




2018 SCORING GUIDELINES

Question 2 (continued)

The skeletal structure of the HNO_2 molecule is shown in the box below.

- (i) Complete the Lewis electron-dot diagram of the HNO_2 molecule in the box below, including any lone pairs of electrons.



See sample response above.
(Line segments can be used to represent electron pairs.)


1 point is earned for a valid diagram.

- (ii) Based on your completed diagram above, identify the hybridization of the nitrogen atom in the HNO_2 molecule.


sp^2

1 point is earned for the correct answer.

To produce an aqueous solution of HNO_2 , the student bubbles $\text{N}_2\text{O}_3\text{(g)}$ into distilled water. Assume that the reaction goes to completion and that HNO_2 is the only species produced. To determine the concentration of $\text{HNO}_2\text{(aq)}$ in the resulting solution, the student titrates a 100. mL sample of the solution with 0.100 KOH(aq) . The neutralization reaction is represented below.



The following titration curve shows the change in pH of the solution during the titration.



2018 SCORING GUIDELINES

Question 2 (continued)

Use the titration curve and the information above to

- (i) determine the initial concentration of the $\text{HNO}_2(aq)$ solution

$\text{mL KOH} \times \frac{0.100 \text{ mol KOH}}{1000 \text{ mL KOH}} = 0.0020 \text{ mol KOH added}$ $\Rightarrow 0.0020 \text{ mol HNO}_2 \text{ in } 100. \text{ mL of solution because the stoichiometry of the neutralization reaction is 1 to 1.}$ $\frac{0.0020 \text{ mol HNO}_2}{0.100 \text{ L}} = 0.020 \text{ M HNO}_2$	1 point is earned for the correct calculation of the initial concentration.
---	---

- (ii) estimate the value of $\text{p}K_a$ for $\text{HNO}_2(aq)$

The value of $\text{p}K_a$ is about 3.4.	1 point is earned for an acceptable estimate for the value of $\text{p}K_a$.
--	---

During the titration, after a volume of 15 mL of 0.100 M $\text{KOH}(aq)$ has been added, which species, $\text{HNO}_2(aq)$ or $\text{NO}_2^-(aq)$, is present at a higher concentration in the solution? Justify your answer.

(aq) The titration is past the half-equivalence point; therefore, there will be more conjugate base present than acid.	1 point is earned for the correct choice <u>and</u> a valid justification.
--	--

2018 SCORING GUIDELINES

Question 3

Answer the following questions relating to Fe and its ions, Fe^{2+} and Fe^{3+} .

Write the ground-state electron configuration of the Fe^{2+} ion.

$1s^2 2s^2 2p^6 3s^2 3p^6 3d^6$ OR $[\text{Ar}] 3d^6$	1 point is earned for a correct electron configuration.
---	---

Ion	Ionic Radius (pm)
Fe^{2+}	92
Fe^{3+}	79


The radii of the ions are given in the table above. Using principles of atomic structure, explain why the radius of the Fe^{2+} ion is larger than the radius of the Fe^{3+} ion.

Both ions have the same nuclear charge; however, the greater number of electrons in the outermost shell of Fe^{2+} results in greater electron-electron repulsion within that shell, leading to a larger radius.	1 point is earned for a valid explanation.
---	--

Fe^{3+} ions interact more strongly with water molecules in aqueous solution than Fe^{2+} ions do. Give one reason for this stronger interaction, and justify your answer using Coulomb's law.

Coulomb's law: $F \propto \frac{q_1 q_2}{r^2}$ (need not be explicitly stated) In comparison to the Fe^{2+} ion, the Fe^{3+} ion has a higher charge. The smaller size of Fe^{3+} allows it to get closer to a water molecule.	1 point is earned for a valid explanation.
--	--

A student obtains a solution that contains an unknown concentration of $\text{Fe}^{2+}(aq)$. To determine the concentration of $\text{Fe}^{2+}(aq)$ in the solution, the student titrates a sample of the solution with MnO_4^- which converts $\text{Fe}^{2+}(aq)$ to $\text{Fe}^{3+}(aq)$, as represented by the following equation.



Write the balanced equation for the half-reaction for the oxidation of $\text{Fe}^{2+}(aq)$ to $\text{Fe}^{3+}(aq)$.

$\text{Fe}^{2+}(aq) \rightarrow \text{Fe}^{3+}(aq) + e^-$	1 point is earned for the correct half-reaction.
---	--

2018 SCORING GUIDELINES

Question 3 (continued)

The student titrates a 10.0 mL sample of the $\text{Fe}^{2+}(\text{aq})$ solution. Calculate the value of $[\text{Fe}^{2+}]$ in solution if it takes 17.48 mL of added 0.0350 M $\text{KMnO}_4(\text{aq})$ to reach the equivalence point of the titration.

$17.48 \text{ mL} \times \frac{0.0350 \text{ mol KMnO}_4}{1000 \text{ mL}} = 0.000612 \text{ mol KMnO}_4$ $0.000612 \text{ mol KMnO}_4 \times \frac{5 \text{ mol Fe}^{2+}}{1 \text{ mol KMnO}_4} = 0.003059 \text{ mol Fe}^{2+}$ $\frac{0.003059 \text{ mol Fe}^{2+}}{0.0100 \text{ L}} = 0.306 \text{ M Fe}^{2+}$	1 point is earned for calculating the number of moles of KMnO_4 (may be implicit). 1 point is earned for the correct concentration of Fe^{2+}.
--	--

To deliver the 10.0 mL sample of the $\text{Fe}^{2+}(\text{aq})$ solution in part (e), the student has the choice of using one of the pieces of glassware listed below.

- 25 mL buret
- 25 mL beaker
- 25 mL graduated cylinder
- 25 mL volumetric flask

Explain why the 25 mL volumetric flask would be a poor choice to use for delivering the required volume of the $\text{Fe}^{2+}(\text{aq})$ solution.

The volumetric flask is designed to contain only 25.00 mL precisely.	1 point is earned for a valid explanation.
--	--

2018 SCORING GUIDELINES

Question 3 (continued)

In a separate experiment, the student is given a sample of powdered $\text{Fe}(\text{s})$ that contains an inert impurity. The student uses a procedure to oxidize the $\text{Fe}(\text{s})$ in the sample to $\text{Fe}_2\text{O}_3(\text{s})$. The student collects the following data during the experiment.

Mass of $\text{Fe}(\text{s})$ with inert impurity	6.724 g
Mass of $\text{Fe}_2\text{O}_3(\text{s})$ produced	7.531 g

Calculate the number of moles of Fe in the $\text{Fe}_2\text{O}_3(\text{s})$ produced.

$7.531 \text{ g Fe}_2\text{O}_3 \times \frac{1 \text{ mol Fe}_2\text{O}_3}{159.70 \text{ g Fe}_2\text{O}_3} = 0.04716 \text{ mol Fe}_2\text{O}_3$ $0.04716 \text{ mol Fe}_2\text{O}_3 \times \frac{2 \text{ mol Fe}}{1 \text{ mol Fe}_2\text{O}_3} = 0.09431 \text{ mol Fe}$	1 point is earned for correct calculation.
--	--

Calculate the percent by mass of Fe in the original sample of powdered $\text{Fe}(\text{s})$ with the inert impurity.


$0.09431 \text{ mol Fe} \times \frac{55.85 \text{ g Fe}}{1 \text{ mol}} = 5.267 \text{ g Fe}$ $\frac{5.267 \text{ g Fe}}{6.724 \text{ g sample}} \times 100 = 78.33\%$	1 point is earned for correct calculation of the mass percent based on the answer to part (g).
--	--

If the oxidation of the $\text{Fe}(\text{s})$ in the original sample was incomplete so that some of the 7.531 g of product was $\text{FeO}(\text{s})$ instead of $\text{Fe}_2\text{O}_3(\text{s})$, would the calculated mass percent of $\text{Fe}(\text{s})$ in the original sample be higher, lower, or the same as the actual mass percent of $\text{Fe}(\text{s})$? Justify your answer.

The calculated mass percent of Fe would be lower than the actual mass percent of Fe. A sample that contains any FeO (rather than Fe_2O_3) will have a lower <u>actual</u> mass percent of Fe than a completely oxidized sample would have. Therefore, when the moles of Fe are calculated (assuming all the mass of the sample is Fe_2O_3) the <u>calculated</u> number of moles of Fe, and hence the <u>calculated</u> mass percent of Fe, will be lower.	1 point is earned for the correct answer and a valid explanation.
--	---

2018 SCORING GUIDELINES

Question 4



Compound	Molecular Structure	Boiling Point at 1 atm (K)
CS ₂		319
COS		223

The table above gives the molecular structures and boiling points for the compounds CS₂ and COS.

In terms of the types and relative strengths of all the intermolecular forces in each compound, explain why the boiling point of CS₂(l) is higher than that of COS(l).

CS₂ has only London dispersion forces, while COS has London dispersion forces and dipole-dipole forces.

The London dispersion forces in CS₂ are stronger than the combination of London dispersion forces and dipole-dipole forces in COS.

1 point is earned for correctly identifying all of the intermolecular forces in **both** molecules.

1 point is earned for a valid explanation.

A 10.0 g sample of CS₂(l) is put in an evacuated 5.0 L rigid container. The container is sealed and heated to 325 K, at which temperature all of the CS₂(l) has vaporized. What is the pressure in the container once all of the CS₂(l) has vaporized?

$$10.0 \text{ g CS}_2 \times \frac{1 \text{ mol CS}_2}{76.13 \text{ g CS}_2} = 0.131 \text{ mol CS}_2$$

$$\therefore \frac{nRT}{V} = \frac{(0.131 \text{ mol})(0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1})(325 \text{ K})}{5.0 \text{ L}}$$


$$= 0.70 \text{ atm}$$

1 point is earned for the correct number of moles of

1 point is earned for the correct calculation of pressure with appropriate units.

2018 SCORING GUIDELINES

Question 5



The ionization of HF(aq) in water is represented by the equation above. In a 0.0350 M HF(aq) solution, the percent ionization of HF is 13.0 percent.

Two particulate representations of the ionization of HF molecules in the 0.0350 M HF(aq) solution shown below in Figure 1 and Figure 2. Water molecules are not shown. Explain why the representation of the ionization of HF molecules in water in Figure 1 is more accurate than the representation in Figure 2. (The key below identifies the particles in the representations.)




Figure 1




Figure 2

HF is a weak acid and is only partially ionized. This fact is consistent with Figure 1, which shows that one out of eight (~13%) HF molecules is ionized (to form one H₃O⁺ and one F⁻).


1 point is earned for a valid explanation.

Figure 2 cannot represent HF because it represents 100% ionization of the acid.

Use the percent ionization data above to calculate the value of K_a for HF.

Assume $[\text{H}_3\text{O}^+] = [\text{F}^-]$ in HF(aq).

$$\frac{[\text{H}_3\text{O}^+]}{0.0350 \text{ M}} = 0.130 \Rightarrow [\text{H}_3\text{O}^+] = 0.00455 \text{ M}$$



0.0350	0	~0
-0.00455	+0.00455	+0.00455
0.0304	0.00455	0.00455

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{F}^-]}{[\text{HF}]} = \frac{(0.00455)^2}{(0.0304)} = 6.81 \times 10^{-4}$$

1 point is earned for the correct calculation of $[\text{H}_3\text{O}^+]$.

1 point is earned for a value of K_a consistent with the calculated value of $[\text{H}_3\text{O}^+]$.

2018 SCORING GUIDELINES

Question 5 (continued)

If 50.0 mL of distilled water is added to 50.0 mL of 0.035 M $\text{HF}(aq)$, will the percent ionization $\text{HF}(aq)$ in the solution increase, decrease, or remain the same? Justify your answer with an explanation or calculation.

The percent ionization of HF in the solution would increase.

Doubling the volume of the solution decreases the initial concentration of each species by one-half; therefore,

$$\frac{(\frac{1}{2}[\text{H}_3\text{O}^+]_i)(\frac{1}{2}[\text{F}^-]_i)}{\frac{1}{2}[\text{HF}]_i} = \frac{1}{2}K_a \Rightarrow Q < K_a.$$

Consequently the equilibrium position will shift toward the products and increase the percent ionization.

New volume = twice original volume, thus new $[\text{HF}]_i = \frac{0.035}{2} = 0.0175 M$

$$= \frac{[\text{H}_3\text{O}^+][\text{F}^-]}{[\text{HF}]} = 6.81 \times 10^{-4} \text{ (value from part (b))}$$

Let $[\text{H}_3\text{O}^+] = [\text{F}^-] = x$

$$\text{Then } 6.81 \times 10^{-4} = \frac{(x)(x)}{(0.0175 - x)} \approx \frac{x^2}{(0.0175)} \Rightarrow x \approx 0.00345 M$$


$$\text{Percent ionization} = \frac{0.00345 M}{0.0175 M} \times 100 = 20.0\%$$

$\% > 13.0\%$; therefore, the percent ionization increases.

1 point is earned for correct answer valid explanation or calculation.

2018 SCORING GUIDELINES

Question 6



A student sets up a galvanic cell at 298 K that has an electrode of $\text{Ag}(s)$ immersed in a 1.0 M solution of $\text{Ag}^+(aq)$ and an electrode of $\text{Cr}(s)$ immersed in a 1.0 M solution of $\text{Cr}^{3+}(aq)$, as shown in the diagram above.

The student measures the voltage of the cell shown above and discovers that it is zero. Identify the missing component of the cell, and explain its importance for obtaining a nonzero voltage.

alt bridge is missing. The salt bridge allows for the migration of ions to maintain charge balance in each half-cell.

1 point is earned for the correct answer and a valid explanation.

Half-Reaction	E° (V)
$\text{Ag}^+(aq) + e^- \rightarrow \text{Ag}(s)$	+0.80
$\text{Cr}^{3+}(aq) + 3e^- \rightarrow \text{Cr}(s)$	?


The student adds the missing component to the cell and measures E°_{cell} to be +1.54 V. As the cell operates, Ag^+ ions are reduced. Use this information and the information in the table above to do the following.

(i) Calculate the value of E° for the half-reaction $\text{Cr}^{3+}(aq) + 3e^- \rightarrow \text{Cr}(s)$.

$$\begin{aligned} E^\circ_{\text{cell}} &= E^\circ_{\text{red}}(\text{cathode}) - E^\circ_{\text{red}}(\text{anode}) \\ +1.54 \text{ V} &= +0.80 \text{ V} - x \\ &= +0.80 \text{ V} - (+1.54 \text{ V}) = -0.74 \text{ V} \end{aligned}$$

1 point is earned for a correct calculation of E°_{red} .

(ii) Write the balanced net-ionic equation for the overall reaction that occurs as the cell operates.



1 point is earned for the correctly balanced equation.

2018 SCORING GUIDELINES

Question 6 (continued)

(iii) Calculate the value of ΔG° for the overall cell reaction in $\text{J/mol}_{\text{rxn}}$.


$$\Delta G^\circ = -nFE^\circ = -\left(\frac{3 \text{ mol } e^-}{1 \text{ mol}_{\text{rxn}}}\right)\left(96,485 \frac{\text{C}}{\text{mol } e^-}\right)\left(1.54 \frac{\text{J}}{\text{C}}\right)$$

$$= -4.46 \times 10^5 \text{ J/mol}_{\text{rxn}}$$

1 point is earned for the correct calculation of the value of Δ

2018 SCORING GUIDELINES

Question 7



The complete photoelectron spectrum of an element is represented above.

Identify the element.

The element is nitrogen, N.

1 point is earned for correctly identifying the element.

A radioactive isotope of the element decays with a half-life of 10. minutes.

Calculate the value of the rate constant, k , for the radioactive decay. Include units with your answer.

$$k = \frac{0.693}{t_{1/2}} = \frac{0.693}{10. \text{ min}} = 0.069 \text{ min}^{-1}$$

1 point is earned for the correct numerical answer.

1 point is earned for the correct unit.

If 64 atoms of the radioactive isotope are originally present in a sample, what is the expected amount of time that will pass until only one atom of the isotope remains? Show how you arrived at your answer.

$$64 \rightarrow 32 \rightarrow 16 \rightarrow 8 \rightarrow 4 \rightarrow 2 \rightarrow 1$$

6 half-lives are required.

$$6 \times 10. \text{ min} = 60. \text{ min}$$

OR

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$t = \frac{\ln(1) - \ln(64)}{-0.069 \text{ min}^{-1}} = 60. \text{ min}$$

1 point is earned for the correct answer and a valid method.

AP[®] Chemistry

2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

SCORING GUIDELINES

Question 1

A student investigates the enthalpy of solution, ΔH_{soln} , for two alkali metal halides, LiCl and NaCl. In addition to the salts, the student has access to a calorimeter, a balance with a precision of ± 0.1 g, and a thermometer with a precision of $\pm 0.1^\circ\text{C}$.

- (a) To measure ΔH_{soln} for LiCl, the student adds 100.0 g of water initially at 15.0°C to a calorimeter and adds 10.0 g of LiCl(s), stirring to dissolve. After the LiCl dissolves completely, the maximum temperature reached by the solution is 35.6°C .
- (i) Calculate the magnitude of the heat absorbed by the solution during the dissolution process, assuming that the specific heat capacity of the solution is $4.18 \text{ J/(g}\cdot^\circ\text{C)}$. Include units with your answer.

$$q = mc\Delta T = (110.0 \text{ g})(4.18 \text{ J/(g}\cdot^\circ\text{C)})(35.6^\circ\text{C} - 15.0^\circ\text{C})$$

$$= 9,470 \text{ J} = 9.47 \text{ kJ}$$

1 point is earned for the correct setup.

1 point is earned for the correct answer with units.

- (ii) Determine the value of ΔH_{soln} for LiCl in kJ/mol_{rxn} .

$$10.0 \text{ g LiCl} \times \frac{1 \text{ mol LiCl}}{42.39 \text{ g LiCl}} = 0.236 \text{ mol LiCl}$$

$$\frac{-9.47 \text{ kJ}}{0.236 \text{ mol LiCl}} = -40.1 \text{ kJ/mol}_{rxn}$$


1 point is earned for the number of moles of LiCl.

1 point is earned for the correct ΔH_{soln} and the correct sign.

To explain why ΔH_{soln} for NaCl is different than that for LiCl, the student investigates factors that affect ΔH_{soln} and finds that ionic radius and lattice enthalpy (which can be defined as the ΔH associated with the separation of a solid crystal into gaseous ions) contribute to the process. The student consults references and collects the data shown in the table below.

Ion	Ionic Radius (pm)
Li ⁺	76
Na ⁺	102

- (b) Write the complete electron configuration for the Na⁺ ion in the ground state.



1 point is earned for the complete correct configuration.

SCORING GUIDELINES

Question 1 (continued)

- (c) Using principles of atomic structure, explain why the
- Na^+
- ion is larger than the
- Li^+
- ion.

The valence electrons in the Na^+ ion are in a higher principal energy level than the valence electrons in the Li^+ ion. Electrons in higher principal energy levels are, on average, farther from the nucleus.


1 point is earned for a correct explanation based on occupied principal energy levels.

- (d) Which salt,
- LiCl
- or
- NaCl
- , has the greater lattice enthalpy? Justify your answer.

LiCl . Because the Li^+ ion is smaller than the Na^+ ion, the Coulombic attractions between ions in LiCl are stronger than in NaCl . This results in a greater lattice enthalpy.

1 point is earned for the correct choice and justification.

- (e) Below is a representation of a portion of a crystal of
- LiCl
- . Identify the ions in the representation by writing the appropriate formulas (
- Li^+
- or
- Cl^-
-) in the boxes below.



See diagram above.

1 point is earned for both identifications.

- (f) The lattice enthalpy of
- LiCl
- is positive, indicating that it takes energy to break the ions apart. However, the dissolution of
- LiCl
- in water is an exothermic process. Identify all particle-particle interactions that contribute significantly to the exothermic dissolution process being exothermic. For each interaction, include the particles that interact and the specific type of intermolecular force between those particles.

There are interactions between Li^+ ions and polar water molecules and between Cl^- ions and polar water molecules.


These are ion-dipole interactions.

1 point is earned for identifying the particles that interact.

1 point is earned for correctly identifying the type of interaction.

2016 SCORING GUIDELINES

Question



A student designs an experiment to study the reaction between NaHCO_3 and $\text{HC}_2\text{H}_3\text{O}_2$. The reaction is represented by the equation above. The student places 2.24 g of NaHCO_3 in a flask and adds 60.0 mL of 0.875 M $\text{HC}_2\text{H}_3\text{O}_2$. The student observes the formation of bubbles and that the flask gets cooler as reaction proceeds.

Identify the reaction represented above as an acid-base reaction, precipitation reaction, or redox reaction. Justify your answer.

It is an acid-base reaction. The weak acid $\text{HC}_2\text{H}_3\text{O}_2$ reacts with the weak base HCO_3^- with $\text{HC}_2\text{H}_3\text{O}_2$ donating a proton.

1 point is earned for identifying the reaction as acid-base.

It is an acid-base reaction. No solid precipitates, so it is not a precipitation reaction. None of the oxidation numbers change, so it is not a redox reaction.

1 point is earned for the justification.

Based on the information above, identify the limiting reactant. Justify your answer with calculations.

$$2.24 \text{ g NaHCO}_3 \times \frac{1 \text{ mol NaHCO}_3}{84.0 \text{ g}} = 0.0267 \text{ mol NaHCO}_3$$

1 point is earned for calculating the number of moles of each reactant.

$$60.0 \text{ mL} \times \frac{0.875 \text{ mol HC}_2\text{H}_3\text{O}_2}{1000 \text{ mL}} = 0.0525 \text{ mol HC}_2\text{H}_3\text{O}_2$$

The $\text{NaHCO}_3(s)$ and $\text{HC}_2\text{H}_3\text{O}_2(aq)$ react in a 1:1 ratio, so the limiting reactant is $\text{NaHCO}_3(s)$.

1 point is earned for identifying the limiting reactant consistent with the calculations.

The student observes that the bubbling is rapid at the beginning of the reaction and gradually slows as the reaction continues. Explain this change in the reaction rate in terms of the collisions between reactant particles.

As the reaction proceeds, both reactants are consumed and their concentrations decrease. Collisions between reactant particles become less likely as their concentrations decrease, thus the reaction rate slows.

1 point is earned for a valid explanation.

In thermodynamic terms, a reaction can be driven by enthalpy, entropy, or both.

- (i) Considering that the flask gets cooler as the reaction proceeds, what drives the chemical reaction between
- $\text{NaHCO}_3(s)$
- and
- $\text{HC}_2\text{H}_3\text{O}_2(aq)$
- ? Answer by drawing a circle around one of the choices below.

Enthalpy only

Entropy only

Both enthalpy and entropy

Entropy only should be circled.

1 point is earned for circling Entropy only.

2016 SCORING GUIDELINES

Question 2 (continued)


(ii) Justify your selection in part (d)(i) in terms of ΔG° .

$$\Delta = \Delta H^\circ - T\Delta S^\circ$$

Reactions are thermodynamically favorable when ΔG° is negative. Since the reaction is endothermic (the flask gets cooler, ΔH° is positive), the reaction is not driven by enthalpy, because enthalpy does not help make ΔG° negative. Because there are no gases in the reactants and one of the products is a gas, ΔS° must be positive, which helps make ΔG° negative.

1 point is earned for a valid justification.

The HCO_3^- ion has three carbon-to-oxygen bonds. Two of the carbon-to-oxygen bonds have the same length and the third carbon-to-oxygen bond is longer than the other two. The hydrogen atom is bonded to one of the oxygen atoms. In the box below, draw a Lewis electron-dot diagram (or diagrams) HCO_3^- ion that is (are) consistent with the given information.




See diagram above.


1 point is earned for a correct Lewis structure of HCO_3^- .

1 point is earned for indicating resonance (e.g., two diagrams, or diagram with an arrow between the two appropriate oxygen atoms)

A student prepares a solution containing equimolar amounts of $\text{HC}_2\text{H}_3\text{O}_2$ and $\text{NaC}_2\text{H}_3\text{O}_2$. The pH of the solution is measured to be 4.7. The student adds two drops of 3.0 M $\text{HNO}_3(aq)$ and stirs the sample, observing that the pH remains at 4.7. Write a balanced, net-ionic equation for the reaction between $\text{HNO}_3(aq)$ and the chemical species in the sample that is responsible for the pH remaining at 4.




OR




1 point is earned for a correct equation.

SCORING GUIDELINES

Question



To determine the molar mass of an unknown metal, M, a student reacts iodine with an excess of the metal to form the water-soluble compound MI_2 , as represented by the equation above. The reaction proceeds until all of the I_2 is consumed. The $\text{MI}_2(aq)$ solution is quantitatively collected and heated to remove the water, and the product is dried and weighed to constant mass. The experimental steps are represented below, followed by a data table.



Data for Unknown Metal Lab	
Mass of beaker	125.457 g
Mass of beaker + metal M	126.549 g
Mass of beaker + metal M + I_2	127.570 g
Mass of MI_2 , first weighing	1.284 g
Mass of MI_2 , second weighing	1.284 g

SCORING GUIDELINES

Question 3 (continued)

- (a) Given that the metal M is in excess, calculate the number of moles of
- I_2
- that reacted.

$127.570 - 126.549 = 1.021 \text{ g } I_2$ $1.021 \text{ g } I_2 \times \frac{1 \text{ mol } I_2}{253.80 \text{ g } I_2} = 0.004023 \text{ mol } I_2$	1 point is earned for the number of moles.
---	--

- (b) Calculate the molar mass of the unknown metal M.

Number of moles of I_2 = number of moles of M $1.284 \text{ g } MI_2 - 1.021 \text{ g } I_2 = 0.263 \text{ g M}$ $\text{Molar mass of M} = \frac{0.263 \text{ g M}}{0.004023 \text{ mol M}} = 65.4 \text{ g/mol}$	1 point is earned for the number of grams of M. 1 point is earned for the molar mass.
---	--

The student hypothesizes that the compound formed in the synthesis reaction is ionic.

- (c) Propose an experimental test the student could perform that could be used to support the hypothesis. Explain how the results of the test would support the hypothesis if the substance was ionic.

The student could dissolve the compound in water or melt the compound and see if the solution/melt conducts electricity. If the solution/melt conducts electricity, mobile ions capable of carrying charge must be present, thus the compound is likely to be ionic. OR The student could heat the compound until it melts or boils. If the melting/boiling point is very high, then the compound is likely to be ionic.	1 point is earned for an appropriate test. 1 point is earned for explaining how the results would support the hypothesis.
--	--

The student hypothesizes that Br_2 will react with metal M more vigorously than I_2 did because Br_2 is a liquid at room temperature.

SCORING GUIDELINES

Question 3 (continued)

- (d) Explain why
- I_2
- is a solid at room temperature whereas
- Br_2
- is a liquid. Your explanation should clearly reference the types and relative strengths of the intermolecular forces present in each substance.

Both Br_2 and I_2 molecules are nonpolar molecules, therefore the only possible intermolecular forces are London dispersion forces. The London dispersion forces are stronger in I_2 because it is larger in size with more electrons and/or a more polarizable electron cloud. The stronger London dispersion forces in I_2 result in a higher melting point, which makes I_2 a solid at room temperature.	1 point is earned for identifying the forces in each substance as London dispersion forces. 1 point is earned for explaining why the forces are stronger in I_2 than in Br_2 .
--	---

While cleaning up after the experiment, the student wishes to dispose of the unused solid I_2 in a responsible manner. The student decides to convert the solid I_2 to $I^-(aq)$ anion. The student has access to three solutions, $H_2O_2(aq)$, $Na_2S_2O_3(aq)$, and $Na_2S_4O_6(aq)$, and the standard reduction table shown below.

Half-reaction	E° (V)
$S_4O_6^{2-}(aq) + 2 e^- \rightarrow 2 S_2O_3^{2-}(aq)$	0.08
$I_2(s) + 2 e^- \rightarrow 2 I^-(aq)$	0.54
$O_2(g) + 2 H^+(aq) + 2 e^- \rightarrow H_2O_2(aq)$	0.68

- (e) Which solution should the student add to
- $I_2(s)$
- to reduce it to
- $I^-(aq)$
- ? Circle your answer below. Justify your answer and include a calculation of
- E°
- for the overall reaction.


$H_2O_2(aq)$	$Na_2S_2O_3(aq)$	$Na_2S_4O_6(aq)$
[Na $S_2O_3(aq)$ should be circled.] The reaction between $S_2O_3^{2-}(aq)$ and $I_2(s)$ will be thermodynamically favorable because E° for the reaction is positive ($E^\circ = 0.54 - 0.08 = +0.46 \text{ V}$), from which it follows that ΔG° is negative because $\Delta G^\circ = -nFE^\circ$.	1 point is earned for the correct choice. 1 point is earned for a correct justification.	

- (f) Write the balanced net-ionic equation for the reaction between
- I_2
- and the solution you selected in part (e).

$I_2 + 2 S_2O_3^{2-} \rightarrow 2 I^- + S_4O_6^{2-}$	1 point is earned for the correct equation.
---	---

SCORING GUIDELINES

Question



Phenol is a weak acid that partially dissociates in water, according to the equation above.

(a) What is the pH of a 0.75 M $\text{C}_6\text{H}_5\text{OH}(aq)$ solution?

$K_a = \frac{[\text{C}_6\text{H}_5\text{O}^-][\text{H}_3\text{O}^+]}{[\text{C}_6\text{H}_5\text{OH}]}$ $1.12 \times 10^{-10} = \frac{x^2}{(0.75 - x)} \quad \text{Assume that } x \ll 0.75.$ $x^2 = 8.4 \times 10^{-11}$ $x = \sqrt{8.4 \times 10^{-11}}$ $x = 9.2 \times 10^{-6} \text{ M}$ $\text{pH} = -\log[\text{H}^+] = -\log(9.2 \times 10^{-6}) = 5.04$	1 point is earned for a correct setup and calculation of $[\text{H}^+]$. 1 point is earned for the correct setup and calculation of pH based on a correct setup for the $[\text{H}^+]$ calculation.
---	---


(b) For a certain reaction involving $\text{C}_6\text{H}_5\text{OH}(aq)$ to proceed at a significant rate, the phenol must be primarily in its deprotonated form, $\text{C}_6\text{H}_5\text{O}^-(aq)$. In order to ensure that the $\text{C}_6\text{H}_5\text{OH}(aq)$ is deprotonated, the reaction must be conducted in a buffered solution. On the number scale below, circle each pH for which more than 50 percent of the phenol molecules are in the deprotonated form ($\text{C}_6\text{H}_5\text{O}^-(aq)$). Justify your answer.

Numbers 10 through 14 should be circled. When $\text{pH} > \text{p}K_a$, the deprotonated form will predominate. $\text{p}K_a = -\log(1.12 \times 10^{-10}) = 9.95$, therefore at pH 10 and above, $[\text{C}_6\text{H}_5\text{O}^-] > [\text{C}_6\text{H}_5\text{OH}]$.	1 point is earned for circling 10–14. 1 point is earned for the justification.
--	--




1 2 3 4 5 6 7 8 9 10 11 12 13 14

SCORING GUIDELINES

Question



At high temperatures the compound C_4H_6 (1,3-butadiene) reacts according to the equation above. The rate of the reaction was studied at 625 K in a rigid reaction vessel. Two different trials, each with a different starting concentration, were carried out. The data were plotted in three different ways, as shown below.



SCORING GUIDELINES

Question 5 (continued)

- (a) For trial 1, calculate the initial pressure, in atm, in the vessel at 625 K. Assume that initially all the gas present in the vessel is C_4H_6 .

For trial 1, $\frac{n}{V} = 0.020 \text{ mol/L}$ (or assume the volume of the vessel is 1.0 L; the number of moles of C_4H_6 in the vessel would then be 0.020 mol).	1 point is earned for a correct setup.
$PV = nRT$	
$P = \frac{nRT}{V} = \frac{(0.020 \text{ mol})(0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1})(625 \text{ K})}{1.0 \text{ L}} = 1.0 \text{ atm}$	1 point is earned for the correct answer.

- (b) Use the data plotted in the graphs to determine the order of the reaction with respect to C_4H_6 .

Second order (because the plot of $1/[C_4H_6]$ is a straight line).	1 point is earned for the correct order.
---	--

- (c) The initial rate of the reaction in trial 1 is 0.0010 mol/(L·s). Calculate the rate constant, k , for the reaction at 625 K.

From the second-order rate law (differential form): $\text{rate} = k[C_4H_6]^2$ $\Rightarrow k = \frac{\text{rate}}{([C_4H_6])^2} = \frac{0.0010 \text{ mol/(L} \cdot \text{s)}}{(0.020 \text{ mol/L})^2} = 2.5 \text{ L/(mol} \cdot \text{s)}$ OR From the second-order rate law (integrated form): $\frac{1}{[C_4H_6]_t} = 2kt + \frac{1}{[C_4H_6]_0}$ The coefficient of t is equal to $2k$ because of the reaction stoichiometry. The slope of the line in the plot of $\frac{1}{[C_4H_6]}$ versus time is $2k$. Thus slope = $5.0 \text{ L/(mol} \cdot \text{s)} = 2k$, therefore $k = 2.5 \text{ L/(mol} \cdot \text{s)}$. <u>Note:</u> Students who choose the second method of determining k but omit the factor of 2, thereby getting an answer of $5.0 \text{ L/(mol} \cdot \text{s)}$, still earn the point.	1 point is earned for the correct value.
---	--

SCORING GUIDELINES

Question



The polyatomic ion $C_{10}H_{12}N_2O_8^{4-}$ is commonly abbreviated as $EDTA^{4-}$. The ion can form complexes with metal ions in aqueous solutions. A complex of $EDTA^{4-}$ with Ba^{2+} ion forms according to the equation above. A 50.0 mL volume of a solution that has an $EDTA^{4-}(aq)$ concentration of 0.30 M is mixed with 50.0 mL of 0.20 M $Ba(NO_3)_2$ to produce 100.0 mL of solution.

- (a) Considering the value of K for the reaction, determine the concentration of $Ba(EDTA)^{2-}(aq)$ in the 100.0 mL of solution. Justify your answer.

Based on the K value, the reaction goes essentially to completion. $Ba^{2+}(aq)$ is the limiting reactant. The concentration of Ba^{2+} when the solutions are first mixed but before any reaction takes place is $0.20 \text{ M}/2 = 0.10 \text{ M}$. Thus the equilibrium concentration of $Ba(EDTA)^{2-}(aq)$ is 0.10 M.	1 point is earned for indicating that the equilibrium concentration of $Ba(EDTA)^{2-}(aq)$ is the same as the original concentration of Ba^{2+} when the solutions are mixed. 1 point is earned for the concentration with appropriate calculations.
--	---

- (b) The solution is diluted with distilled water to a total volume of 1.00 L. After equilibrium has been reestablished, is the number of moles of $Ba^{2+}(aq)$ present in the solution greater than, less than, or equal to the number of moles of $Ba^{2+}(aq)$ present in the original solution before it was diluted? Justify your answer.


The number of moles of $Ba^{2+}(aq)$ increases because the percent dissociation of $Ba(EDTA)^{2-}(aq)$ increases as the solution is diluted. OR A mathematical justification such as the following: The dilution from 100.0 mL to 1.00 L reduces the concentrations of all species to one tenth of their original values. Immediately after the dilution, the reaction quotient, Q , can be determined as shown below. $Q = \frac{\frac{1}{10}[Ba(EDTA)^{2-}]}{\frac{1}{10}[Ba^{2+}] \times \frac{1}{10}[EDTA^{4-}]} = 10K$ Because $Q > K$, the net reaction will produce more reactants to move toward equilibrium, so the number of moles of $Ba^{2+}(aq)$ will be greater than the number in the original solution.	1 point is earned for stating that the number of moles of $Ba^{2+}(aq)$ will increase. 1 point is earned for a valid justification.
---	--

SCORING GUIDELINES

Question

A student is given a 25.0 mL sample of a solution of an unknown monoprotic acid and asked to determine the concentration of the acid by titration. The student uses a standardized solution of 0.110 M NaOH(aq), a buret, a flask, an appropriate indicator, and other laboratory equipment necessary for the titration.

- (a) The images below show the buret before the titration begins (below left) and at the end point (below right). What should the student record as the volume of NaOH(aq) delivered to the flask?



$$37.30 \text{ mL} - 5.65 \text{ mL} = 31.65 \text{ mL}$$

1 point is earned for the correct volume.

- (b) Based on the given information and your answer to part (a), determine the value of the concentration of the acid that should be recorded in the student's lab report.

At the equivalence point, moles OH^- added = moles of H^+ consumed.
Because HA is monoprotic:

$$(0.110 \text{ M})(0.03165 \text{ L}) \times \frac{1 \text{ mol HA}}{1 \text{ mol NaOH}} \times \frac{1}{0.0250 \text{ L}} = 0.139 \text{ M}$$

OR

$$\text{moles of } \text{H}^+ \text{ consumed} = M_a V_a$$

$$M_a V_a = M_b V_b$$

$$\text{Therefore, } M_a = \frac{M_b V_b}{V_a} = \frac{(0.110 \text{ M})(0.03165 \text{ L})}{0.0250 \text{ L}} = 0.139 \text{ M}$$

1 point is earned for the correct setup and molarity.

- (c) In a second trial, the student accidentally added more NaOH(aq) to the flask than was needed to reach the end point, and then recorded the final volume. Would this error increase, decrease, or have no effect on the calculated acid concentration for the second trial? Justify your answer.

The error would increase the calculated acid concentration.

A volume of NaOH(aq) larger than the actual volume needed to reach the equivalence point, would lead to a calculation of moles of base that would be greater than the moles of acid actually present in the solution. The assumption that the moles of acid are the same as the moles of base would lead to a calculated concentration of acid that would be higher than the actual concentration.

1 point is earned for indicating an increase.

1 point is earned for a valid justification.

AP[®] Chemistry

2014 Scoring Guidelines

© 2014 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

4 SCORING GUIDELINES

Question 1
(10 points)

Mass of KI tablet	0.425 g
Mass of thoroughly dried filter paper	1.462 g
Mass of filter paper + precipitate after first drying	1.775 g
Mass of filter paper + precipitate after second drying	1.699 g
Mass of filter paper + precipitate after third drying	1.698 g

A student is given the task of determining the I^- content of tablets that contain KI and an inert, water-soluble sugar as a filler. A tablet is dissolved in 50.0 mL of distilled water, and an excess of 0.20 M $Pb(NO_3)_2(aq)$ is added to the solution. A yellow precipitate forms, which is then filtered, washed, and dried. The data from the experiment are shown in the table above.

(a) For the chemical reaction that occurs when the precipitate forms,

(i) write a balanced, net-ionic equation for the reaction, and

$Pb^{2+} + 2 I^- \rightarrow PbI_2$	1 point is earned for a balanced net-ionic equation.
-------------------------------------	--

(ii) explain why the reaction is best represented by a net-ionic equation.

The net-ionic equation shows the formation of the $PbI_2(s)$ from $Pb^{2+}(aq)$ and $I^-(aq)$ ions, omitting the non-reacting species (spectator ions), $K^+(aq)$ and $NO_3^-(aq)$.	1 point is earned for a valid explanation.
--	--

(b) Explain the purpose of drying and weighing the filter paper with the precipitate three times.

The filter paper and precipitate must be dried several times (to a constant mass) to ensure that all the water has been driven off.	1 point is earned for a valid explanation.
---	--

(c) In the filtrate solution, is $[K^+]$ greater than, less than, or equal to $[NO_3^-]$? Justify your answer.

$[K^+]$ is less than $[NO_3^-]$ because the source of the NO_3^- , the 0.20 M $Pb(NO_3)_2(aq)$, was added in excess.	1 point is earned for a correct comparison with a valid explanation.
---	--

4 SCORING GUIDELINES

Question 1 (continued)

(d) Calculate the number of moles of precipitate that is produced in the experiment.

$1.698 \text{ g} - 1.462 \text{ g} = 0.236 \text{ g } PbI_2(s)$ $0.236 \text{ g } PbI_2 \times \frac{1 \text{ mol } PbI_2}{461.0 \text{ g } PbI_2} = 5.12 \times 10^{-4} \text{ mol } PbI_2$	1 point is earned for the correct number of moles of $PbI_2(s)$ precipitate.
---	--

(e) Calculate the mass percent of I^- in the tablet.

$5.12 \times 10^{-4} \text{ mol } PbI_2 \times \frac{2 \text{ mol } I^-}{1 \text{ mol } PbI_2} = 1.02 \times 10^{-3} \text{ mol } I^-$ $1.02 \times 10^{-3} \text{ mol } I^- \times \frac{126.91 \text{ g } I^-}{1 \text{ mol } I^-} = 0.130 \text{ g } I^- \text{ in one tablet}$ $\frac{0.130 \text{ g } I^-}{0.425 \text{ g KI tablet}} = 0.306 = 30.6\% I^- \text{ per KI tablet}$	1 point is earned for determining the number of moles of I^- in one tablet. 1 point is earned for calculating the mass percent of I^- in the KI tablet.
--	---

(f) In another trial, the student dissolves a tablet in 55.0 mL of water instead of 50.0 mL of water. Predict whether the experimentally determined mass percent of I^- will be greater than, less than, or equal to the amount calculated in part (e). Justify your answer.

The mass percent of I^- will be the same. $Pb^{2+}(aq)$ was added in excess, ensuring that essentially no I^- remained in solution. The additional water is removed by filtration and drying, leaving the same mass of dried precipitate.	1 point is earned for correct comparison with a valid justification.
---	--

(g) A student in another lab also wants to determine the I^- content of a KI tablet but does not have access to $Pb(NO_3)_2$. However, the student does have access to 0.20 M $AgNO_3$, which reacts with $I^-(aq)$ to produce $AgI(s)$. The value of K_{sp} for AgI is 8.5×10^{-17} .

(i) Will the substitution of $AgNO_3$ for $Pb(NO_3)_2$ result in the precipitation of the I^- ion from solution? Justify your answer.

Yes. Addition of an excess of 0.20 M $AgNO_3(aq)$ will precipitate all of the I^- ion present in the solution because AgI is insoluble, as evidenced by its low value of K_{sp} .	1 point is earned for the correct answer with a valid justification.
---	--

(ii) The student only has access to one KI tablet and a balance that can measure to the nearest 0.01 g. Will the student be able to determine the mass of AgI produced to three significant figures? Justify your answer.


4 SCORING GUIDELINES

Question 1 (continued)

No. If masses can be measured to ± 0.01 g, then the mass of the dry AgI(s) precipitate (which is less than 1 g) will be known to only two significant figures.


1 point is earned for a correct answer with a valid justification.

2014 SCORING GUIDELINES

Question
(10 points)

Propanoic acid, $\text{CH}_3\text{CH}_2\text{COOH}$, is a carboxylic acid that reacts with water according to the equation above. At 25°C the pH of a 50.0 mL sample of 0.20 M $\text{CH}_3\text{CH}_2\text{COOH}$ is 2.79.

- (a) Identify a Brønsted-Lowry conjugate acid-base pair in the reaction. Clearly label which is the acid and which is the base.



1 point is earned for writing (or naming) either of the Brønsted-Lowry conjugate acid-base pairs with a clear indication of which is the acid and which is the base.

- (b) Determine the value of K_a for propanoic acid at 25°C .

$$[\text{H}_3\text{O}^+] = 10^{-\text{pH}} = 10^{-2.79} = 1.6 \times 10^{-3} M$$

$$[\text{CH}_3\text{CH}_2\text{COO}^-] = [\text{H}_3\text{O}^+]$$

AND

$$[\text{CH}_3\text{CH}_2\text{COOH}] = 0.20 M - [\text{H}_3\text{O}^+], \text{ OR } [\text{CH}_3\text{CH}_2\text{COOH}] \approx 0.20 M$$

(state or assume that $[\text{H}_3\text{O}^+] \ll 0.20 M$)

$$K_a = \frac{[\text{CH}_3\text{CH}_2\text{COO}^-][\text{H}_3\text{O}^+]}{[\text{CH}_3\text{CH}_2\text{COOH}]} = \frac{(1.6 \times 10^{-3} M)^2}{0.20 M} = 1.3 \times 10^{-5}$$

1 point is earned for correctly solving for $[\text{H}_3\text{O}^+]$.

1 point is earned for the K_a expression for propanoic acid
OR

1 point is earned for substituting values into the K_a expression.

1 point is earned for correctly solving for the value of K_a .

- (c) For each of the following statements, determine whether the statement is true or false. In each case, explain the reasoning that supports your answer.

- (i) The pH of a solution prepared by mixing the 50.0 mL sample of 0.20 M $\text{CH}_3\text{CH}_2\text{COOH}$ with a 50.0 mL sample of 0.20 M NaOH is 7.00.

False. The conjugate base of a weak acid undergoes hydrolysis (see equation below) at equivalence to form a solution with a pH > 7 .



1 point is earned for noting that the statement is false AND providing a supporting explanation.

2014 SCORING GUIDELINES

Question 2 (continued)

- (ii) If the pH of a hydrochloric acid solution is the same as the pH of a propanoic acid solution, then the molar concentration of the hydrochloric acid solution must be less than the molar concentration of the propanoic acid solution.

True. HCl is a strong acid that ionizes completely. Fewer moles of HCl are needed to produce the same $[\text{H}_3\text{O}^+]$ as the propanoic acid solution, which only partially ionizes.

1 point is earned for noting that the statement is true and providing a supporting explanation.

A student is given the task of determining the concentration of a propanoic acid solution of unknown concentration. A 0.173 M NaOH solution is available to use as the titrant. The student uses a 25.00 mL volumetric pipet to deliver the propanoic acid solution to a clean, dry flask. After adding an appropriate indicator to the flask, the student titrates the solution with the 0.173 M NaOH, reaching the end point after 20.52 mL of the base solution has been added.

- (d) Calculate the molarity of the propanoic acid solution.

$$\begin{aligned} \text{Let } x &= \text{moles of propanoic acid} \\ \text{then } x &= (0.02052 \text{ L NaOH}) \times \frac{0.173 \text{ mol NaOH}}{1 \text{ L NaOH}} \times \frac{1 \text{ mol acid}}{1 \text{ mol NaOH}} \\ &= 3.55 \times 10^{-3} \text{ mol propanoic acid} \\ \frac{3.55 \times 10^{-3} \text{ mol acid}}{0.02500 \text{ L acid}} &= 0.142 \text{ M} \end{aligned}$$

OR

Since $\text{CH}_3\text{CH}_2\text{COOH}$ is monoprotic and, at the equivalence point, moles H^+ = moles OH^- , then

$$\begin{aligned} M_A V_A &= M_B V_B \\ M_A &= \frac{M_B V_B}{V_A} = \frac{(0.173 \text{ M NaOH})(20.52 \text{ mL NaOH})}{25.00 \text{ mL acid}} = 0.142 \text{ M} \end{aligned}$$

1 point is earned for correctly calculating the number of moles of acid that reacted at the equivalence point.

1 point is earned for the correct molarity of acid.

- (e) The student is asked to redesign the experiment to determine the concentration of a butanoic acid solution instead of a propanoic acid solution. For butanoic acid the value of $\text{p}K_a$ is 4.83. The student claims that a different indicator will be required to determine the equivalence point of the titration accurately. Based on your response to part (b), do you agree with the student's claim? Justify your answer.

2014 SCORING GUIDELINES

Question 2 (continued)


Disagree with the student's claim

From part (b) above, $\text{p}K_a$ for propanoic acid is $\log(1.3 \times 10^{-5}) = 4.89$. Because 4.83 is so close to 4.89, the pH at the equivalence point in the titration of butanoic acid should be close enough to the pH in the titration of propanoic acid to make the original indicator appropriate for the titration of butanoic acid.

1 point is earned for disagreeing with the student's claim and making a valid justification using $\text{p}K_a$, K_a , or pH arguments.

1 point is earned for numerically comparing either: the two $\text{p}K_a$ values, the two K_a values, or the two pH values at the equivalence point.

SCORING GUIDELINES

Question
(10 points)

A student is given a standard galvanic cell, represented above, that has a Cu electrode and a Sn electrode. As current flows through the cell, the student determines that the Cu electrode increases in mass and the Sn electrode decreases in mass.

- (a) Identify the electrode at which oxidation is occurring. Explain your reasoning based on the student's observations.

Since the Sn electrode is losing mass, Sn atoms must be forming $\text{Sn}^{2+}(aq)$. This process is oxidation.

OR

because the cell operates, E° must be positive and, based on the E° values from the table, it must be Sn that is oxidized.


1 point is earned for the correct answer with justification.

- (b) As the mass of the Sn electrode decreases, where does the mass go?

The atoms on the Sn electrode are going into the solution as Sn^{2+} ions.

1 point is earned for the correct answer.

- (c) In the expanded view of the center portion of the salt bridge shown in the diagram below, draw and label a particle view of what occurs in the salt bridge as the cell begins to operate. Omit solvent molecules and use arrows to show the movement of particles.



The response should show at least one K^+ ion moving toward the Cu compartment on the left and at least one NO_3^- ion moving in the opposite direction.

1 point is earned for correct representation of both K^+ and NO_3^- ions. (Including free electrons loses this point.)

1 point is earned for correctly indicating the direction of movement of both ions.

SCORING GUIDELINES

Question 3 (continued)

- (d) A nonstandard cell is made by replacing the 1.0 M solutions of $\text{Cu}(\text{NO}_3)_2$ and $\text{Sn}(\text{NO}_3)_2$ in the standard cell with 0.50 M solutions of $\text{Cu}(\text{NO}_3)_2$ and $\text{Sn}(\text{NO}_3)_2$. The volumes of solutions in the nonstandard cell are identical to those in the standard cell.

- (i) Is the cell potential of the nonstandard cell greater than, less than, or equal to the cell potential of the standard cell? Justify your answer.

It is the same. In the cell reaction $Q = \frac{[\text{Sn}^{2+}]}{[\text{Cu}^{2+}]}$, and the concentrations of Sn^{2+} and Cu^{2+} are equal to each other in both cases.

1 point is earned for the correct answer with justification.

- (ii) Both the standard and nonstandard cells can be used to power an electronic device. Would the nonstandard cell power the device for the same time, a longer time, or a shorter time as compared with the standard cell? Justify your answer.

The nonstandard cell would power the device for a shorter time because the supply of Cu^{2+} ions will be exhausted more quickly.

OR

The nonstandard cell would power the device for a shorter time because the reaction will reach $E = 0$ more quickly.

1 point is earned for the correct answer with justification.

- (e) In another experiment, the student places a new Sn electrode into a fresh solution of 1.0 M $\text{Cu}(\text{NO}_3)_2$.

Half-Reaction	E° (V)
$\text{Cu}^+ + e^- \rightarrow \text{Cu}(s)$	0.52
$\text{Cu}^{2+} + 2e^- \rightarrow \text{Cu}(s)$	0.34
$\text{Sn}^{4+} + 2e^- \rightarrow \text{Sn}^{2+}$	0.15
$\text{Sn}^{2+} + 2e^- \rightarrow \text{Sn}(s)$	-0.14

- (i) Using information from the table above, write a net-ionic equation for the reaction between the Sn electrode and the $\text{Cu}(\text{NO}_3)_2$ solution that would be thermodynamically favorable. Justify that the reaction is thermodynamically favorable.

SCORING GUIDELINES


Question 3 (continued)

$\text{Cu}^{2+}(\text{aq}) + \text{Sn}(\text{s}) \rightarrow \text{Cu}(\text{s}) + \text{Sn}^{2+}(\text{aq})$ E° is positive ($0.34 \text{ V} + 0.14 \text{ V} = 0.48 \text{ V}$), therefore the reaction is thermodynamically favorable. OR The cell observations from earlier parts of the question are evidence that the Sn is oxidized and Cu is reduced, therefore E° must be positive.	1 point is earned for the correct net-ionic equation. 1 point is earned for a correct justification (unit not needed in calculation).
--	---


(ii) Calculate the value of ΔG° for the reaction. Include units with your answer.

$\Delta G^\circ = -nFE^\circ$ $\Delta G^\circ = -\frac{2 \text{ mol } e^-}{\text{mol}_{\text{rxn}}} \times \frac{96,485 \text{ C}}{\text{mol } e^-} \times \frac{0.48 \text{ J}}{\text{C}} = -93,000 \text{ J/mol}_{\text{rxn}} = -93 \text{ kJ/mol}_{\text{rxn}}$	1 point is earned for the correct number of electrons. 1 point is earned for the correct answer with unit.
--	--

SCORING GUIDELINES

Question
(4 points)

When heated, calcium carbonate decomposes according to the equation above. In a study of the decomposition of calcium carbonate, a student added a 50.0 g sample of powdered $\text{CaCO}_3(\text{s})$ to a 1.00 L rigid container. The student sealed the container, pumped out all the gases, then heated the container in an oven at 1100 K. As the container was heated, the total pressure of the $\text{CO}_2(\text{g})$ in the container was measured over time. The data are plotted in the graph below.



The student repeated the experiment, but this time the student used a 100.0 g sample of powdered $\text{CaCO}_3(\text{s})$. In this experiment, the final pressure in the container was 1.04 atm, which was the same final pressure as in the first experiment.

(a) Calculate the number of moles of $\text{CO}_2(\text{g})$ present in the container after 20 minutes of heating.

$PV = nRT$ $\frac{PV}{RT} = n = \frac{(1.04 \text{ atm})(1.00 \text{ L})}{(0.0821 \frac{\text{L atm}}{\text{mol K}})(1100 \text{ K})} = 0.0115 \text{ mol CO}_2$	1 point is earned for the proper setup using the ideal gas law and an answer that is consistent with the setup.
--	--

SCORING GUIDELINES

Question continued)

- (b) The student claimed that the final pressure in the container in each experiment became constant because all of the $\text{CaCO}_3(s)$ had decomposed. Based on the data in the experiments, do you agree with this claim? Explain.

Do not agree with claim Explanation I: In experiment 1, the moles of $\text{CaCO}_3 = 50.0 \text{ g}/100.09 \text{ g/mol} = 0.500 \text{ mol } \text{CaCO}_3$. If the reaction had gone to completion, 0.500 mol of CO_2 would have been produced. From part (a) only 0.0115 mol was produced. Hence, the student's claim was false. Explanation II: The two different experiments (one with 50.0 g of CaCO_3 and one with 100.0 g of CaCO_3) reached the same constant, final pressure of 1.04 atm. Since increasing the amount of reactant did not produce more product, there is no way that all of the CaCO_3 reacted. Instead, an equilibrium condition has been achieved and there must be some solid CaCO_3 in the container.	1 point is earned for disagreement with the claim <u>and</u> for a correct justification using stoichiometry or a discussion of the creation of an equilibrium condition.
--	--

- (c) After 20 minutes some $\text{CO}_2(g)$ was injected into the container, initially raising the pressure to 1.5 atm. Would the final pressure inside the container be less than, greater than, or equal to 1.04 atm? Explain your reasoning.

The final pressure would be equal to 1.04 atm. Equilibrium was reached in both experiments; the equilibrium pressure at this temperature is 1.04 atm. As the reaction shifts toward the reactant, the amount of $\text{CO}_2(g)$ in the container will decrease until the pressure returns to 1.04 atm.	1 point is earned for the correct answer with justification.
---	---

- (d) Are there sufficient data obtained in the experiments to determine the value of the equilibrium constant, K_p , for the decomposition of $\text{CaCO}_3(s)$ at 1100 K? Justify your answer.

SCORING GUIDELINES

Question continued)

Yes. For the equilibrium reaction represented by the chemical equation in this problem, at a given temperature the equilibrium pressure of CO_2 determines the equilibrium constant. Since the measured pressure of CO_2 is also the equilibrium pressure of CO_2 , $K_p = P_{\text{CO}_2} = 1.04$.

Note: If the response in part (b) indicates “yes”, that all of the $\text{CaCO}_3(s)$ had decomposed, then the point can be earned by stating that the system did not reach equilibrium in either experiment and hence the value of K_p cannot be calculated from the data.

1 point is earned for correct explanation that is consistent with the student's answer to part (b).

SCORING GUIDELINES

Question 5
(4 points)


Nonmetal	C	N	O	Ne	Si	P	S	Ar
Formula of Compound	CF ₄	NF ₃	OF ₂	No compound	SiF ₄	PF ₃	SF ₂	No compound

Some binary compounds that form between fluorine and various nonmetals are listed in the table above. A student examines the data in the table and poses the following hypothesis: the number of F atoms that will bond to a nonmetal is always equal to 8 minus the number of valence electrons in the nonmetal atom.

- (a) Based on the student's hypothesis, what should be the formula of the compound that forms between chlorine and fluorine?

ClF	1 point is earned for the correct formula.
-----	--

- (b) In an attempt to verify the hypothesis, the student researches the fluoride compounds of the other halogens and finds the formula ClF₃. In the box below, draw a complete Lewis electron-dot diagram for a molecule of ClF₃.



See diagram above.	1 point is earned for a central Cl atom surrounded by three bonding pairs with F atoms and two nonbonding (lone) pairs of electrons. F atoms must have three nonbonding pairs each. Electron pairs can be depicted as dots or line segments.
--------------------	--

SCORING GUIDELINES

Question 5 (continued)

- (c) Two possible geometric shapes for the ClF₃ molecule are trigonal planar and T-shaped. The student does some research and learns that the molecule has a dipole moment. Which of the two shapes is consistent with the fact that the ClF₃ molecule has a dipole moment? Justify your answer in terms of bond polarity and molecular structure.

The molecule is T-shaped because a T-shaped structure is asymmetric with dipoles that do not cancel out, but produce a net dipole (i.e., a polar molecule). OR because, if the molecule had a trigonal planar structure, the molecule would be symmetric with dipoles that cancel out and produce a net dipole of zero (i.e., a nonpolar molecule), which is not consistent with the observation that the ClF₃ molecule does have a dipole moment.	1 point is earned for indicating that the molecule is T-shaped with an acceptable explanation.
---	--

In an attempt to resolve the existence of the ClF₃ molecule with the hypothesis stated above, the student researches the compounds that form between halogens and fluorine, and assembles the following list.

Halogen	Formula(s)
F	F ₂
Cl	
Br	BrF, BrF ₃ , BrF ₅
I	IF, IF ₃ , IF ₅ , IF ₇


- (d) Based on concepts of atomic structure and periodicity, propose a modification to the student's previous hypothesis to account for the compounds that form between halogens and fluorine.

An acceptable hypothesis (descriptive or formulaic) must include the following ideas: 1. Atomic Structure: e.g., odd number of F atoms 2. Periodicity: e.g., as the atomic number of the central halogen atom increases, the number of F atoms increases.	1 point is earned for an acceptably modified hypothesis that addresses both atomic structure and periodicity.
---	---

SCORING GUIDELINES

Question
(4 points)

A student places a mixture of plastic beads consisting of polypropylene (PP) and polyvinyl chloride (PVC) in a 1.0 L beaker containing distilled water. After stirring the contents of the beaker vigorously, the student observes that the beads of one type of plastic sink to the bottom of the beaker and the beads of the other type of plastic float on the water. The chemical structures of PP and PVC are represented by the diagrams below, which show segments of each polymer.





- (a) Given that the spacing between polymer chains in PP and PVC is similar, the beads that sink are made of which polymer? Explain.

The PVC beads sink. The spacing between chains is similar, but a Cl atom has a greater mass than CH₃.

1 point is earned for the correct polymer with a correct explanation.

PP is synthesized from propene, C₃H₆, and PVC is synthesized from vinyl chloride, C₂H₃Cl. The structures of the molecules are shown below.



- (b) The boiling point of liquid propene (226 K) is lower than the boiling point of liquid vinyl chloride (260 K). Account for this difference in terms of the types and strengths of intermolecular forces present in each liquid.


Both substances have dipole-dipole interactions and London dispersion forces (or propene is essentially nonpolar with only LDFs while vinyl chloride has both LDFs and dipole-dipole forces). Propene contains a CH₃ group, but vinyl chloride contains a Cl atom. Vinyl chloride thus has a larger electron cloud, is more polarizable, and has a larger dipole moment. Thus intermolecular attractions are stronger in vinyl chloride, which results in it having the higher boiling point.

1 point is earned for a discussion of intermolecular forces and for a comparison of their relative strengths.

SCORING GUIDELINES

Question 6 (continued)

In a separate experiment, the student measures the enthalpies of combustion of propene and vinyl chloride. The student determines that the combustion of 2.00 mol of vinyl chloride releases 2300 kJ of energy, according to the equation below.



- (c) Using the table of standard enthalpies of formation below, determine whether the combustion of 2.00 mol of propene releases more, less, or the same amount of energy that 2.00 mol of vinyl chloride releases. Justify your answer with a calculation. The balanced equation for the combustion of 2.00 mol of propene is $2 \text{C}_3\text{H}_6(g) + 9 \text{O}_2(g) \rightarrow 6 \text{CO}_2(g) + 6 \text{H}_2\text{O}(g)$.

Substance	C ₂ H ₃ Cl(g)	C ₃ H ₆ (g)	CO ₂ (g)	H ₂ O(g)	HCl(g)	O ₂ (g)
Standard Enthalpy of Formation (kJ/mol)	37	21	-394	-242	-92	0


$$\Delta H^\circ = 6(-394) + 6(-242) - 2(21) = -3858 \text{ kJ/mol}_{\text{rxn}}$$

The combustion of 2.00 mol of propene releases more energy.

1 point is earned for the calculation of the enthalpy of combustion of propene.

1 point is earned for the comparison of propene to vinyl chloride that is consistent with the calculated value.

SCORING GUIDELINES

Question 7
(4 points)

The half-life ($t_{1/2}$) of the catalyzed isomerization of *cis*-2-butene gas to produce *trans*-2-butene gas, represented above, was measured under various conditions, as shown in the table below.

Trial Number	Initial $P_{\text{cis-2-butene}}$ (torr)	V (L)	T (K)	$t_{1/2}$ (s)
1	300.	2.00	350.	100.
2	600.	2.00	350.	100.
3	300.	4.00	350.	100.
4	300.	2.00	365	50.

(a) The reaction is first order. Explain how the data in the table are consistent with a first-order reaction.

For a first-order reaction, the half-life is independent of reactant concentration (or pressure) at constant T , as shown in trials 1, 2, and 3.

1 point is earned for a correct explanation.

(b) Calculate the rate constant, k , for the reaction at 350. K. Include appropriate units with your answer.

$$k = \frac{0.693}{t_{1/2}} = \frac{0.693}{100. \text{ s}} = 0.00693 \text{ s}^{-1}$$

1 point is earned for correct numerical answer with units.

(c) Is the initial rate of the reaction in trial 1 greater than, less than, or equal to the initial rate in trial 2? Justify your answer.

The initial rate in trial 1 is less than that in trial 2 because $\text{rate} = k[\text{cis-2-butene}]$ or $\text{rate} = kP_{\text{cis-2-butene}}$ (with reference to values from both trials).

OR

because the initial concentration of *cis*-2-butene in trial 1 is less than that in trial 2 and k is constant.

1 point is earned for the correct answer with justification.

(d) The half-life of the reaction in trial 4 is less than the half-life in trial 1. Explain why, in terms of activation energy.

The temperature is higher in trial 4, meaning that the KE_{avg} of the molecules is greater. Consequently, in this trial a greater fraction of collisions have sufficient energy to overcome the activation energy barrier, thus the rate is greater.

1 point is earned for a correct answer with justification.