Chemical Reactions

AP Chemistry ▪ Unit 4

AP Chemistry



What we will cover



1 Recognizing

Recognizing a chemical reaction

A **chemical reaction** 化学反应 rearranges atoms: old **bonds** 化学键 break, new ones form – atoms are conserved.

> A **chemical change** 化学变化 makes new substances; a **physical change** 物理变化 only moves molecules apart.

Evidence of reaction: heat, light, gas, solid

2 Net ionic

Net ionic equations

A **precipitation reaction** 沉淀反应 makes an insoluble solid. Strip out the **spectator ions** 旁观离子 that do not change:

Silver + chloride

molecular: $AgNO_3 + NaCl \rightarrow AgCl\downarrow + NaNO_3$

$$Ag^+(aq) + Cl^-(aq) \rightarrow AgCl(s)$$

Use **solubility rules** 溶解度规则 to see which product is insoluble.



AgCl solid

3 Stoichiometry

Stoichiometry

Use the **mole ratio** 摩尔比 from the balanced equation: moles → ratio → moles.

Worked example

$8.0\,\mathrm{g\ H_2}$ burns $(2\mathrm{H_2} + \mathrm{O_2} \rightarrow 2\mathrm{H_2O})$:

$$n = \tfrac{8.0}{2.02} = 3.96\,\mathrm{mol} \Rightarrow 71\,\mathrm{g\ H_2O}$$

The **limiting reactant** 限量反应物 runs out first and caps the product.

Percent yield

$$\frac{\text{actual}}{\text{theoretical}} \times 100\%$$

4 Titration

Titration

A **titration** 滴定 finds an unknown concentration. Add **standard solution** 标准溶液 until the reaction is just complete – the **equivalence point** 等当点 (an **indicator** 指示剂 shows it).

Worked example

$25\,\mathrm{mL}$ of $0.100\,M$ NaOH neutralizes $20\,\mathrm{mL}$ HCl:

$$n = 0.00250 \Rightarrow c_{\mathrm{HCl}} = 0.125\,M$$



Titration finds concentration by volume

5 Types

Types of chemical reactions

Knowing the family helps you predict the products:

- **Synthesis** 化合: $\mathrm{A} + \mathrm{B} \rightarrow \mathrm{AB}$
- **Decomposition** 分解: $\mathrm{AB} \rightarrow \mathrm{A} + \mathrm{B}$
- **Precipitation**: an insoluble solid forms
- **Acid-base**: transfers a **proton** 质子 $\mathrm{H^+}$
- **Redox**: transfers **electrons** 电子

6 Acid-base

Acid-base reactions

Bronsted-Lowry 布朗斯特-劳里: an **acid** 酸 donates a proton, a **base** 碱 accepts one.

$$\underset{\text{acid}}{\mathrm{HCl}} + \underset{\text{base}}{\mathrm{H_2O}} \rightarrow \underset{\text{conj. acid}}{\mathrm{H_3O^+}} + \underset{\text{conj. base}}{\mathrm{Cl^-}}$$

> **Conjugate pairs** 共轭酸碱对 differ by one $\mathrm{H^+}$.

- **strong** 强 – ionizes almost completely (HCl, NaOH)
- **weak** 弱 – ionizes only slightly (vinegar)

Acid + base → salt + water (**neutralization** 中和).

7 Redox

Oxidation-reduction

A **redox** 氧化还原反应 transfers electrons, tracked by **oxidation numbers** 氧化数:

OIL RIG

Oxidation Is Loss, Reduction Is Gain

The **oxidizing agent** 氧化剂 is itself reduced; the **reducing agent** 还原剂 is oxidized.



8 Recap

Unit 4 – key takeaways

1. **Net ionic** — drop spectator ions; keep only what changes
2. **Stoichiometry** — mole ratio; limiting reactant caps the yield
3. **Acid-base** — Bronsted-Lowry proton transfer; conjugate pairs
4. **Redox** — OIL RIG electron transfer; oxidation numbers change

Check yourself

1. What are the ions removed from a net ionic equation called?
2. Which reactant caps how much product can form?
3. In Bronsted-Lowry terms, what does an acid do?
4. $Zn \rightarrow Zn^{2+}$: is zinc oxidized or reduced?



Answers **1.** spectator ions **2.** the limiting reactant **3.** donates a proton **4.** oxidized